Surname			Other	Names			
Centre Number				Candidate Number			
Candidate Signatur	re						

I eave blank

General Certificate of Education January 2004 Advanced Subsidiary Examination



CHEMISTRY CHM1 **Atomic Structure, Bonding and Periodicity** Unit 1

Friday 9 January 2004 Morning Session

In addition to this paper you will require: a calculator.

Time allowed: 1 hour

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

Information

- The maximum mark for this paper is 60.
- Mark allocations are shown in brackets.
- This paper carries 30 per cent of the total marks for AS. For Advanced Level this paper carries 15 per cent of the total marks.
- You are expected to use a calculator where appropriate.
- The following data may be required. Gas constant $R = 8.31 \,\mathrm{J \, K^{-1} \, mol^{-1}}$
- Your answers to the question in Section B should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

Advice

• You are advised to spend about 45 minutes on **Section A** and about 15 minutes on **Section B**.

	For Exam	iner's Use)
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
Total (Column	1)	→	
Total (Column	2)	→	
TOTAL			
Examine	r's Initials		

SECTION A

Answer all questions in the spaces provided.

(a)	One	isotope of sodium has a relative mass of 23.
	(i)	Define, in terms of the fundamental particles present, the meaning of the term isotopes.
	(ii)	Explain why isotopes of the same element have the same chemical properties.
	(iii)	Calculate the mass, in grams, of a single atom of this isotope of sodium. (The Avogadro constant, L , is $6.023 \times 10^{23} \text{ mol}^{-1}$)
		(5 marks
(b)	Give	the electronic configuration, showing all sub-levels, for a sodium atom.
(c)	Expl	(1 mark) ain why chromium is placed in the d block in the Periodic Table.
		(1 mark
(d)	an at	atom has half as many protons as an atom of ²⁸ Si and also has six fewer neutrons that com of ²⁸ Si. Give the symbol, including the mass number and the atomic number, of atom.



The Periodic Table of the Elements

The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

						3							
0	4.0 He Helium 2	20.2 Ne	Neon 10	39.9 Ar	Argon 18	83.8 Kr	Krypton 36	131.3 Xe	Xenon 54	222.0 Rn	Radon 86		
=		^о .	luorine	تې	hlorine	_ම ක	romine	6.9 —	lodine	0:0 A	statine		
>		16.0 O	Oxygen 8	32.1 S	Sulphur 16	79.0 Se	Selenium 34	127.6 Te	Tellurium 52	210.0 Po	Polonium 84		
>		14.0 N	Nitrogen 7	31.0 P	Phosphorus 15	74.9 As	Arsenic 33	121.8 Sb	Antimony 51	209.0 Bi	Bismuth 83		
≥		ا2.0 د	Carbon 6	28.1 Si	Silicon 14	72.6 Ge	Germanium 32	118.7 Sn	Tin 50	207.2 Pb	Lead 82		
≡		10.8 B	Boron Carbon Nitrogen Oxygen F 5 6 7 8 9	27.0 AI	Aluminium 13	69.7 Ga	Gallium 31	114.8 In	Indium 49	204.4 T	Thallium 81		
						65.4 Zn	Zinc 30	112.4 Cd	Cadmium 48	200.6 Hg	Mercury 80		
						58.7 Ni	Nickel 28	106.4 Pd	Palladium 46	195.1 Pt	Platinum 78		
						58.9 C	Cobalt 27	102.9 Rh	Rhodium 45	192.2 Ir	Iridium 77		
						55.8 Fe	Iron 26	101.1 Ru	Ruthenium 44	190.2 Os	Osmium 76		
		6.9 Li	Lithium 3			54.9 Mn	Manganese 25	98.9 Tc	umMolybdenumTechnetiumRutheniumRhodiumPalladiumSilver424344454647	186.2 Re	Rhenium 75		
		ssr				52.0 Ç	Chromium 24	95.9 Mo	Molybdenum 42	183.9 W	Tungsten 74		
		relative atomic mass	umber —			50.9 V	Vanadium 23	92.9 Nb	Niobium 41	180.9 Ta	Tantalum 73		
	Key	relative a	atomic number			47.9 Ti	Titanium 22	91.2 Zr	_	178.5 H	Hafnium 72		
						45.0 Sc		88.9 ×		138.9 La	≅ *	227 Ac	Actinium 89 †
=		9.0 Be	Beryllium 4	24.3 Mg		40.1 Ca	Calcium 20	87.6 Sr	_	137.3 Ba		226.0 Ra	Radium 88
-	1.0 H Hydrogen	6.9 Li	Lithium 3	23.0 Na	Sodium 11	39.1 X	_	85.5 Rb	_	132.9 Cs	_	223.0 Fr	Francium 87

175.0	_ ~	(260)	Lawrencium
Lu		Lr	103
, Yb Yb	Ytterbium 7	(259) No	Nobelium 1
168.9	Thulium	(258) (259) (3	Mendelevium
Tm	69	Md No	101
167.3 Er	ium	_E	mium
164.9	Holmium	(252)	Einsteinium
H3	67	Es	99
162.5	Dysprosium	247.1 252.1 (252) (257) (188	Californium
Dy	66		98
158.9	Terbium	247.1	Berkelium
Tb	65	Bk	97
157.3	dolinium	247.1	Curium
Gd		Cm	96
152.0	Europium	243.1	Americium
Eu	63	Am	95
150.4	Samarium	239.1	Plutonium
Sm	62	Pu	94
144.9	lum Promethium Samarium Europium Gadolinium Terbium Dysprosium Holmium Ert 61 62 63 64 65 66 67 68	237.0	Neptunium
Pm		Np	93
144.2	Neodymium	238.0	Uranium
Nd	60	U	92
140.1 140.9 144.2 144.9 150.4 152.0 155 Ce Pr Nd Pm Sm Eu	Praseodymium 59	232.0 231.0 238.0 237.0 Th Pa U Np	Thorium Protactinium Uranium 0
140.1	Cerium	232.0	Thorium
Ce	58	Th	90

† 90 – 103 Actinides

* 58 - 71 Lanthanides

Table 1 Proton n.m.r chemical shift data

Type of proton	δ/ppm
RCH ₃	0.7–1.2
R_2CH_2	1.2–1.4
R_3 CH	1.4–1.6
RCOCH ₃	2.1–2.6
$ROCH_3$	3.1–3.9
$RCOOCH_3$	3.7–4.1
ROH	0.5-5.0

Table 2 Infra-red absorption data

Bond	Wavenumber/cm ⁻¹
С—Н	2850-3300
С—С	750–1100
C=C	1620–1680
C=O	1680–1750
С—О	1000-1300
O—H (alcohols)	3230–3550
O—H (acids)	2500–3000

2

.)	Describ spectro	be briefly how positive ion ometer.	ns are form	ned from g	aseous chr	omium ato	oms in a m
	•••••						
							(2 mar
)	What is	s used in a mass spectrom	eter to acc	elerate the	positive ic	ons?	
	•••••						(1 ma
)	What is	s used in a mass spectrom	eter to def	lect the po	sitive ions	?	
)	The ma	ass spectrum of a sample	of chromiu	ım shows f	our peaks.	Use the d	·
)	calcula	ass spectrum of a sample te the relative atomic mas					lata below
)	calcula	te the relative atomic mas					lata below
)	calcula	te the relative atomic mas il places.	s of chrom	ium in the	sample. C	Give your a	
)	calcula	te the relative atomic mas l places.	ss of chrom	52	sample. C	Sive your a	lata below
)	calcula	te the relative atomic mas l places.	ss of chrom	52	sample. C	Sive your a	lata below
	calcula	te the relative atomic mas l places.	ss of chrom	52	sample. C	Sive your a	lata below
)	calcula	te the relative atomic mas l places.	ss of chrom	52	sample. C	Sive your a	lata below

Turn over ▶

The equation for the reaction between magnesium carbonate and hydrochloric acid is given below.

$$MgCO_3 + 2HCl \rightarrow MgCl_2 + H_2O + CO_2$$

When 75.0 cm³ of 0.500 mol dm⁻³ hydrochloric acid were added to 1.25 g of impure MgCO₃ some acid was left unreacted. This unreacted acid required 21.6 cm³ of a 0.500 mol dm⁻³ solution of sodium hydroxide for complete reaction.

(i)	Calculate the number of moles of HCl in 75.0 cm ³ of 0.500 mol dm ⁻³ hydrochloric
	acid.

(ii)	Calculate the number of moles of NaOH used to neutralise the unreacted HCl.

(iii)	Show that the number of moles of HCl which reacted with the MgCO ₃ in the
	sample was 0.0267

(iv)	Calculate the number of moles and the mass of MgCO ₃ in the sample, and hence
	deduce the percentage by mass of MgCO ₃ in the sample.

Moles of $MgCO_3$
Mass of $MgCO_3$
Percentage of MgCO ₃

(8 marks)

APW/0104/CHM1

(b)	A compound contains 36.5% of sodium and 25.5% of sulphur by mass, the rest being
	oxygen.

7

(i)	Use this information to show that the empirical formula of the compound is Na ₂ SO ₃
(ii)	When Na ₂ SO ₃ is treated with an excess of hydrochloric acid, aqueous sodium chloride is formed and sulphur dioxide gas is evolved. Write an equation to represent this reaction.
	(4 marks

TURN OVER FOR THE NEXT QUESTION


Turn over

4	(a)	Com	pound \mathbf{A} is an oxide of sulphur. At 415 K, a gaseous sample of \mathbf{A} , of mass 0.304 g, pied a volume of 127 cm ³ at a pressure of 103 kPa.
		samp	the ideal gas equation and use it to calculate the number of moles of A in the ble, and hence calculate the relative molecular mass of A . gas constant $R = 8.31 \mathrm{J K^{-1} mol^{-1}})$
		Ideal	gas equation
		Calc	ulation
		•••••	
		•••••	
		•••••	(5 marks)
	(b)		presence of sulphate ions in an aqueous solution can be shown by means of a simple nical test.
		(i)	Identify a reagent you would use in this chemical test.
		(ii)	State what you would observe if the test were positive.
		(iii)	Write an ionic equation for the reaction occurring when the test is positive.
			(3 marks)



The diagram below shows the values of the first ionisation energies of some of the elements in Period 3.

First ionisation energy/kJ mol⁻¹



(a) On the above diagram, use crosses to mark the approximate positions of the values of the first ionisation energies for the elements Na, P and S. Complete the diagram by joining the crosses. (3 marks)

(b)	Explain the general increase in the values of the first ionisation energies of the elemen
	Va–Ar.

	 		•••••
•••••	 	•••••	(3 marks)

(c) In terms of the electron sub-levels involved, explain the position of aluminium and the position of sulphur in the diagram.

Explanation for aluminium
Explanation for sulphur

(4 marks)



SECTION B

Answer the question below in the space provided on pages 10 to 12 of this booklet.

- 6 (a) Iodine and graphite crystals both contain covalent bonds and yet the physical properties of their crystals are very different. For iodine and graphite, state and explain the differences in their melting points and in their electrical conductivities. (*9 marks*)
 - (b) Draw the shape of the BeCl₂ molecule and explain why it has this shape. State and explain the effect that an isolated Be²⁺ ion would have on an isolated Cl⁻ ion and explain how this effect would lead to the formation of a covalent bond. Give one chemical property of Be(OH)₂ which is atypical of the chemistry of Group II hydroxides. (6 marks)

END OF QUESTIONS