

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

Leave blank
-------------

General Certificate of Education  
June 2006  
Advanced Subsidiary Examination



**CHEMISTRY**  
**Unit 1 Atomic Structure, Bonding and Periodicity**

**CHM1**

Wednesday 7 June 2006 9.00 am to 10.00 am

**For this paper you must have**

- a calculator.

Time allowed: 1 hour

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Answer questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

**Information**

- The maximum mark for this paper is 60.
- The marks for questions are shown in brackets.
- You are expected to use a calculator where appropriate.
- Your answers to the question in **Section B** should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

**Advice**

- You are advised to spend about 45 minutes on **Section A** and about 15 minutes on **Section B**.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
Total (Column 1) →			
Total (Column 2) →			
TOTAL			
Examiner's Initials			

## SECTION A

Answer **all** questions in the spaces provided.

- 1 (a) State, in terms of the fundamental particles present, the meaning of the term *isotopes*.

.....  
 .....  
 (1 mark)

- (b) An atom contains one more proton than, but the same number of neutrons as, an atom of  $^{36}\text{S}$ . Deduce the symbol, including the mass number and the atomic number, of this atom.

.....  
 (2 marks)

- (c) The table below gives the relative abundance of each isotope in a mass spectrum of a sample of germanium, Ge.

$m/z$	70	72	74
Relative abundance (%)	24.4	32.4	43.2

- (i) Complete the electron arrangement of a Ge atom.

$1s^2$  .....

- (ii) Use the data above to calculate the relative atomic mass of this sample of germanium. Give your answer to one decimal place.

.....  
 .....  
 .....  
 .....

# The Periodic Table of the Elements

- The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

		I	II	III	IV	V	VI	VII	0					
1.0	<b>H</b> Hydrogen								4.0 <b>He</b> Helium 2					
6.9	<b>Li</b> Lithium 3	9.0 <b>Be</b> Beryllium 4	6.9 <b>Li</b> Lithium 3		10.8 <b>B</b> Boron 5	12.0 <b>C</b> Carbon 6	14.0 <b>N</b> Nitrogen 7	16.0 <b>O</b> Oxygen 8	19.0 <b>F</b> Fluorine 9	20.2 <b>Ne</b> Neon 10				
23.0	<b>Na</b> Sodium 11	24.3 <b>Mg</b> Magnesium 12	atomic number		27.0 <b>Al</b> Aluminium 13	28.1 <b>Si</b> Silicon 14	31.0 <b>P</b> Phosphorus 15	32.1 <b>S</b> Sulphur 16	35.5 <b>Cl</b> Chlorine 17	39.9 <b>Ar</b> Argon 18				
39.1	<b>K</b> Potassium 19	40.1 <b>Ca</b> Calcium 20	54.9 <b>Mn</b> Manganese 25	55.8 <b>Fe</b> Iron 26	58.7 <b>Ni</b> Nickel 28	63.5 <b>Cu</b> Copper 29	65.4 <b>Zn</b> Zinc 30	69.7 <b>Ga</b> Gallium 31	72.6 <b>Ge</b> Germanium 32	74.9 <b>As</b> Arsenic 33	79.0 <b>Se</b> Selenium 34	79.9 <b>Br</b> Bromine 35	83.8 <b>Kr</b> Krypton 36	
85.5	<b>Rb</b> Rubidium 37	87.6 <b>Sr</b> Strontium 38	98.9 <b>Tc</b> Technetium 43	101.1 <b>Ru</b> Ruthenium 44	106.4 <b>Pd</b> Palladium 46	107.9 <b>Ag</b> Silver 47	112.4 <b>Cd</b> Cadmium 48	114.8 <b>In</b> Indium 49	118.7 <b>Sn</b> Tin 50	121.8 <b>Sb</b> Antimony 51	127.6 <b>Te</b> Tellurium 52	126.9 <b>I</b> Iodine 53	131.3 <b>Xe</b> Xenon 54	
132.9	<b>Cs</b> Caesium 55	137.3 <b>Ba</b> Barium 56	186.2 <b>Re</b> Rhenium 75	190.2 <b>Os</b> Osmium 76	195.1 <b>Pt</b> Platinum 78	197.0 <b>Au</b> Gold 79	200.6 <b>Hg</b> Mercury 80	204.4 <b>Tl</b> Thallium 81	207.2 <b>Pb</b> Lead 82	209.0 <b>Bi</b> Bismuth 83	210.0 <b>Po</b> Polonium 84	210.0 <b>At</b> Astatine 85	222.0 <b>Rn</b> Radon 86	
223.0	<b>Fr</b> Francium 87	226.0 <b>Ra</b> Radium 88												
										88 – 71 Lanthanides				
										89 – 103 Actinides				
140.1	<b>Ce</b> Cerium 58	140.9 <b>Pr</b> Praseodymium 59	144.2 <b>Nd</b> Neodymium 60	144.9 <b>Pm</b> Promethium 61	150.4 <b>Sm</b> Samarium 62	157.3 <b>Gd</b> Gadolinium 64	158.9 <b>Tb</b> Terbium 65	162.5 <b>Dy</b> Dysprosium 66	164.9 <b>Ho</b> Holmium 67	167.3 <b>Er</b> Erbium 68	168.9 <b>Tm</b> Thulium 69	173.0 <b>Yb</b> Ytterbium 70	175.0 <b>Lu</b> Lutetium 71	
232.0	<b>Th</b> Thorium 90	231.0 <b>Pa</b> Protactinium 91	238.0 <b>U</b> Uranium 92	237.0 <b>Np</b> Neptunium 93	239.1 <b>Pu</b> Plutonium 94	247.1 <b>Cm</b> Curium 96	247.1 <b>Bk</b> Berkelium 97	252.1 <b>Cf</b> Californium 98	(252) <b>Es</b> Einsteinium 99	(257) <b>Fm</b> Fermium 100	(258) <b>Md</b> Mendelevium 101	(259) <b>No</b> Nobelium 102	(260) <b>Lr</b> Lawrencium 103	

Gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

**Table 1**  
Proton n.m.r chemical shift data

Type of proton	$\delta/\text{ppm}$
$\text{RCH}_3$	0.7–1.2
$\text{R}_2\text{CH}_2$	1.2–1.4
$\text{R}_3\text{CH}$	1.4–1.6
$\text{RCOCH}_3$	2.1–2.6
$\text{ROCH}_3$	3.1–3.9
$\text{RCOOCH}_3$	3.7–4.1
$\text{ROH}$	0.5–5.0

**Table 2**  
Infra-red absorption data

Bond	Wavenumber/ $\text{cm}^{-1}$
$\text{C—H}$	2850–3300
$\text{C—C}$	750–1100
$\text{C=C}$	1620–1680
$\text{C=O}$	1680–1750
$\text{C—O}$	1000–1300
$\text{O—H}$ (alcohols)	3230–3550
$\text{O—H}$ (acids)	2500–3000

- (iii) State what is adjusted in a mass spectrometer in order to direct ions with different  $m/z$  values onto the detector. Explain your answer.

*Adjustment* .....

*Explanation* .....

.....

.....

- (iv) One of the isotopes of Ge, given in the table in part (c), has an ion that forms a small peak in the mass spectrum which is indistinguishable from a peak produced by  $^{36}\text{S}^+$  ions. Identify this Ge ion and explain your answer.

*Ion* .....

*Explanation* .....

.....


(8 marks)

11


**Turn over for the next question**

2 At room temperature, both sodium metal and sodium chloride are crystalline solids which contain ions.

(a) On the diagrams for sodium metal and sodium chloride below, mark the charge for each ion.



Sodium metal



Sodium chloride

(2 marks)

(b) (i) Explain how the ions are held together in solid sodium metal.

.....  
 .....

(ii) Explain how the ions are held together in solid sodium chloride.

.....  
 .....

(iii) The melting point of sodium chloride is much higher than that of sodium metal. What can be deduced from this information?

.....  
 .....

(3 marks)

(c) Compare the electrical conductivity of solid sodium metal with that of solid sodium chloride. Explain your answer.

*Comparison* .....

.....

*Explanation* .....

.....

.....

(3 marks)

(d) Explain why sodium metal is malleable (can be hammered into shape).

.....  
.....

(1 mark)

(e) Sodium chlorate(V),  $\text{NaClO}_3$ , contains 21.6% by mass of sodium, 33.3% by mass of chlorine and 45.1% by mass of oxygen.

(i) Use the above data to show that the empirical formula of sodium chlorate(V) is  $\text{NaClO}_3$

.....  
.....  
.....  
.....  
.....

(ii) Sodium chlorate(V) may be prepared by passing chlorine into hot aqueous sodium hydroxide. Balance the equation for this reaction below.




(3 marks)

12
----

**Turn over for the next question**

- 3 Nitroglycerine,  $C_3H_5N_3O_9$ , is an explosive which, on detonation, decomposes rapidly to form a large number of gaseous molecules. The equation for this decomposition is given below.



- (a) A sample of nitroglycerine was detonated and produced 0.350 g of oxygen gas.

- (i) State what is meant by the term *one mole* of molecules.

.....

- (ii) Calculate the number of moles of oxygen gas produced in this reaction, and hence deduce the total number of moles of gas formed.

*Moles of oxygen gas* .....

*Total moles of gas* .....

.....

.....

- (iii) Calculate the number of moles, and the mass, of nitroglycerine detonated.

*Moles of nitroglycerine* .....

.....

*Mass of nitroglycerine* .....

.....

.....

(7 marks)

- (b) A second sample of nitroglycerine was placed in a strong sealed container and detonated. The volume of this container was  $1.00 \times 10^{-3} \text{ m}^3$ . The resulting decomposition produced a total of 0.873 mol of gaseous products at a temperature of 1100 K.

State the ideal gas equation and use it to calculate the pressure in the container after detonation.

(The gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

*Ideal gas equation* .....

*Pressure* .....


.....

.....

.....



4 (a) The diagram below shows the melting points of some of the elements in Period 3.



(i) On the diagram, use crosses to mark the approximate positions of the melting points for the elements silicon, chlorine and argon. Complete the diagram by joining the crosses.

(ii) By referring to its structure and bonding, explain your choice of position for the melting point of silicon.

.....

.....

.....

(iii) Explain why the melting point of sulphur, S<sub>8</sub>, is higher than that of phosphorus, P<sub>4</sub>

.....

.....

(8 marks)

(b) State and explain the trend in melting point of the Group II elements Ca–Ba.

*Trend* .....

*Explanation* .....

.....

.....

(3 marks)





A series of horizontal dotted lines for writing, spanning the width of the page.