

General Certificate of Education

Chemistry 5421

CHM2 Foundation Physical and Inorganic Chemistry

Mark Scheme

2007 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2007 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

CHM2

Question 1

(a)	(i)	Curve starts at the origin and does not touch the <i>x</i> axis on the right hand side. (should be asymptotic)	
		Curve is skewed left	(1)
	(ii)	Minimum energy needed for a reaction to occur	(2)
	(iii)	Area under candidates curve from $E_{\rm a}$ to the right of it should be shaded	(1)
(b)	(i)	None	(1)
	(ii)	More molecules have energy greater than the activation energy	(1)
		Therefore there are more successful collisions	(1)
(c)	Increa	ses rate	(1)
		vs an alternative route has a lower activation energy	(1) (1)
Ques	tion 2		
(a)	(i)	Reduction is the gain of electrons	(1)
		A reducing agent donates electrons (not pairs of electrons)	(1)
	(ii)	C or CO	(1)
		Fe ₂ O ₃ + 3C \rightarrow 2Fe + 3CO Or Fe ₂ O ₃ + 3CO \rightarrow 2Fe + 3CO ₂ Or 2 Fe ₂ O ₃ + 3C \rightarrow 4Fe + 3CO ₂	(1)
	(iii)	High temperature needed or in range 700°C to 2000°C	(1)
(b)	Blast with oxygen		
	so the carbon is oxidised (or use equation e.g. $C + O_2 \rightarrow CO_2$)		(1)
(c)	(i)	$TiO_2 + 2C + 2CI_2 \rightarrow TiCI_4 + 2CO$ Or $TiO_2 + C + 2CI_2 \rightarrow TiCI_4 + CO_2$	(1)
		$TiCl_4 + 4Na \rightarrow Ti + 4NaCl$ Or $TiCl_4 + 2Mg \rightarrow Ti + 2MgCl_2$	(1)

	(ii)	Argon	(1)	
		Prevent Ti reacting with oxygen, nitrogen, air Or prevent Na/Mg reacting with oxygen, nitrogen, water, air Or prevent TiCl ₄ hydrolysing/ reacting with water	(1)	
(d)	Any tv	wo from Mg/Na/Cl ₂ have to be made first by electrolysis Batch process Argon atmosphere needed High temperatures needed in both stages	(2)	
Ques	stion 3			
(a)	Is forr	actants and products in their standard states	(1) (1) (1)	
(b)	$\frac{1}{2}$ H ₂ (g) + $\frac{1}{2}$ N ₂ (g) + $\frac{3}{2}$ O ₂ (g) \rightarrow HNO ₃ (I)	(1) for equation 1) for state symbols	
(c)	Entha	lpy change is independent of the route taken	(1)	
(d)	(i)	It is an element	(1)	
	(ii)	$\Delta H = \Sigma \Delta H_{\rm f}$ products - $\Sigma \Delta H_{\rm f}$ reactants or correct cycle	(1)	
		-128 = 2x - (-286 + 2(34))	(1)	
		-346 + 2 <i>x</i>	(1)	
		$x = -173 \text{ kJmol}^{-1}$	(1)	
Ques	stion 4			
(a)	Rate	of forward reaction = rate of backward reaction	(1)	
	Conce	entration of reactants and products are constant (QoL)	(1)	
(b)	Increa	ases	(1)	
	More moles on the left hand side of equation			
	Equilibrium moves to reduce pressure/ oppose the change			
(c)	Decre	eases	(1)	
	Reaction is exothermic			
	Equilibrium moves to absorb heat/ lower the temperature/ oppose the change (

(d) (+) 5 (1)

(e) (+) 5 (1)

(f) (1)

Question 5

(a) Increase (1)

$$2H^* + H_2SO_4 + 2e^- \rightarrow SO_2 + 2H_2O$$

$$2H^* + H_2SO_4 + 2l^- \rightarrow SO_2 + 2H_2O + l_2$$
(allow alternative correct equations with $SO_4^{2^+}$)

$$H_2S \text{ or } S$$
(1)

(c) Yellowy solution turns to a brown solution/ black ppt (need both colours) (1)

$$Br_2 + 2KI \rightarrow l_2 + 2KBr$$
(1)

$$Br_2 \text{ is an oxidising agent}$$
(1)

(d) Add silver nitrate solution (1)

$$KBr \text{ forms creamy ppt}$$
(1)

$$KI \text{ forms yellow ppt}$$
(1)

$$AgNO_3 + KBr \rightarrow AgBr + KNO_3 \text{ or } AgNO_3 + KI \rightarrow AgI + KNO_3 + KI \rightarrow AgI + KNO_$$

Alternative answers to d (1) Either add chlorine /Cl₂ Yellow brown/yellow/ orange brown solution formed with KBr (1) Brown solution/black ppt formed with KI (1) $Cl_2 +2 KBr \rightarrow Br_2 + 2 KCl$ (1) or $Cl_2 + 2 KI \rightarrow l_2 + 2 KCI$ add bromine/Br₂ or (1) no reaction with bromide ions (1) Brown solution/black ppt formed with KI (1) $Br_2 + 2 KI \rightarrow I_2 + 2 KBr$ (1) Confirmatory test for either alternative above answers Either Add starch (1) (1) No change with Bromine formed Goes black with iodine formed (1) Or add CCI₄/ organic solvent (1) Bromine goes yellow/orange brown (1) lodine goes purple (1)