

|                     |  |  |  |  |  |                  |  |  |  |  |  |
|---------------------|--|--|--|--|--|------------------|--|--|--|--|--|
| Surname             |  |  |  |  |  | Other Names      |  |  |  |  |  |
| Centre Number       |  |  |  |  |  | Candidate Number |  |  |  |  |  |
| Candidate Signature |  |  |  |  |  |                  |  |  |  |  |  |

|             |
|-------------|
| Leave blank |
|-------------|

General Certificate of Education  
January 2006  
Advanced Subsidiary Examination



**CHEMISTRY** **CHM2**  
**Unit 2 Foundation Physical and Inorganic Chemistry**

Wednesday 11 January 2006 9.00 am to 10.00 am

**For this paper you must have**

- a calculator.

Time allowed: 1 hour

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Answer questions in **Section A** and **Section B** in the spaces provided.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

**Information**

- The maximum mark for this paper is 60.
- The marks for each question are shown in brackets.
- This paper carries 30 per cent of the total marks for AS. For Advanced Level this paper carries 15 per cent of the total marks.
- You are expected to use a calculator where appropriate.
- The following data may be required.  
Gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
- Your answers to the question in **Section B** should be written in continuous prose, where appropriate.
- You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

**Advice**

- You are advised to spend about 45 minutes on **Section A** and about 15 minutes on **Section B**.

| For Examiner's Use  |      |        |      |
|---------------------|------|--------|------|
| Number              | Mark | Number | Mark |
| 1                   |      |        |      |
| 2                   |      |        |      |
| 3                   |      |        |      |
| 4                   |      |        |      |
| 5                   |      |        |      |
|                     |      |        |      |
|                     |      |        |      |
|                     |      |        |      |
| Total (Column 1) →  |      |        |      |
| Total (Column 2) →  |      |        |      |
| TOTAL               |      |        |      |
| Examiner's Initials |      |        |      |

## SECTION A

Answer **all** questions in the spaces provided.


- 1 (a) Define the term *standard enthalpy of combustion*,  $\Delta H_c^\ominus$

.....  
 .....  
 .....

(3 marks)

- (b) Use the mean bond enthalpy data from the table and the equation given below to calculate a value for the standard enthalpy of combustion of propene. All substances are in the gaseous state.

| Bond                                      | C=C | C—C | C—H | O=O | O=C | O—H |
|---|-----|-----|-----|-----|-----|-----|
| Mean bond enthalpy / kJ mol <sup>-1</sup> | 612 | 348 | 412 | 496 | 743 | 463 |



.....  
 .....  
 .....  
 .....

(3 marks)

- (c) State why the standard enthalpy of formation,  $\Delta H_f^\ominus$ , of oxygen is zero.

.....

(1 mark)

## The Periodic Table of the Elements

■ The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

|       |                             | I     | II                           | Key   |                                 |       |                              |       |                              |       |                               |       |                               | III   | IV                           | V     | VI                            | VII   | 0                            |       |                                |       |                                |       |                             |       |                                 |       |                              |       |                                |       |                             |       |                            |  |
|-------|-----------------------------|-------|------------------------------|-------|---------------------------------|-------|------------------------------|-------|------------------------------|-------|-------------------------------|-------|-------------------------------|-------|------------------------------|-------|-------------------------------|-------|------------------------------|-------|--------------------------------|-------|--------------------------------|-------|-----------------------------|-------|---------------------------------|-------|------------------------------|-------|--------------------------------|-------|-----------------------------|-------|----------------------------|--|
| 1.0   | <b>H</b><br>Hydrogen<br>1   | 9.0   | <b>Be</b><br>Beryllium<br>4  | 45.0  | <b>Sc</b><br>Scandium<br>21     | 47.9  | <b>Ti</b><br>Titanium<br>22  | 50.9  | <b>V</b><br>Vanadium<br>23   | 52.0  | <b>Cr</b><br>Chromium<br>24   | 54.9  | <b>Mn</b><br>Manganese<br>25  | 55.8  | <b>Fe</b><br>Iron<br>26      | 58.9  | <b>Co</b><br>Cobalt<br>27     | 58.7  | <b>Ni</b><br>Nickel<br>28    | 63.5  | <b>Cu</b><br>Copper<br>29      | 65.4  | <b>Zn</b><br>Zinc<br>30        | 69.7  | <b>Ga</b><br>Gallium<br>31  | 72.6  | <b>Ge</b><br>Germanium<br>32    | 74.9  | <b>As</b><br>Arsenic<br>33   | 79.0  | <b>Se</b><br>Selenium<br>34    | 79.9  | <b>Br</b><br>Bromine<br>35  | 83.8  | <b>Kr</b><br>Krypton<br>36 |  |
| 6.9   | <b>Li</b><br>Lithium<br>3   | 23.0  | <b>Na</b><br>Sodium<br>11    | 88.9  | <b>Y</b><br>Yttrium<br>39       | 91.2  | <b>Zr</b><br>Zirconium<br>40 | 92.9  | <b>Nb</b><br>Niobium<br>41   | 95.9  | <b>Mo</b><br>Molybdenum<br>42 | 98.9  | <b>Tc</b><br>Technetium<br>43 | 101.1 | <b>Ru</b><br>Ruthenium<br>44 | 102.9 | <b>Rh</b><br>Rhodium<br>45    | 106.4 | <b>Pd</b><br>Palladium<br>46 | 107.9 | <b>Ag</b><br>Silver<br>47      | 112.4 | <b>Cd</b><br>Cadmium<br>48     | 114.8 | <b>In</b><br>Indium<br>49   | 118.7 | <b>Sn</b><br>Tin<br>50          | 121.8 | <b>Sb</b><br>Antimony<br>51  | 127.6 | <b>Te</b><br>Tellurium<br>52   | 126.9 | <b>I</b><br>Iodine<br>53    | 131.3 | <b>Xe</b><br>Xenon<br>54   |  |
| 39.1  | <b>K</b><br>Potassium<br>19 | 40.1  | <b>Ca</b><br>Calcium<br>20   | 138.9 | <b>La</b><br>Lanthanum<br>57    | 178.5 | <b>Hf</b><br>Hafnium<br>72   | 180.9 | <b>Ta</b><br>Tantalum<br>73  | 183.9 | <b>W</b><br>Tungsten<br>74    | 186.2 | <b>Re</b><br>Rhenium<br>75    | 190.2 | <b>Os</b><br>Osmium<br>76    | 192.2 | <b>Ir</b><br>Iridium<br>77    | 195.1 | <b>Pt</b><br>Platinum<br>78  | 197.0 | <b>Au</b><br>Gold<br>79        | 200.6 | <b>Hg</b><br>Mercury<br>80     | 204.4 | <b>Tl</b><br>Thallium<br>81 | 207.2 | <b>Pb</b><br>Lead<br>82         | 209.0 | <b>Bi</b><br>Bismuth<br>83   | 210.0 | <b>Po</b><br>Polonium<br>84    | 210.0 | <b>At</b><br>Astatine<br>85 | 222.0 | <b>Rn</b><br>Radon<br>86   |  |
| 85.5  | <b>Rb</b><br>Rubidium<br>37 | 87.6  | <b>Sr</b><br>Strontium<br>38 | 227   | <b>Ac</b><br>Actinium<br>89     |       |                              |       |                              |       |                               |       |                               |       |                              |       |                               |       |                              |       |                                |       |                                |       |                             |       |                                 |       |                              |       |                                |       |                             |       |                            |  |
| 132.9 | <b>Cs</b><br>Caesium<br>55  | 137.3 | <b>Ba</b><br>Barium<br>56    | 140.9 | <b>Pr</b><br>Praseodymium<br>59 | 140.9 | <b>Ce</b><br>Cerium<br>58    | 144.2 | <b>Nd</b><br>Neodymium<br>60 | 144.9 | <b>Pm</b><br>Promethium<br>61 | 150.4 | <b>Sm</b><br>Samarium<br>62   | 152.0 | <b>Eu</b><br>Europium<br>63  | 157.3 | <b>Gd</b><br>Gadolinium<br>64 | 158.9 | <b>Tb</b><br>Terbium<br>65   | 162.5 | <b>Dy</b><br>Dysprosium<br>66  | 164.9 | <b>Ho</b><br>Holmium<br>67     | 167.3 | <b>Er</b><br>Erbium<br>68   | 168.9 | <b>Tm</b><br>Thulium<br>69      | 173.0 | <b>Yb</b><br>Ytterbium<br>70 | 175.0 | <b>Lu</b><br>Lutetium<br>71    |       |                             |       |                            |  |
| 223.0 | <b>Fr</b><br>Francium<br>87 | 226.0 | <b>Ra</b><br>Radium<br>88    | 231.0 | <b>Pa</b><br>Protactinium<br>91 | 231.0 | <b>Th</b><br>Thorium<br>90   | 238.0 | <b>U</b><br>Uranium<br>92    | 237.0 | <b>Np</b><br>Neptunium<br>93  | 239.1 | <b>Pu</b><br>Plutonium<br>94  | 243.1 | <b>Am</b><br>Americium<br>95 | 247.1 | <b>Cm</b><br>Curium<br>96     | 247.1 | <b>Bk</b><br>Berkelium<br>97 | 252.1 | <b>Cf</b><br>Californium<br>98 | (252) | <b>Es</b><br>Einsteinium<br>99 | (257) | <b>Fm</b><br>Fermium<br>100 | (258) | <b>Md</b><br>Mendelevium<br>101 | (259) | <b>No</b><br>Nobelium<br>102 | (260) | <b>Lr</b><br>Lawrencium<br>103 |       |                             |       |                            |  |

\* 58 – 71 Lanthanides

† 90 – 103 Actinides

**Table 1**  
Proton n.m.r chemical shift data

| Type of proton          | $\delta/\text{ppm}$ |
|-------------------------|---------------------|
| $\text{RCH}_3$          | 0.7–1.2             |
| $\text{R}_2\text{CH}_2$ | 1.2–1.4             |
| $\text{R}_3\text{CH}$   | 1.4–1.6             |
| $\text{RCOCH}_3$        | 2.1–2.6             |
| $\text{ROCH}_3$         | 3.1–3.9             |
| $\text{RCOOCH}_3$       | 3.7–4.1             |
| $\text{ROH}$            | 0.5–5.0             |

**Table 2**  
Infra-red absorption data

| Bond                    | Wavenumber/ $\text{cm}^{-1}$ |
|-------------------------|------------------------------|
| $\text{C—H}$            | 2850–3300                    |
| $\text{C—C}$            | 750–1100                     |
| $\text{C=C}$            | 1620–1680                    |
| $\text{C=O}$            | 1680–1750                    |
| $\text{C—O}$            | 1000–1300                    |
| $\text{O—H}$ (alcohols) | 3230–3550                    |
| $\text{O—H}$ (acids)    | 2500–3000                    |

- (d) Use the data from the table below to calculate a more accurate value for the standard enthalpy of combustion of propene.

| Compound  | C <sub>3</sub> H <sub>6</sub> (g) | CO <sub>2</sub> (g) | H <sub>2</sub> O(g) |
|---|-----------------------------------|---------------------|---------------------|
| Standard enthalpy of formation, $\Delta H_f^\ominus / \text{kJ mol}^{-1}$ | +20                               | -394                | -242                |

.....

.....

.....

.....

.....

(3 marks)

- (e) Explain why your answer to part (b) is a less accurate value than your answer to part (d).

.....

.....


(2 marks)

12

Turn over for the next question

Turn over 

- 2 The gas-phase reaction between hydrogen and chlorine is very slow at room temperature.



- (a) Define the term *activation energy*.

.....  
.....  
(2 marks)

- (b) Give **one** reason why the reaction between hydrogen and chlorine is very slow at room temperature.

.....  
.....  
(1 mark)

- (c) Explain why an increase in pressure, at constant temperature, increases the rate of reaction between hydrogen and chlorine.

.....  
.....  
(2 marks)

- (d) Explain why a small increase in temperature can lead to a large increase in the rate of reaction between hydrogen and chlorine.

.....  
.....  
(2 marks)

- (e) Give the meaning of the term *catalyst*.

.....  
.....  
(1 mark)

- (f) Suggest **one** reason why a solid catalyst for a gas-phase reaction is often in the form of a powder.

.....  
(1 mark)

- 3 At high temperatures, nitrogen is oxidised by oxygen to form nitrogen monoxide in a reversible reaction as shown in the equation below.



- (a) In terms of electrons, give the meaning of the term *oxidation*.

.....  
(1 mark)

- (b) State and explain the effect of an increase in pressure, and the effect of an increase in temperature, on the yield of nitrogen monoxide in the above equilibrium.

*Effect of an increase in pressure on the yield* .....

*Explanation* .....

.....  
.....

*Effect of an increase in temperature on the yield* .....

*Explanation* .....

.....  
.....

(6 marks)

- (c) Nitrogen monoxide, NO, is formed when silver metal reduces nitrate ions, NO<sub>3</sub><sup>-</sup>, in acid solution.

- (i) Deduce the oxidation state of nitrogen in NO and in NO<sub>3</sub><sup>-</sup>

NO .....

NO<sub>3</sub><sup>-</sup> .....

- (ii) Write a half-equation for the reduction of NO<sub>3</sub><sup>-</sup> ions in acid solution to form nitrogen monoxide and water.

.....

- (iii) Write a half-equation for the oxidation of silver metal to Ag<sup>+</sup>(aq) ions.

.....

- (iv) Hence, deduce an overall equation for the reaction between silver metal and nitrate ions in acid solution.

.....

(5 marks)

- 4 (a) State the trend in electronegativity of the elements down Group VII. Explain this trend.

*Trend* .....

*Explanation* .....

.....

.....

(3 marks)

- (b) (i) State the trend in reducing ability of the halide ions down Group VII.

.....

- (ii) Give an example of a reagent which could be used to show that the reducing ability of bromide ions is different from that of chloride ions.

.....

(2 marks)

- (c) The addition of silver nitrate solution followed by dilute aqueous ammonia can be used as a test to distinguish between chloride and bromide ions. For each ion, state what you would observe if an aqueous solution containing the ion was tested in this way.

*Observations with chloride ions* .....

.....

*Observations with bromide ions* .....

.....

(4 marks)

- (d) Write an equation for the reaction between chlorine and cold, dilute aqueous sodium hydroxide. Give two uses of the resulting solution.

*Equation* .....

*Use 1* .....

*Use 2* .....

(3 marks)



**SECTION B**

Answer the question in the space provided.

**5** In this question, where appropriate, illustrate your answer with equations.

- (a) Explain how iron is produced in the Blast Furnace from an iron ore that does not contain sulphur impurities. In your answer, state the source of the energy for this process and mention any environmental problems that may arise from the operation of the Blast Furnace.

Explain why limestone is added to the Blast Furnace.

*(10 marks)*

- (b) Phosphorus is a significant impurity in iron from the Blast Furnace. Explain how this impurity is removed in the steel-making process.

*(5 marks)*

**END OF QUESTIONS**

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....


.....

.....

.....



.....


Turn over 



LEAVE  
MARGIN  
BLANK



A large rectangular area containing 25 horizontal dotted lines, intended for writing answers.






LEAVE  
MARGIN  
BLANK

Blank writing area with horizontal dotted lines for text entry.



Turn over 



LEAVE  
MARGIN  
BLANK

Area containing horizontal dotted lines for writing.

