Surname					Other	Names			
Centre Nu	mber			Candidate Number					
Candidate	Signat	ure							·

For Examiner's Use

General Certificate of Education June 2007 Advanced Subsidiary Examination



CHEMISTRY CHM2

Unit 2 Foundation Physical and Inorganic Chemistry

Wednesday 6 June 2007 9.00 am to 10.00 am

For this paper you must have

a calculator.

Time allowed: 1 hour

## **Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer all questions.
- Answer questions in **Section A** and **Section B** in the spaces provided.
- All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

## **Information**

- The maximum mark for this paper is 60.
- The marks for each question are shown in brackets.
- You are expected to use a calculator where appropriate.
- Write your answers to the question in Section B in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

### **Advice**

• You are advised to spend about 45 minutes on **Section A** and about 15 minutes on **Section B**.

For Examiner's Use					
Question	Mark	Question	Mark		
1					
2					
3					
4					
5					
Total (Column 1)					
Total (Column 2) —>					
TOTAL					
Examine	r's Initials				

# SECTION A

Answer all questions in the spaces provided.

1	(a)	(i)	On the axes below, draw a Maxwell-Boltzmann distribution of molecul energies for a gas at temperature $T$ . $E_a$ is the activation energy for a reaction involving this gas.	ar
			umber of olecules	
			$E_a$ Energy	(2 marks)
		(ii)	State the meaning of the term activation energy.	
				(2 marks)
		(iii)	Shade on the graph the area that represents the number of molecules with	,
		()	react at temperature $T$ .	(1 mark)
	(b)	(i)	State the effect on the activation energy of increasing the temperature.	
				(1 mark)
		(ii)	Explain why reactions involving gases become faster as the temperature increases.	e
				(2 marks)
	(c)		nixture of gases is allowed to react in the presence of a catalyst. te and explain the effect of a catalyst on the rate of this reaction.	
		Effec	ect	
		Expl	olanation	

www.theallpapers.com

# The Periodic Table of the Elements

■ The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

_	=											=	≥	>	>	<b>=</b>	0
1.0 <b>H</b> Hydrogen		_	Key														4.0 <b>He</b> Helium 2
6.9 <b>Li</b> Lithium	9.0 <b>Be</b> Beryllium 4		relative atomic	relative atomic mass		6.9 <b>Li</b> Lithium						10.8 <b>B</b> Boron	12.0 <b>C</b> Carbon	14.0 <b>N</b> Nitrogen	16.0 <b>O</b> Oxygen	19.0 <b>F</b> Fluorine	20.2 <b>Ne</b> Neon
23.0 23 2 23.0 23 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	H.3 Mg agnesium											_	.8.1 <b>Si</b> Silicon	31.0 <b>P</b> Phosphorus 15	32.1 <b>S</b> Sulphur 16		39.9 <b>Ar</b> Argon
	_	Scandium 21	_ ا	_	_	ı w	<b>Fe</b> 155.8 150 150 150 150 150 150 150 150 150 150	58.9 <b>Co</b> Cobalt 27	58.7 <b>Nickel</b> 28	63.5 <b>Cu</b> Copper 29	65.4 <b>Zn</b> Zinc 30	69.7 <b>Ga</b> Gallium 31	.2.6 <b>Ge</b> Sermanium	74.9 <b>As</b> Arsenic 33	79.0 <b>Se</b> Selenium 34		83.8 <b>Kr</b> Krypton 36
85.5 <b>Rb</b> Rubidium 37	87.6 Srontium 38	88.9 <b>Y</b> Yttrium 39	91.2 <b>Zr</b> Zirconium 40	92.9 <b>Nb</b> Niobium 41	95.9         98.9         101.1         102.9           Mo         Tc         Ru         Rh           Molybdenum         Technetium         Ruthenium         Rhodium           42         43         44         45	98.9 <b>Tc</b> Technetium	Ruthenium 44		_	107.9 <b>Ag</b> Silver 47		114.8 <b>In</b> Indium 49		≥	127.6 <b>Te</b> Te Tellurium 52		131.3 <b>Xe</b> Xenon 54
	137.3 <b>Ba</b> Barium 56	138.9 <b>La</b> La Lanthanum 57 *	178.5 <b>Hf</b> Hafnium 72	180.9 <b>Ta</b> Tantalum 73	183.9 W Tungsten 74	186.2 <b>Re</b> Rhenium 75	190.2 <b>Os</b> Osmium 76	192.2 <b>    r</b>   <b>  r</b>   Iridium	195.1 <b>Pt</b> Platinum 78	197.0 <b>Au</b> Gold 79	200.6 <b>Hg</b> Mercury 80		207.2 <b>Pb</b> Lead Lead	209.0 <b>Bi</b> Bismuth 83	210.0 <b>Po</b> Polonium 84	210.0 <b>At</b> Astatine 85	222.0 <b>Rn</b> Radon 86
223.0	226.0 <b>Ra</b> Radium 88	227 <b>Ac</b> Actinium 89 †															
thealthanides ada ada ada ada ada ada ada ada ada ad	Lantha	nides		_	140.9 <b>Pr</b> Praseodymium 1	Neodymium   60 (	144.9 Pm Promethium 631 (	150.4 <b>Sm</b> Samarium 62	152.0 <b>Eu</b> Europium (63	157.3 <b>Gd</b> Gadolinium 64	158.9 <b>Tb</b> Terbium	162.5 164.9 <b>Dy Ho</b> Dysprosium Holmium  66  67	164.9 <b>Ho</b> Holmium 67	167.3 <b>Er</b> bit 58	168.9 <b>Tm</b> Thulium 69	173.0 <b>Yb</b> Ytterbium 70	. Tm Yterbium Lutetium 69 70 70 70 70 70 70 70 70 70 70 70 70 70
rs.eom	3 Actini	səp		<b>Th</b> Thorium 90	Protactinium 91	.236.0 <b>U</b> Uranium 92	Np	Pu Pu Plutonium 94	Americium	Curium 96	Bk Bk Berkelium	247.1   252.1   (252)   (252	Einsteinium 99	(257) Fermi 100	Md Mendelevium 101	Nobelium	Lawrencium

Gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ 

**Table 1** Proton n.m.r chemical shift data

Type of proton	δ/ppm
$RCH_3$	0.7–1.2
$R_2CH_2$	1.2–1.4
$R_3CH$	1.4–1.6
$RCOCH_3$	2.1–2.6
$ROCH_3$	3.1–3.9
$RCOOCH_3$	3.7–4.1
ROH	0.5–5.0

**Table 2** Infra-red absorption data

Bond	Wavenumber/cm <sup>-1</sup>
С—Н	2850–3300
С—С	750–1100
C=C	1620–1680
C=O	1680–1750
С—О	1000-1300
O—H (alcohols)	3230–3550
O—H (acids)	2500–3000

(a)			Blast
	(i)	In terms of electrons, state what is meant by reduction and reducing age	ent.
		Reduction	•••••
		Reducing agent	•••••
	(ii)	Identify a reducing agent that can reduce $Fe_2O_3$ to iron in the Blast Furn Write an equation for the reaction between $Fe_2O_3$ and the reducing agen have stated.	
		Reducing agent	•••••
		Equation	•••••
	(iii)	Give one essential condition needed for this reduction.	
			(5 marks)
(b)		± • •	Explain
(c)	(i)	Titanium is extracted from titanium(IV) oxide, TiO <sub>2</sub> , in a two-stage proc Write equations for the reaction occurring in stage 1 and stage 2 of this extraction.	cess.
		Equation for stage 1	
		Equation for stage 2	
	(ii)	Give one essential condition, other than temperature, for the second stag state why it is necessary.	ge, and
		Condition	
		Reason	
(d)	Give		(4 marks)
	Reas	son 1	
	(c)	(ii) (iii) (b) Molhow (c) (i) (d) Give	Furnace by a reducing agent.  (i) In terms of electrons, state what is meant by reduction and reducing agent Reduction Reducing agent Reduction Reducing agent Reducing agent that can reduce Fe <sub>2</sub> O <sub>3</sub> to iron in the Blast Furn Write an equation for the reaction between Fe <sub>2</sub> O <sub>3</sub> and the reducing agent have stated.  Reducing agent Reducing Re

(a)	Defi	ne the term stand	ard enthalpy o	of formation.		
	•••••					
	•••••					(3 marks)
(b)		e an equation, inc all to the standard				an enthalpy change HNO <sub>3</sub>
	•••••					(2 marks)
(c)	State	e Hess's Law.				
	•••••					
	•••••					(1 mark)
(d)	the f	ic acid can be marginal control of $O(1) + 2NO_2(g)$	n.			oxygen according to 8 kJ mol <sup>-1</sup>
	Som	e standard enthal	pies of format	ion, $\Delta H_{\rm f}^{\Theta}$ , are	given in the tal	ole below.
		Substance	H <sub>2</sub> O(1)	NO <sub>2</sub> (g)	O <sub>2</sub> (g)	
	ΔΙ	$H_{\rm f}^{\Theta}/{ m kJmol}^{-1}$	-286	+34	0	
	(i)	State why the st	andard enthal	py of formation	of $O_2(g)$ is ze	ero.
						(1 mark)
	(ii)	TT 4 1 . 1	ove to calculat			
		nitric acid.	ove to calculat	e a value for th	e standard entl	halpy of formation of
				e a value for th	e standard entl	halpy of formation of
				e a value for th	e standard entl	halpy of formation of
				e a value for th	e standard ent	halpy of formation of

www.theallpapers.com

4	Whe	n nitrogen monoxide reacts with oxygen, a dynamic equilibrium is established.
		$2NO(g) + O_2(g) \rightleftharpoons 2NO_2(g) \qquad \Delta H^{\circ} = -115 \text{ kJ mol}^{-1}$
	(a)	State what is meant by dynamic equilibrium.
		(2 marks)
	(b)	State and explain how the total pressure in this equilibrium reaction should be changed to give a higher equilibrium yield of $NO_2$
		Change in pressure
		Explanation
		(3 marks)
	(c)	State and explain the effect of an increase in temperature on the yield of $NO_2$ in this equilibrium reaction.
		Effect
		Explanation
		(3 marks)
	(d)	Deduce the oxidation state of nitrogen in NO <sub>3</sub> and in NO <sub>2</sub> <sup>+</sup>
		NO <sub>3</sub>
		$\mathrm{NO}_2^+$
		(2 marks)

# **SECTION B**

Answer question 5 in the space provided on pages 8 to 12 of this booklet.

_	( )	
5	(a)	State the trend in the reducing ability of the halide ions from fluoride to iodide.  (1 mark)
	(b)	Concentrated sulphuric acid reacts with solid potassium iodide to form a mixture of products. These products include sulphur dioxide and iodine.
		Write half-equations for the formation of iodine from iodide ions, and for the formation of sulphur dioxide from sulphuric acid. Hence write an overall equation for the formation of these products from iodide ions and sulphuric acid.
		Identify one other reduction product formed in the reaction between sulphuric acid and solid potassium iodide.
		(4 marks)
	(c)	State what you would observe when aqueous bromine reacts with a solution of potassium iodide. Write an equation for the reaction.  State the role of bromine in the reaction.
		(3 marks)
	(d)	Give a reagent which could be used to distinguish between separate solutions of potassium bromide and potassium iodide. State what would be observed when this reagent is added to each of the separate solutions of potassium bromide and potassium iodide. Write an equation for <b>one</b> of the reactions. Identify a reagent which could be added to the mixtures from the first test to confirm the identity of the halide ions. State what would be observed in each case.  (7 marks)
		END OF QUESTIONS
•••••	••••••	
•••••	•••••	
•••••	••••••	
•••••	•••••	

••••••••••••
•••••
 •••••
•••••
 •••••

 •••••
 •••••
 •••••
 •••••
 •••••

•••••
•••••
•••••
 •••••
 •••••
•••••
•••••
•••••
•••••
