

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

Leave blank
-------------

General Certificate of Education  
January 2003  
Advanced Level Examination



**CHEMISTRY** **CHM4**  
**Unit 4 Further Physical and Organic Chemistry**

Wednesday 22 January 2003 Morning Session

In addition to this paper you will require: a calculator.
--

Time allowed: 1 hour 30 minutes

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

**Information**

- The maximum mark for this paper is 90.
- Mark allocations are shown in brackets.
- This paper carries 15 per cent of the total marks for Advanced Level.
- You are expected to use a calculator where appropriate.
- The following data may be required.  
Gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
- Your answers to questions in **Section B** should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

**Advice**

- You are advised to spend about 1 hour on **Section A** and about 30 minutes on **Section B**.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
7			
8			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

## The Periodic Table of the Elements

- The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

		I	II	III	IV	V	VI	VII	0								
<b>H</b> Hydrogen									4.0 <b>He</b> Helium 2								
<b>Li</b> Lithium 3	<b>Be</b> Beryllium 4	relative atomic mass — 6.9 atomic number — 3		<b>Li</b> Lithium 3	<b>B</b> Boron 5	<b>C</b> Carbon 6	<b>N</b> Nitrogen 7	<b>O</b> Oxygen 8	<b>F</b> Fluorine 9	<b>Ne</b> Neon 10							
<b>Na</b> Sodium 11	<b>Mg</b> Magnesium 12				<b>Al</b> Aluminium 13	<b>Si</b> Silicon 14	<b>P</b> Phosphorus 15	<b>S</b> Sulphur 16	<b>Cl</b> Chlorine 17	<b>Ar</b> Argon 18							
<b>K</b> Potassium 19	<b>Ca</b> Calcium 20	<b>Sc</b> Scandium 21	<b>Ti</b> Titanium 22	<b>V</b> Vanadium 23	<b>Cr</b> Chromium 24	<b>Mn</b> Manganese 25	<b>Fe</b> Iron 26	<b>Co</b> Cobalt 27	<b>Ni</b> Nickel 28	<b>Cu</b> Copper 29	<b>Zn</b> Zinc 30	<b>Ga</b> Gallium 31	<b>Ge</b> Germanium 32	<b>As</b> Arsenic 33	<b>Se</b> Selenium 34	<b>Br</b> Bromine 35	<b>Kr</b> Krypton 36
<b>Rb</b> Rubidium 37	<b>Sr</b> Strontium 38	<b>Y</b> Yttrium 39	<b>Zr</b> Zirconium 40	<b>Nb</b> Niobium 41	<b>Mo</b> Molybdenum 42	<b>Tc</b> Technetium 43	<b>Ru</b> Ruthenium 44	<b>Rh</b> Rhodium 45	<b>Pd</b> Palladium 46	<b>Ag</b> Silver 47	<b>Cd</b> Cadmium 48	<b>In</b> Indium 49	<b>Sn</b> Tin 50	<b>Sb</b> Antimony 51	<b>Te</b> Tellurium 52	<b>I</b> Iodine 53	<b>Xe</b> Xenon 54
<b>Cs</b> Cesium 55	<b>Ba</b> Barium 56	<b>La</b> Lanthanum 57	<b>Hf</b> Hafnium 72	<b>Ta</b> Tantalum 73	<b>W</b> Tungsten 74	<b>Re</b> Rhenium 75	<b>Os</b> Osmium 76	<b>Ir</b> Iridium 77	<b>Pt</b> Platinum 78	<b>Au</b> Gold 79	<b>Hg</b> Mercury 80	<b>Tl</b> Thallium 81	<b>Pb</b> Lead 82	<b>Bi</b> Bismuth 83	<b>Po</b> Polonium 84	<b>At</b> Astatine 85	<b>Rn</b> Radon 86
<b>Fr</b> Francium 87	<b>Ra</b> Radium 88	<b>Ac</b> Actinium 89															
		Lanthanides															
		<b>Ce</b> Cerium 58	<b>Pr</b> Praseodymium 59	<b>Nd</b> Neodymium 60	<b>Pm</b> Promethium 61	<b>Sm</b> Samarium 62	<b>Eu</b> Europium 63	<b>Gd</b> Gadolinium 64	<b>Tb</b> Terbium 65	<b>Dy</b> Dysprosium 66	<b>Ho</b> Holmium 67	<b>Er</b> Erbium 68	<b>Tm</b> Thulium 69	<b>Yb</b> Ytterbium 70	<b>Lu</b> Lutetium 71		
		<b>Th</b> Thorium 90	<b>Pa</b> Protactinium 91	<b>U</b> Uranium 92	<b>Np</b> Neptunium 93	<b>Pu</b> Plutonium 94	<b>Am</b> Americium 95	<b>Cm</b> Curium 96	<b>Bk</b> Berkelium 97	<b>Cf</b> Californium 98	<b>Es</b> Einsteinium 99	<b>Fm</b> Fermium 100	<b>Md</b> Mendelevium 101	<b>No</b> Nobelium 102	<b>Lr</b> Lawrencium 103		

**Table 1**  
Proton n.m.r chemical shift data

Type of proton	$\delta/\text{ppm}$
$\text{RCH}_3$	0.7–1.2
$\text{R}_2\text{CH}_2$	1.2–1.4
$\text{R}_3\text{CH}$	1.4–1.6
$\text{RCOCH}_3$	2.1–2.6
$\text{ROCH}_3$	3.1–3.9
$\text{RCOOCH}_3$	3.7–4.1
$\text{ROH}$	0.5–5.0

**Table 2**  
Infra-red absorption data

Bond	Wavenumber/ $\text{cm}^{-1}$
$\text{C—H}$	2850–3300
$\text{C—C}$	750–1100
$\text{C=C}$	1620–1680
$\text{C=O}$	1680–1750
$\text{C—O}$	1000–1300
$\text{O—H}$ (alcohols)	3230–3550
$\text{O—H}$ (acids)	2500–3000

## SECTION A

Answer **all** questions in the spaces provided.

- 1 (a) The initial rate of the reaction between substances **P** and **Q** was measured in a series of experiments and the following rate equation was deduced.

$$\text{rate} = k[\mathbf{P}]^2[\mathbf{Q}]$$

- (i) Complete the table of data below for the reaction between **P** and **Q**.

Experiment	Initial $[\mathbf{P}]/\text{mol dm}^{-3}$	Initial $[\mathbf{Q}]/\text{mol dm}^{-3}$	Initial rate/ $\text{mol dm}^{-3}\text{s}^{-1}$
1	0.20	0.30	$4.8 \times 10^{-3}$
2	0.10	0.10	
3	0.40		$9.6 \times 10^{-3}$
4		0.60	$19.2 \times 10^{-3}$

- (ii) Using the data from experiment 1, calculate a value for the rate constant,  $k$ , and deduce its units.

.....

.....

.....

.....

.....

(6 marks)

- (b) What change in the reaction conditions would cause the value of the rate constant to change?

.....

(1 mark)

- 2 Nitrogen dioxide dissociates according to the following equation.



When 21.3 g of nitrogen dioxide were heated to a constant temperature,  $T$ , in a flask of volume  $11.5 \text{ dm}^3$ , an equilibrium mixture was formed which contained 7.04 g of oxygen.

- (a) (i) Calculate the number of moles of oxygen present in this equilibrium mixture and deduce the number of moles of nitrogen monoxide also present in this equilibrium mixture.

*Number of moles of O<sub>2</sub> at equilibrium* .....

.....

*Number of moles of NO at equilibrium* .....

- (ii) Calculate the number of moles in the original 21.3 g of nitrogen dioxide and hence calculate the number of moles of nitrogen dioxide present in this equilibrium mixture.

*Original number of moles of NO<sub>2</sub>* .....

.....

*Number of moles of NO<sub>2</sub> at equilibrium* .....

.....

(4 marks)

- (b) Write an expression for the equilibrium constant,  $K_c$ , for this reaction. Calculate the value of this constant at temperature  $T$  and give its units.

*Expression for  $K_c$*  .....

.....

*Calculation* .....

.....

.....

.....

.....

(4 marks)

- (c) The total number of moles of gas in the flask is 0.683. Use the ideal gas equation to determine the temperature  $T$  at which the total pressure in the flask is  $3.30 \times 10^5$  Pa. (The gas constant  $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$ )

.....  
.....  
.....  
.....  
.....

(3 marks)

- (d) State the effect on the equilibrium yield of oxygen and on the value of  $K_c$  when the same mass of nitrogen dioxide is heated to the same temperature  $T$ , but in a different flask of greater volume.

Yield of oxygen .....


Value of  $K_c$  .....

(2 marks)

13

**TURN OVER FOR THE NEXT QUESTION**

- 3 The sketch below shows the change in pH when a  $0.200 \text{ mol dm}^{-3}$  solution of sodium hydroxide is added from a burette to  $25.0 \text{ cm}^3$  of a  $0.150 \text{ mol dm}^{-3}$  solution of the weak acid HA at  $25^\circ\text{C}$ .



- (a) The volume of sodium hydroxide solution added at the equivalence point is  $x \text{ cm}^3$ . Calculate the value of  $x$ .

.....  
 .....  
 .....

(2 marks)

- (b) (i) Define the term pH.

.....

- (ii) The pH at the equivalence point is  $c$ . Suggest a value for  $c$ .

.....

- (iii) Identify a suitable indicator for detecting the equivalence point of the titration.

.....

(3 marks)

(c) The value of  $K_a$  for the weak acid HA at  $25^\circ\text{C}$  is  $2.75 \times 10^{-5} \text{ mol dm}^{-3}$ .

(i) Explain the term *weak* as applied to the acid HA.

.....

(ii) Write an expression for  $K_a$  for the acid HA.

.....

.....

(iii) Calculate the pH of the  $0.150 \text{ mol dm}^{-3}$  solution of acid HA before any sodium hydroxide is added, i.e. the pH at point *a*.

.....

.....

.....

.....

.....

(5 marks)

(d) Calculate the pH of the solution formed when  $x/2 \text{ cm}^3$  of the  $0.200 \text{ mol dm}^{-3}$  solution of sodium hydroxide are added to  $25.0 \text{ cm}^3$  of the  $0.150 \text{ mol dm}^{-3}$  solution of HA, i.e. the pH at point *b*.

.....

.....


.....

(3 marks)


**TURN OVER FOR THE NEXT QUESTION**



4 The structures of the amino acids *alanine* and *glycine* are shown below.



*alanine*



*glycine*

(a) Give the systematic name for *alanine*.

.....  
(1 mark)

(b) *Alanine* exists as a pair of stereoisomers.

(i) Explain the meaning of the term *stereoisomers*.

.....  
.....  
.....

(ii) State how you could distinguish between the stereoisomers.

.....  
.....  
.....

(4 marks)

(c) Give the structural formula of the species formed by *glycine* at pH 14.

(1 mark)

- (d) When two amino acids react together, a dipeptide is formed. Give the structural formulae of the **two** dipeptides which are formed when *alanine* and *glycine* react together.

*Dipeptide 1*

*Dipeptide 2*


(2 marks)

- (e) Give the structural formula of the organic compound formed when *glycine* reacts with methanol in the presence of a small amount of concentrated sulphuric acid.

(1 mark)

9

5 The proton n.m.r. spectrum of compound **X** is shown below.



Compound **X**,  $C_7H_{12}O_3$ , contains both a ketone and an ester functional group. The measured integration trace for the peaks in the n.m.r. spectrum of **X** gives the ratio shown in the table below.

Chemical shift, $\delta$ /ppm	4.13	2.76	2.57	2.20	1.26
Integration ratio	0.8	0.8	0.8	1.2	1.2

Refer to the spectrum, the information given above and the data on the reverse of the Periodic Table provided to answer the following questions.

- (a) How many different types of proton are present in compound **X**?

.....  
(1 mark)

- (b) What is the whole-number ratio of each type of proton in compound **X**?

.....  
(1 mark)

- (c) Draw the part of the structure of **X** which can be deduced from the presence of the peak at  $\delta$ 2.20.

.....  
(1 mark)

- (d) The peaks at  $\delta$ 4.13 and  $\delta$ 1.26 arise from the presence of an alkyl group. Identify the group and explain the splitting pattern.

*Alkyl group* .....

*Explanation* .....

.....  
.....  
(3 marks)

- (e) Draw the part of the structure of **X** which can be deduced from the splitting of the peaks at  $\delta$ 2.76 and  $\delta$ 2.57.

.....  
(1 mark)

- (f) Deduce the structure of compound **X**.

.....  
(2 marks)

6 (a) Methylamine is a weak Brønsted–Lowry base and can be used in aqueous solution with one other substance to prepare a basic buffer.

(i) Explain the term *Brønsted–Lowry base* and write an equation for the reaction of methylamine with water to produce an alkaline solution.

*Brønsted–Lowry base* .....

*Equation* .....

(ii) Suggest a substance that could be added to aqueous methylamine to produce a basic buffer.

.....

(iii) Explain how the buffer solution in part (a)(ii) is able to resist a change in pH when a small amount of sodium hydroxide is added.

.....

.....

.....

(5 marks)

(b) Explain why methylamine is a stronger base than ammonia.

.....

.....

.....

(2 marks)

(c) A cation is formed when methylamine reacts with a large excess of bromoethane. Name the mechanism involved in the reaction and draw the structure of the cation formed.

*Name of mechanism* .....


*Structure*

(2 marks)

## SECTION B

Answer **both** questions in the space provided on pages 16 to 20 of this booklet.




7 A possible synthesis of 1,4-diaminobenzene is shown below.



- (a) Identify a suitable reagent or combination of reagents for Step 1. Name and outline a mechanism for the reaction. (6 marks)
- (b) Identify a suitable reagent or combination of reagents for Step 2. Name and outline a mechanism for the reaction. (6 marks)
- (c) Identify a suitable reagent or combination of reagents for Step 4. Draw the repeating unit of the polymer formed by reaction of 1,4-diaminobenzene with pentanedioic acid. (3 marks)

**TURN OVER FOR THE NEXT QUESTION**

- 8 (a) Describe, by giving reagents and stating observations, how you could distinguish between the compounds in the following pairs using a simple test-tube reaction for each pair.



(8 marks)

- (b) State how compounds **E** and **F** in part (a)(iii) above could be distinguished by their infra-red spectra, without using the fingerprint region. Explain how *fingerprinting* is used to identify a compound. (3 marks)
- (c) Suggest the structure of the fragment responsible for the major peak in the mass spectrum of compound **E** and state its  $m/z$  value. Write an equation showing the formation of this fragment from the molecular ion. (4 marks)

### END OF QUESTIONS

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....