

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

Leave blank

General Certificate of Education
June 2002
Advanced Level Examination



CHEMISTRY **CHM4**
Unit 4 Further Physical and Organic Chemistry

Wednesday 19 June 2002 Afternoon Session

In addition to this paper you will require:
the AQA Periodic Table (Reference CHEM/PT/EX);
a calculator.

Time allowed: 1 hour 30 minutes

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

Information

- The maximum mark for this paper is 90.
- Mark allocations are shown in brackets.
- This paper carries 15 per cent of the total marks for Advanced Level.
- You are expected to use a calculator where appropriate.
- The following data may be required.
Gas constant $R = 8.31 \text{ J mol}^{-1} \text{ K}^{-1}$
- Your answers to questions in **Section B** should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

Advice


- You are advised to spend about 1 hour on **Section A** and about 30 minutes on **Section B**.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
7			
8			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

SECTION A

Answer **all** the questions in the spaces provided.

- 1** Iodine and propanone react in acid solution according to the equation



The rate equation for the reaction is found to be

$$\text{rate} = k [\text{CH}_3\text{COCH}_3][\text{H}^+]$$

- (a) Deduce the order of reaction with respect to iodine and the overall order of reaction.

Order with respect to iodine

Overall order

(2 marks)

- (b) At the start of the experiment, the rate of reaction was found to be $2.00 \times 10^{-5} \text{ mol dm}^{-3} \text{ s}^{-1}$ when the concentrations of the reactants were as shown below.

Reactant	Concentration/mol dm ⁻³
CH ₃ COCH ₃	1.50
I ₂	2.00×10^{-2}
H ⁺	3.00×10^{-2}

Use these data to calculate a value for the rate constant and deduce its units.

Rate constant

.....

.....

Units

(3 marks)

- (c) How can you tell that H⁺ acts as a catalyst in this reaction?

.....

.....

.....

(2 marks)

- (d) Calculate the initial rate of reaction if the experiment were to be repeated at the same temperature and with the same concentrations of iodine and propanone as in part (b) but at a pH of 1.25

.....

.....

.....

.....

.....


.....

(3 marks)

$\frac{\quad}{10}$

TURN OVER FOR THE NEXT QUESTION

- 2 Acid **X** reacts with methanol to form ester **Y** according to the following equation.



A mixture of 0.25 mol of **X** and 0.34 mol of methanol was left to reach equilibrium in the presence of a small amount of concentrated sulphuric acid. The equilibrium mixture thus formed contained 0.13 mol of **Y** in a total volume of $V \text{ dm}^3$.

- (a) Name **X**.

.....
(1 mark)

- (b) Using **X** to represent the acid and **Y** to represent the ester, write an expression for the equilibrium constant, K_c , for this reaction.

(1 mark)

- (c) Calculate the number of moles of **X**, the number of moles of methanol and the number of moles of water in the equilibrium mixture.

Moles of X

Moles of methanol

Moles of water

(3 marks)

- (d) State why the volume V need not be known in calculating the value of K_c for the reaction.

.....

.....

(1 mark)

- (e) Calculate the value of K_c for this reaction and deduce its units.

Calculation

.....

.....

.....

Units of K_c

.....

(3 marks)

- (f) State the effect, if any, of increasing the temperature on the value of K_c

.....

(1 mark)

10

TURN OVER FOR THE NEXT QUESTION

3 Ethylbenzene is made by the reaction shown below.



(a) Identify two other substances required as catalysts in this preparation.

Substance 1

Substance 2

(2 marks)

(b) Write an equation for the reaction of these two substances with ethene to form the reactive intermediate involved in the formation of ethylbenzene.

.....

(1 mark)

(c) Name and outline a mechanism for the reaction between this reactive intermediate and benzene.

Name of mechanism

Mechanism

(4 marks)

(d) Draw the structure of the product formed in a similar reaction between benzene and cyclohexene.

(1 mark)

- (e) Ethylbenzene is used to make phenylethene which can be polymerised to form poly(phenylethene). Name this type of polymerisation and draw the structure of the repeating unit in the polymer.

Type of polymerisation


Repeating unit

(2 marks)

10

TURN OVER FOR THE NEXT QUESTION

4 Three sections of the proton n.m.r. spectrum of $\text{CH}_3\text{CHClCOOH}$ are shown below.



(a) Name the compound $\text{CH}_3\text{CHClCOOH}$

.....
(1 mark)

(b) Explain the splitting patterns in the peaks at $\delta 1.72$ and $\delta 4.44$

.....

(2 marks)

(c) Predict the splitting pattern that would be seen in the proton n.m.r. spectrum of the isomeric compound $\text{ClCH}_2\text{CH}_2\text{COOH}$


.....

(1 mark)

- (d) The amino acid *alanine* is formed by the reaction of $\text{CH}_3\text{CHClCOOH}$ with an excess of ammonia. The mechanism is nucleophilic substitution. Outline this mechanism, showing clearly the structure of *alanine*.

(5 marks)

- (e) The amino acid *lysine* has the structure



Draw structures to show the product formed in each case when lysine reacts with

- (i) an excess of aqueous HCl,
- (ii) an excess of aqueous NaOH,
- (iii) another molecule of lysine.

(3 marks)

- 5 (a) **P**, **Q** and **R** have the molecular formula C_6H_{12}
 All three are branched-chain molecules and none is cyclic.
P can represent a pair of optical isomers.
Q can represent a pair of geometrical isomers.
R can represent another pair of geometrical isomers different from **Q**.

Draw one possible structure for one of the isomers of each of **P**, **Q** and **R**.

Structure of P

Structure of Q

Structure of R

(3 marks)

- (b) Butanone reacts with reagent **S** to form compound **T** which exists as a racemic mixture.
 Dehydration of **T** forms **U**, C_5H_7N , which can represent a pair of geometrical isomers.

- (i) State the meaning of the term *racemic mixture* and suggest why such a mixture is formed in this reaction.

Racemic mixture

.....

Explanation.....

.....

.....

- (ii) Identify reagent **S**, and draw a structural formula for each of **T** and **U**.

Reagent S

Compound T

Compound U



- 6 (a) Draw the structure of ethyl propanoate.

(1 mark)

- (b) Name and outline a mechanism for the formation of ethyl propanoate from propanoyl chloride and ethanol.

Name of mechanism

Mechanism

(5 marks)

- (c) The mass spectrum of ethyl propanoate contains a major peak at $m/z = 57$. Write an equation showing the fragmentation of the molecular ion to form the species responsible for the peak at $m/z = 57$. Show the structure of this species in your answer.

.....
(2 marks)

- (d) Draw the structure of another ester which is an isomer of ethyl propanoate and which gives a major peak at $m/z = 71$ in its mass spectrum.


(1 mark)

9

SECTION B

Answer **both** the questions below in the space provided on pages 12 to 16 of this booklet.

- 7 (a) Compound **C**, $\text{H}_2\text{N}(\text{CH}_2)_4\text{NH}_2$, can be synthesised from ethene in three steps as shown below.



Name compound **C** and draw a structure for each of compounds **A** and **B**.
State the reagent(s) required for each step and name the type of reaction involved in the conversion of **B** into **C**. (7 marks)

- (b) Draw the repeating unit of the polyamide formed when **C** reacts with hexanedioic acid. Discuss the interactions between the chains of the polyamide. (4 marks)
- (c) Explain why polyamides are degraded by sodium hydroxide whereas polymers such as poly(ethene) are not. (3 marks)

- 8 A $0.210 \text{ mol dm}^{-3}$ solution of potassium hydroxide was added from a burette to 25.0 cm^3 of a $0.160 \text{ mol dm}^{-3}$ solution of ethanoic acid in a conical flask. Given that the value of the acid dissociation constant, K_a , for ethanoic acid is $1.74 \times 10^{-5} \text{ mol dm}^{-3}$, calculate the pH at 25°C of the solution in the conical flask at the following three points:

- before any potassium hydroxide had been added;
after 8.0 cm^3 of potassium hydroxide solution had been added;
after 40.0 cm^3 of potassium hydroxide solution had been added. (16 marks)

END OF QUESTIONS

.....

.....

.....

.....

.....

.....

Handwriting practice area with 25 horizontal dotted lines.

Handwriting practice area with 25 horizontal dotted lines.

Handwriting practice area with 25 horizontal dotted lines.

