

GCE

Chemistry

Unit CHM5

Copyright © 2003 AQA and its licensors. All rights reserved.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334
Registered address: Addleshaw Booth & Co., Sovereign House, PO Box 8, Sovereign Street, Leeds LS1 1HQ
Kathleen Tattersall: *Director General*

SECTION A


Answer all questions in the spaces provided.

- 1 (a) At high temperatures, aluminium chloride exists in the vapour phase as the molecule AlCl_3 . On cooling, two molecules of AlCl_3 combine by co-ordinate bonding to form molecules of Al_2Cl_6 .

- (i) State the shape of the AlCl_3 molecule and give the bond angle.

Linked to a correct shape { Shape Trigonal planar (1)
 Bond angle 120° (1)

- (ii) Sketch the structure of Al_2Cl_6 and mark on your sketch the value of one of the bond angles.



- (iii) Explain how two AlCl_3 molecules are able to bond together.

Lone electron pair on chlorine (1)

Electron deficient Al (accepts lone pair) (1)
 (6 marks)

- (b) (i) Describe what is observed when anhydrous AlCl_3 is added to an excess of water. Identify the major aluminium-containing species formed and predict the pH of the final solution.

Mark Separately Observation Dissolves / exothermic reaction (1)


Major aluminium-containing species $[\text{Al}(\text{H}_2\text{O})_6]^{3+}$ (1)

pH of final solution 2/3/4 (1)

- (ii) Describe what you would observe when aqueous sodium carbonate is added to aqueous aluminium chloride. Write an equation for the reaction.

Mark Separately Observations Gas evolved (1)

..... white precipitate formed (1)



Species (1) Balanced (1)

- 2 (a) Write equations for the reactions which occur when the following compounds are added separately to water. In each case, predict the approximate pH of the solution formed when one mole of each compound is added to 1 dm³ of water.

Sodium oxide



pH of solution formed 14 (1)

Sulphur dioxide




pH of solution formed 1 - 3 (1)

(4 marks)

- (b) When silicon dioxide and carbon are heated in a stream of chlorine gas, silicon tetrachloride and carbon monoxide are formed.


- (i) Write an equation for this reaction.



- (ii) State what is observed when silicon tetrachloride is added to water. Write an equation for the reaction which occurs.

Observations (Gelatinous) {Solid Gel precipitate formed (1)}

Exothermic reaction or gas evolved (1)



- (iii) Explain, in terms of their structure and bonding, why silicon tetrachloride has a lower melting point than phosphorus pentachloride.

SiCl_4 (is molecular), weak vdw forces between molecules (1); PCl_5 is ionic(1); ion-ion attraction stronger than vdw forces (1)

(7 marks)

11

- 3 Use the data in the table below to answer the questions which follow.

Substance	$\text{Fe}_2\text{O}_3(\text{s})$	$\text{Fe}(\text{s})$	$\text{C}(\text{s})$	$\text{CO}(\text{g})$	$\text{CO}_2(\text{g})$
$\Delta H_f^\ominus/\text{kJ mol}^{-1}$	-824.2	0	0	-110.5	-393.5
$S^\ominus/\text{J K}^{-1}\text{mol}^{-1}$	87.4	27.3	5.7	197.6	213.6

- (a) The following equation shows one of the reactions which can occur in the extraction of iron.



- (i) Calculate the standard enthalpy change and the standard entropy change for this reaction.

Standard enthalpy change $\Delta H_R = \sum \Delta H_f^\ominus \text{products} - \sum \Delta H_f^\ominus \text{reactants}$
 or a cycle (1)

$$\Delta H_R = ([2 \times 0] + [3 \times -393.5]) - (-824.2 + [3 \times -110.5]) \quad (1)$$

$$= -24.8 \quad (\text{kJ mol}^{-1}) \quad (1)$$

Allow + 24.8 max one

Standard entropy change $\Delta S = \sum S^\ominus \text{products} - \sum S^\ominus \text{reactants} \quad (1)$

$$\Delta S = ([2 \times 27.3] + [3 \times 213.6]) - (87.4 + [3 \times 197.6]) \quad (1)$$

$$= (54.6 + 640.8) - (87.4 + 592.8) \quad (1)$$

$$= 15.2 \quad (\text{J K}^{-1}\text{mol}^{-1}) \quad (1)$$

Allow - 15.2 max one

- (ii) Explain why this reaction is feasible at all temperatures.


$$\Delta G = \Delta H - T\Delta S \quad \text{or } \Delta S \text{ positive or correct calc.} \quad (1)$$

ΔH is negative and $-T\Delta S$ is negative (1)

{ Hence ΔG is always negative (1)

or feasible when $\Delta G \leq 0$ (9 marks)

- (b) The reaction shown by the following equation can also occur in the extraction of iron.



The standard entropy change, ΔS^\ominus , for this reaction is $+542.6 \text{ J K}^{-1} \text{ mol}^{-1}$

Use this information to calculate the temperature at which this reaction becomes feasible.

$$\Delta G = 0 = \Delta H - T\Delta S \quad \text{Hence } \Delta H = T\Delta S \quad (1)$$

Penalise missing 1000 by one mark

$$T = \Delta H / \Delta S = 492.7 \times 1000 / 542.6 \quad (1)$$

$$= 908 \text{ K} \quad (1)$$

(3 marks)

- (c) Calculate the temperature at which the standard free-energy change, ΔG^\ominus , has the same value for the reactions in parts (a) and (b).

$$\Delta G_{(b)} = \Delta G_{(a)}$$

$$(492.7 \times 10^3 - T \times 542.6) = (-24.8 \times 10^3 - T \times 15.2) \quad (1)$$

$$517.5 \times 10^3 = 527.4 \text{ T} \quad (1)$$

$$T = 982 \text{ K} \quad \text{Allow } 980 - 982 \quad (1)$$

(3 marks)

15

TURN OVER FOR THE NEXT QUESTION

- 4 Use the standard electrode potential data in the table below to answer the questions which follow.

	E^\ominus/V
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ce}^{3+}(\text{aq})$	+1.70
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	+1.51
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$	+1.36
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{SO}_3(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.17

- (a) Name the standard reference electrode against which all other electrode potentials are measured.

(Standard) hydrogen (electrode) (1)
(1 mark)

- (b) When the standard electrode potential for $\text{Fe}^{3+}(\text{aq})/\text{Fe}^{2+}(\text{aq})$ is measured, a platinum electrode is required.

- (i) What is the function of the platinum electrode?

To allow transfer of electrons/provide a reaction surface (1)

- (ii) What are the standard conditions which apply to $\text{Fe}^{3+}(\text{aq})/\text{Fe}^{2+}(\text{aq})$ when measuring this potential?

298 K (1)

Both $\text{Fe}^{3+}(\text{aq})$ and $\text{Fe}^{2+}(\text{aq})$ have a concentration QOL

of 1 mol dm^{-3} (1) OR $[\text{H}^+] = 1\text{ mol dm}^{-3}$

NOT zero current

or 100 kPa


(3 marks)

- (c) The cell represented below was set up under standard conditions.



Calculate the e.m.f. of this cell and write an equation for the spontaneous cell reaction.

Cell e.m.f. +1.34 V (1)



Correct species/order (1)

..... Balance d and cancelled (1)

Allow one for $2\text{MnO}_4^- + 5\text{H}_2\text{SO}_3 \rightarrow 2\text{Mn}^{2+} + 5\text{SO}_4^{2-}$ (3 marks)

- (d) (i) Which one of the species given in the table is the strongest oxidising agent?

..... $\text{Ce}^{4+}(\text{aq})$ (1)

- (ii) Which of the species in the table could convert $\text{Fe}^{2+}(\text{aq})$ into $\text{Fe}^{3+}(\text{aq})$ but could not convert $\text{Mn}^{2+}(\text{aq})$ into $\text{MnO}_4^-(\text{aq})$?

..... $\text{VO}_2^+(\text{aq})$ (1) ; Cl_2 (1)

Penalise additional answers to zero (3 marks)

- (e) Use data from the table of standard electrode potentials to deduce the cell which would have a standard e.m.f. of 0.93 V. Represent this cell using the convention shown in part (c).




Correct species (1) (2 marks)

Correct order (1) (Deduct one mark for each error)


12

TURN OVER FOR THE NEXT QUESTION


- 5 (a) Give one example of a bidentate ligand.




- (b) Give one example of a linear complex ion formed by a transition metal.




- (c) Write an equation for a substitution reaction in which the complete replacement of ligands in a complex ion occurs with a change in both the co-ordination number and the overall charge of the complex ion.



- (d) Write an equation for a substitution reaction in which the complete replacement of ligands in a complex ion occurs without a change in either the co-ordination number or the overall charge of the complex ion.



- (e) When a solution containing $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ ions is treated with a solution containing EDTA^{4-} ions, a more stable complex is formed. Write an equation for this reaction and explain why the complex is more stable.



6
9


Section B Question 6 Answers

- (a) $\text{Cu} \rightarrow \text{Cu}^{2+} + 2\text{e}^-$ (1)
 $\text{NO}_3^- + 2\text{H}^+ + \text{e}^- \rightarrow \text{H}_2\text{O} + \text{NO}_2$ (1)
-
- (1) 3
- or $\text{Cu} + 2\text{HNO}_3 + 2\text{H}^+ \rightarrow \text{Cu}^{2+} + 2\text{H}_2\text{O} + 2\text{NO}_2$
- (b) Moles Cu = $1.25/63.5 = 0.0197$ to 0.020 (1)
Moles NO = $2/3 \times 0.0197 = 0.0131$ (1)
 $\text{PV} = \text{nRT}$ (1)
 $V = \text{nRT}/P = 0.0131 \times 8.31 \times 330 / 98000$ (1)
- (Allow "n" consequentially to moles NO but not to moles Cu)
= 3.67×10^{-4} (1)
units m³ (1) 6
- or equivalent
- (c) Yellow-green copper species $[\text{CuCl}_4]^{2-}$ (1)
Equation; $\text{Cu} + [\text{CuCl}_4]^{2-} + 4\text{Cl}^- \rightarrow 2[\text{CuCl}_4]^{3-}$ Species (1)
Balance (1)
- Copper acts as a reducing agent (1)
- Not coloured as d sub-shell full (1)
Copper in CuCl is in oxidation state +1 or
No absorption of light as electron transitions not possible (1)

Section B Question 7 Answers

- (a) reactants brought together/ increased concentration on surface
or increased collision frequency (1)
reactants must be correctly orientated (1)
reaction on the surface (1)
products desorbed (1)
example of a catalysed reaction (not a named process) (1)
a suitable catalyst for this reaction (1)
(Penalise incorrect second reactions and catalysts))
If adsorption too weak reactants not brought together (1)
e.g. silver (1)
If adsorption too strong products not desorbed (1)
e.g. tungsten (1) Max 8

(b) Equations



Method

- Titrate measured volume solution against $\text{K}_2\text{Cr}_2\text{O}_7$ (1)
Reduce same volume solution with zinc (1)
Filter off excess zinc (1)
Titrate for total Fe^{n+} using $\text{K}_2\text{Cr}_2\text{O}_7$ (1)
Percentage $\text{Fe}^{3+} = 100 \times (\text{titre 2} - \text{titre 1}) / \text{titre 2}$
or equivalent (1) 7

Section B Question 8 Answers.

(a) Identity of X; 2-methylpropene (1)

Absorption at 1650 cm^{-1} indicates an alkene present (1)
[OR a chemical answer e.g. $\text{Br}_2(\text{aq})$ brown to colourless]


2

(b) Reagents


Step 1 KOH (Allow NaOH) (1) Alcoholic (1) Warm (1)
(Only allow solvent and warm if reagent correct)

Step 2 HBr (1)

Mechanism $A \rightarrow X$



Mechanism $X \rightarrow B$



11

(c) A gives three peaks (1)

B gives one peak (1)

Allow one for


"A has more peaks than B" when no number of peaks given

2


Section B Question 9 Answers



Ammonia reacts as a base or nucleophile (1)



Ammonia acts as a ligand (1)



(Allow one mark if second product is HCl)

Ammonia behaves as a nucleophile (1)

7

(b) Pentane is non-polar (1)

van der Waals forces between molecules (1)

Butanone has a polar C=O bond (1)

Dipole-dipole attraction between molecules (1)

Propanoic acid has very polar O-H groups (1)

Hydrogen bonding between molecules (1)

$\text{vdw} < \text{dipole - dipole attraction} < \text{hydrogen bonding}$ (1)

Link to energy needed to separate molecules (1)

(Last mark only allowed when clearly stated and all three interactions are correct)

8