

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

Leave blank

General Certificate of Education
January 2003
Advanced Level Examination



CHEMISTRY **CHM5**
Unit 5 Thermodynamics and Further Inorganic Chemistry
(including Synoptic Assessment)

Friday 24 January 2003 Afternoon Session

In addition to this paper you will require: a calculator.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
5			
6			
7			
8			
9			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Time allowed: 2 hours

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in **Section A** and **Section B** in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

Information

- The maximum mark for this paper is 120.
- Mark allocations are shown in brackets.
- This paper carries 20 per cent of the total marks for Advanced Level.
- You are expected to use a calculator where appropriate.
- The following data may be required.
Gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$
- Your answers to questions in **Section B** should be written in continuous prose, where appropriate. You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

Advice

- You are advised to spend about 1 hour on **Section A** and about 1 hour on **Section B**.

SECTION A

Answer **all** questions in the spaces provided.

- 1 (a) At high temperatures, aluminium chloride exists in the vapour phase as the molecule AlCl_3 . On cooling, two molecules of AlCl_3 combine by co-ordinate bonding to form molecules of Al_2Cl_6 .

- (i) State the shape of the AlCl_3 molecule and give the bond angle.

Shape

Bond angle

- (ii) Sketch the structure of Al_2Cl_6 and mark on your sketch the value of **one** of the bond angles.

- (iii) Explain how two AlCl_3 molecules are able to bond together.

.....
.....

(6 marks)

- (b) (i) Describe what is observed when anhydrous AlCl_3 is added to an excess of water. Identify the major aluminium-containing species formed and predict the pH of the final solution.

Observation

Major aluminium-containing species

pH of final solution

- (ii) Describe what you would observe when aqueous sodium carbonate is added to aqueous aluminium chloride. Write an equation for the reaction.

Observations

.....

Equation

.....



The Periodic Table of the Elements

■ The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

		I	II	III	IV	V	VI	VII	0							
H Hydrogen 1									He Helium 2							
Li Lithium 3	Be Beryllium 4		Li Lithium 3					F Fluorine 9	Ne Neon 10							
Na Sodium 11	Mg Magnesium 12							Cl Chlorine 17	Ar Argon 18							
K Potassium 19	Ca Calcium 20	V Vanadium 23	Ti Titanium 22	Cr Chromium 24	Mn Manganese 25	Fe Iron 26	Co Cobalt 27	Ni Nickel 28	Cu Copper 29	Zn Zinc 30	Ga Gallium 31	Ge Germanium 32	As Arsenic 33	Se Selenium 34	Br Bromine 35	Kr Krypton 36
Rb Rubidium 37	Sr Strontium 38	Nb Niobium 41	Zr Zirconium 40	Mo Molybdenum 42	Tc Technetium 43	Ru Ruthenium 44	Rh Rhodium 45	Pd Palladium 46	Ag Silver 47	Cd Cadmium 48	In Indium 49	Sn Tin 50	Sb Antimony 51	Te Tellurium 52	I Iodine 53	Xe Xenon 54
Cs Caesium 55	Ba Barium 56	Ta Tantalum 73	Hf Hafnium 72	W Tungsten 74	Re Rhenium 75	Os Osmium 76	Ir Iridium 77	Pt Platinum 78	Au Gold 79	Hg Mercury 80	Tl Thallium 81	Pb Lead 82	Bi Bismuth 83	Po Polonium 84	At Astatine 85	Rn Radon 86
Ra Radium 88	Ac Actinium 89															

Ce Cerium 58	Pr Praseodymium 59	Nd Neodymium 60	Pm Promethium 61	Sm Samarium 62	Eu Europium 63	Gd Gadolinium 64	Tb Terbium 65	Dy Dysprosium 66	Ho Holmium 67	Er Erbium 68	Tm Thulium 69	Yb Ytterbium 70	Lu Lutetium 71
Th Thorium 90	Pa Protactinium 91	U Uranium 92	Np Neptunium 93	Pu Plutonium 94	Am Americium 95	Cm Curium 96	Bk Berkelium 97	Cf Californium 98	Es Einsteinium 99	Fm Fermium 100	Md Mendelevium 101	No Nobelium 102	Lr Lawrencium 103

88 – 71 Lanthanides

90 – 103 Actinides

Table 1
Proton n.m.r chemical shift data

Type of proton	δ/ppm
RCH_3	0.7–1.2
R_2CH_2	1.2–1.4
R_3CH	1.4–1.6
RCOCH_3	2.1–2.6
ROCH_3	3.1–3.9
RCOOCH_3	3.7–4.1
ROH	0.5–5.0

Table 2
Infra-red absorption data

Bond	Wavenumber/ cm^{-1}
C—H	2850–3300
C—C	750–1100
C=C	1620–1680
C=O	1680–1750
C—O	1000–1300
O—H (alcohols)	3230–3550
O—H (acids)	2500–3000

- 2 (a) Write equations for the reactions which occur when the following compounds are added separately to water. In each case, predict the approximate pH of the solution formed when one mole of each compound is added to 1 dm³ of water.

Sodium oxide

Equation

pH of solution formed

Sulphur dioxide

Equation

pH of solution formed

(4 marks)

- (b) When silicon dioxide and carbon are heated in a stream of chlorine gas, silicon tetrachloride and carbon monoxide are formed.

- (i) Write an equation for this reaction.

.....

- (ii) State what is observed when silicon tetrachloride is added to water. Write an equation for the reaction which occurs.

Observations

.....

Equation

- (iii) Explain, in terms of their structure and bonding, why silicon tetrachloride has a lower melting point than phosphorus pentachloride.

.....

.....

.....


.....

(7 marks)

3 Use the data in the table below to answer the questions which follow.

Substance	Fe ₂ O ₃ (s)	Fe(s)	C(s)	CO(g)	CO ₂ (g)
$\Delta H_f^\ominus / \text{kJ mol}^{-1}$	-824.2	0	0	-110.5	-393.5
$S^\ominus / \text{J K}^{-1} \text{mol}^{-1}$	87.4	27.3	5.7	197.6	213.6

- (a) The following equation shows one of the reactions which can occur in the extraction of iron.



- (i) Calculate the standard enthalpy change and the standard entropy change for this reaction.

Standard enthalpy change

.....

.....

.....

.....

.....

Standard entropy change

.....

.....

.....

.....

- (ii) Explain why this reaction is feasible at all temperatures.


.....

.....

.....

(9 marks)

- (b) The reaction shown by the following equation can also occur in the extraction of iron.



The standard entropy change, ΔS^\ominus , for this reaction is $+542.6 \text{ J K}^{-1} \text{ mol}^{-1}$

Use this information to calculate the temperature at which this reaction becomes feasible.

.....
.....
.....

(3 marks)

- (c) Calculate the temperature at which the standard free-energy change, ΔG^\ominus , has the same value for the reactions in parts (a) and (b).

.....
.....
.....
.....

(3 marks)

15

TURN OVER FOR THE NEXT QUESTION

- 4 Use the standard electrode potential data in the table below to answer the questions which follow.

	E^\ominus/V
$\text{Ce}^{4+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ce}^{3+}(\text{aq})$	+1.70
$\text{MnO}_4^-(\text{aq}) + 8\text{H}^+(\text{aq}) + 5\text{e}^- \rightleftharpoons \text{Mn}^{2+}(\text{aq}) + 4\text{H}_2\text{O}(\text{l})$	+1.51
$\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$	+1.36
$\text{VO}_2^+(\text{aq}) + 2\text{H}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{VO}^{2+}(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+1.00
$\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$	+0.77
$\text{SO}_4^{2-}(\text{aq}) + 4\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{SO}_3(\text{aq}) + \text{H}_2\text{O}(\text{l})$	+0.17

- (a) Name the standard reference electrode against which all other electrode potentials are measured.

.....
(1 mark)

- (b) When the standard electrode potential for $\text{Fe}^{3+}(\text{aq})/\text{Fe}^{2+}(\text{aq})$ is measured, a platinum electrode is required.

- (i) What is the function of the platinum electrode?

.....

- (ii) What are the standard conditions which apply to $\text{Fe}^{3+}(\text{aq})/\text{Fe}^{2+}(\text{aq})$ when measuring this potential?


.....

.....

.....

(3 marks)

- (c) The cell represented below was set up under standard conditions.



Calculate the e.m.f. of this cell and write an equation for the spontaneous cell reaction.

Cell e.m.f.

Equation

.....

.....

(3 marks)

- (d) (i) Which one of the species given in the table is the strongest oxidising agent?

.....

- (ii) Which of the species in the table could convert $\text{Fe}^{2+}(\text{aq})$ into $\text{Fe}^{3+}(\text{aq})$ but could not convert $\text{Mn}^{2+}(\text{aq})$ into $\text{MnO}_4^-(\text{aq})$?

.....

(3 marks)

- (e) Use data from the table of standard electrode potentials to deduce the cell which would have a standard e.m.f. of 0.93 V. Represent this cell using the convention shown in part (c).

.....

(2 marks)

12

TURN OVER FOR THE NEXT QUESTION

- 5 (a) Give **one** example of a bidentate ligand.

.....
(1 mark)

- (b) Give **one** example of a linear complex ion formed by a transition metal.

.....
(1 mark)

- (c) Write an equation for a substitution reaction in which the complete replacement of ligands in a complex ion occurs with a change in **both** the co-ordination number and the overall charge of the complex ion.

.....
(2 marks)

- (d) Write an equation for a substitution reaction in which the complete replacement of ligands in a complex ion occurs without a change in either the co-ordination number or the overall charge of the complex ion.

.....
(2 marks)

- (e) When a solution containing $[\text{Co}(\text{H}_2\text{O})_6]^{2+}$ ions is treated with a solution containing EDTA^{4-} ions, a more stable complex is formed. Write an equation for this reaction and explain why the complex is more stable.

Equation

Explanation

.....
(3 marks)


SECTION B

Answer **all** questions in the space provided on pages 13 to 20 of this booklet.

- 6 (a) Water, copper(II) ions and nitrogen dioxide are formed when copper metal reacts with concentrated nitric acid.

Write half-equations for the reactions occurring and use these to construct an overall ionic equation for the reaction. (3 marks)

- (b) When copper reacts with dilute nitric acid, gaseous nitrogen monoxide is formed as shown by the following equation.



Calculate the volume of nitrogen monoxide, measured at 330K and 98.0 kPa, which is formed when 1.25g of copper metal reacts completely with an excess of dilute nitric acid. (6 marks)

- (c) When copper(II) chloride dissolves in concentrated hydrochloric acid, a yellow-green copper-containing complex is formed.

When an excess of copper metal is added to this solution and the mixture is warmed, the complex species $[\text{CuCl}_4]^{3-}$ is formed.

When the solution containing the complex $[\text{CuCl}_4]^{3-}$ is poured into water, CuCl is formed as a white solid.

Identify the yellow-green copper-containing complex, write an equation for the reaction in which $[\text{CuCl}_4]^{3-}$ is formed and deduce the role of copper in this reaction.

Explain why CuCl is not coloured. (6 marks)


- 7 (a) "The strength of adsorption onto the active sites on the surface of a heterogeneous catalyst helps to determine the activity of the catalyst."

Explain how heterogeneous catalysts work, give **one** example of a reaction catalysed in this way and discuss why different catalysts have different activities. (8 marks)

- (b) Outline a plan of an experiment to determine the percentage of iron present as iron(III) in a solution containing $\text{Fe}^{3+}(\text{aq})$ and $\text{Fe}^{2+}(\text{aq})$ ions. You are provided with zinc, a standard solution of potassium dichromate(VI) and dilute sulphuric acid. Zinc can reduce $\text{Fe}^{3+}(\text{aq})$ to $\text{Fe}^{2+}(\text{aq})$.

Write equations for all the reactions that occur. Explain how you would use the zinc and how you would calculate the final answer. (7 marks)

- 8 The conversion of compound **A** into compound **B** can be achieved in two steps as shown below.



The intermediate compound, **X**, has an absorption at 1650 cm^{-1} in its infra-red spectrum.

- (a) Identify compound **X**. Explain your answer. (2 marks)
- (b) For each step in this conversion, give the reagents and essential conditions required and outline a mechanism. (11 marks)
- (c) Show how the number of peaks in their proton n.m.r. spectra would enable you to distinguish between compounds **A** and **B**. (2 marks)
- 9 (a) The lone pair of electrons on the nitrogen atom is involved in the separate reactions of ammonia with hydrogen chloride, silver chloride and ethanoyl chloride.
- Write equations for the reaction of ammonia with each of these compounds. State the role of ammonia in each of these reactions. (7 marks)
- (b) Explain, in terms of the forces between particles, why the following compounds, which have similar relative molecular masses, have different melting points.

Compound	Formula	M_r	Melting point/K
Pentane	$\text{CH}_3(\text{CH}_2)_3\text{CH}_3$	72	143
Butanone	$\text{CH}_3\text{CH}_2\text{COCH}_3$	72	187
Propanoic acid	$\text{CH}_3\text{CH}_2\text{CO}_2\text{H}$	74	252

(8 marks)

END OF QUESTIONS