

Surname						Other Names					
Centre Number						Candidate Number					
Candidate Signature											

For Examiner's Use

General Certificate of Education
June 2008
Advanced Level Examination



CHEMISTRY
Unit 6(b) Practical Examination

CHM6/P

Thursday 22 May 2008 9.00 am to 11.00 am

For this paper you must have

- a calculator.

Time allowed: 2 hours

Instructions

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Carry out **all three** exercises.
- Answer **all** questions.
- Answer questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want to be marked.
- Take careful note of all the instructions given in each exercise.
- The Periodic Table/Data Sheet is provided on pages 3 and 4. Detach this perforated sheet at the start of the examination.

Information

- You must **not** use note books and laboratory books.
- The maximum mark for this paper is 30.
- The skills which are being assessed are
Skill 1 Planning (8 marks)
Skill 2 Implementing (8 marks)
Skill 3 Analysing (8 marks)
Skill 4 Evaluating (6 marks)
- You will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary, where appropriate.

Advice

- You are advised to spend about 40 minutes on each of the three exercises.
- You are advised to carry out Exercise 1 first.

For Examiner's Use			
Number	Mark	Number	Mark
Skill 1			
Skill 2			
Skill 3			
Skill 4			
Total (Column 1) →			
Total (Column 2) →			
TOTAL			
Examiner's Initials			

This paper consists of the following.

Exercise 1	Implementing	Titration of a solution of iron(II) sulphate
Exercise 2	Analysing and Evaluating	The reaction between hydrogen peroxide and iodide ions
Exercise 3	Planning	Determination of the dissociation constant, K_a , of a weak acid.

An essential part of any practical work is to plan for the most efficient use of the time available. There is enough time to complete the exercises set provided that a sensible approach is used.

You are advised to spend approximately

40 minutes on Exercise 1
40 minutes on Exercise 2
40 minutes on Exercise 3

■ The atomic numbers and approximate relative atomic masses shown in the table are for use in the examination unless stated otherwise in an individual question.

www.theallpapers.com

Gas constant $R = 8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

Table 1
Proton n.m.r chemical shift data

Type of proton	δ/ppm
RCH_3	0.7–1.2
R_2CH_2	1.2–1.4
R_3CH	1.4–1.6
RCOCH_3	2.1–2.6
ROCH_3	3.1–3.9
RCOOCH_3	3.7–4.1
ROH	0.5–5.0

Table 2
Infra-red absorption data

Bond	Wavenumber/ cm^{-1}
C—H	2850–3300
C—C	750–1100
C=C	1620–1680
C=O	1680–1750
C—O	1000–1300
O—H (alcohols)	3230–3550
O—H (acids)	2500–3000

Exercise 1 Titration of a solution of iron(II) sulphate**Skill assessed** **Implementing** (8 marks)**Introduction**

You are provided with an aqueous solution of iron(II) sulphate of concentration approximately 0.1 mol dm^{-3} . Titrate this solution, after acidification, with the $0.0200 \text{ mol dm}^{-3}$ solution of potassium manganate(VII) provided.

Wear eye protection at all times.

Assume that all of the solutions are toxic and corrosive.

Procedure

- 1 Rinse the burette with the potassium manganate(VII) solution provided. Set up the burette and, using a funnel, fill it with the potassium manganate(VII) solution. Record the initial burette reading in the table below.
- 2 Rinse a pipette with the iron(II) sulphate solution provided. Using this pipette and a pipette filler, transfer 25.0 cm^3 of the iron(II) sulphate solution to a 250 cm^3 conical flask.
- 3 Using a measuring cylinder, transfer approximately 10 cm^3 of dilute sulphuric acid to the conical flask.
- 4 Add the potassium manganate(VII) solution from the burette until the mixture in the conical flask has a permanent pink colour. Record your final burette reading in the table below.
- 5 Rinse the conical flask with water and repeat the titration until you obtain **two** titres which are within 0.10 cm^3 of each other. (You should do no more than five titrations.)
Have one of your final burette readings checked by your supervisor.
- 6 Calculate and record the average titre.

Results


Final burette reading/ cm^3					
Initial burette reading/ cm^3					
Volume of potassium manganate(VII) solution used/ cm^3					
Tick the titres to be used in calculating the average titre					

Average titre = cm^3

For Examiner's use only					
M		C		P	
T		A			

Exercise 2 The reaction between hydrogen peroxide and iodide ions**Skills assessed** **Analysing** (8 marks) **and Evaluating** (6 marks)**Introduction**

Acidified hydrogen peroxide reacts with iodide ions to form iodine according to the following equation.



In an experiment to determine the order of the reaction with respect to iodide ions, measured amounts of sodium thiosulphate solution and starch solution are added to an acidified mixture of hydrogen peroxide and potassium iodide. The **initial rate** of this reaction is investigated by measuring the time taken to produce sufficient iodine to react with the thiosulphate ions present, and then produce a blue colour with starch solution.


A series of experiments is carried out, in which the concentration of iodide ions is varied, while keeping the concentrations of all of the other reagents the same. In each experiment the time taken for the reaction mixture to turn blue is recorded. The results obtained are used to determine the order of reaction with respect to iodide ions.

The rate of the reaction can be represented as (1/time), and the concentration of iodide ions can be represented by the volume of potassium iodide solution used.

A graph of $\log_{10} (1/\text{time})$ on the y axis against $\log_{10} (\text{volume of KI(aq)})$ is a straight line. The gradient of this straight line is equal to the order of the reaction with respect to iodide ions.

A set of results is given in the table below. The volumes of potassium iodide solution were measured using a measuring cylinder. The time taken for each mixture to turn blue was recorded on a stopclock graduated in seconds.

Expt.	Volume of KI(aq)/cm ³	$\log_{10} (\text{volume of KI(aq)})$	time/s	$\log_{10} \left(\frac{1}{\text{time}} \right)$
1	5	0.70	68	-1.83
2	8	0.90	45	-1.65
3	10	1.00	36	-1.56
4	15	1.18	25	-1.40
5	20	1.30	22	-1.34
6	25	1.40	16	-1.20



Analysis Full marks can only be scored if you show all of your working.

- 1 Use the results given in the table to plot a graph of $\log_{10} (1/\text{time})$ on the y axis against $\log_{10} (\text{volume of KI(aq)})$.

Draw a straight line of best fit on the graph, ignoring any anomalous points.

- 2 Determine the gradient of the line you have drawn.

.....

.....

.....

- 3 For the measuring cylinder and the clock, the maximum total errors are shown below. These errors take into account multiple measurements.

measuring cylinder	$\pm 0.5 \text{ cm}^3$
clock	$\pm 1 \text{ second}$

Estimate the maximum percentage error in using these pieces of apparatus in Experiment 3.
Hence calculate the maximum overall percentage error in Experiment 3.

.....

.....

.....

.....

Evaluation **Full marks can only be scored if you show all of your working.**

- 1 Consider your graph and comment on the results obtained from the experiments. Is your line of best fit good enough for you to deduce an order with confidence? Identify any anomalous results.

.....

.....

.....

.....

- 2 State **two** ways in which the method used in these experiments could be improved, other than by repeating the experiments. In each case explain why the accuracy of the experiment would be improved.

Improvement 1

.....

.....

Explanation

.....

.....

Improvement 2

.....

.....

Explanation

.....

.....

Exercise 3 Determining the dissociation constant, K_a , of a weak acid

Skill assessed **Planning** (8 marks)

Introduction

A suitable method for determining the dissociation constant, K_a , of a weak acid involves measuring the pH values of the solution obtained as aqueous alkali is added to a known volume of the aqueous acid, until the alkali is present in considerable excess.

A pH curve can be plotted from the results. This curve can be used to determine the volume of alkali needed at the equivalence point (end-point). When half of this volume of alkali has been added, the pH of the solution is equal to the pK_a of the acid.

Question

Describe how you could determine the dissociation constant, K_a , of the acid.

You are provided with a crystalline sample of the weak monoprotic acid HA and a $0.100 \text{ mol dm}^{-3}$ solution of sodium hydroxide. The M_r of the acid is 150 and the acid is soluble in water.

Your answer must include

- (a) the scale you would choose for the experiment and a calculation of the mass of acid you would use to prepare your solution.
- (b) a detailed description of the experiment you would carry out. You do **not** need to describe how you would prepare the acid solution.
- (c) a sketch of the pH curve and an explanation of how you would use this curve to determine the dissociation constant, K_a , of the acid.
- (d) details of potential hazards and the relevant safety precautions you would take.

END OF QUESTIONS

Handwriting practice area with 25 horizontal dotted lines.

Handwriting practice lines consisting of 25 horizontal dotted lines.

Handwriting practice lines consisting of 25 horizontal dotted lines.

Handwriting practice lines consisting of 25 horizontal dotted lines.

Handwriting practice lines consisting of 25 horizontal dotted lines.

This image shows a full page of a worksheet designed for handwriting practice. It features 20 evenly spaced, horizontal dashed lines across the entire width of the page. The background is plain white, and there are no margins, text, or other markings present.