



ASSESSMENT and  
QUALIFICATIONS  
ALLIANCE

---

# Mark scheme January 2004

## GCE

# Mathematics A

## Unit MAM2

Copyright © 2004 AQA and its licensors. All rights reserved.

## Key to mark scheme

<b>M</b>	mark is for	method
<b>m</b>	mark is dependent on one or more M marks and is for	method
<b>A</b>	mark is dependent on M or m mark and is for	accuracy
<b>B</b>	mark is independent of M or m marks and is for	method and accuracy
<b>E</b>	mark is for	explanation
<b>√ or ft or F</b>		follow through from previous incorrect result
<b>CAO</b>		correct answer only
<b>AWFW</b>		anything which falls within
<b>AWRT</b>		anything which rounds to
<b>AG</b>		answer given
<b>SC</b>		special case
<b>OE</b>		or equivalent
<b>A2,1</b>		2 or 1 (or 0) accuracy marks
<b>- x EE</b>		Deduct $x$ marks for each error
<b>NMS</b>		No method shown
<b>PI</b>		Perhaps implied
<b>c</b>		Candidate


## Abbreviations used in marking

<b>MC - <math>x</math></b>	deducted $x$ marks for miscopy
<b>MR - <math>x</math></b>	deducted $x$ marks for misread
<b>ISW</b>	ignored subsequent working
<b>BOD</b>	gave benefit of doubt
<b>WR</b>	work replaced by candidate


## Application of mark scheme



Correct answer without working	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise



Award method and accuracy marks as appropriate to an alternative solution using a correct method or partially correct method.

Q	Solution	Marks	Total	Comments
1 (a)	Impulse = area $= 2000 \times 5 + \frac{1}{2}(2000) \times 5$	M1	3	AG
	$= 15\,000 \text{ N s}$	M1		
		A1		
1 (b)	Change in momentum  = impulse $1000 v - 0 = 15000$	M1	2	Attempt at $ mv - mu  = \text{impulse from (a)}$  Must state units
	$v = 15 \text{ m s}^{-1}$	A1		
<b>Total</b>			<b>5</b>	
2 (a)		B1	1	
2 (b)	$F = mr\omega^2$ $= 0.01(0.4)\omega^2$ $= 0.004\omega^2$	M1	2	Formula quoted and attempt at use
		A1		
2 (c)	$F = \mu N$ $= 0.8(0.01g)$ $= 0.008g$ Limiting, so $0.004\omega^2 = 0.008g$ $\omega^2 = 2g$ $\omega = 4.43$	M1	4	Use of limiting fraction  or 0.0784 seen  Equates their expressions  AG
		A1		
<b>Total</b>			<b>7</b>	

Q	Solution	Marks	Total	Comments		
3 (a)	$EPE = \frac{1}{2}(50)(0.03)^2$	M1	2	Use of correct formula		
	$= 0.0225 \text{ J}$	A1				
(b)	KE + PE = EPE					
	$\frac{1}{2}mv^2 + mgh = EPE$	B1		KE or PE term correct		
	$\frac{1}{2}(0.02)v^2 + (0.02)(9.8)(0.03) = 0.0225$	M1 A1✓		Equation formed Equation correct; ft EPE		
(c)	Energy remains constant – No KE at end, no EPE at end		4	AG		
	$\therefore (0.02)(9.8)h = 0.0225$	M1A1				
	$\therefore h = 0.11 \text{ m}$ i.e. 11cm	A1✓			3	ft EPE      2 sig fig accuracy
	<b>Alternatives to part (c)</b> Energy remains constant KE at point of release = PE on reaching max height					
$\frac{1}{2}(0.02)(1.29)^2 = (0.02)gh$	(M1)					
$\therefore h = 0.849\dots$	(A1)					
$\therefore \text{height above initial position} = 0.849\dots + 3$ $= 0.11\text{m}$	(A1✓)		ft their $h$ value + 3			
<b>or</b> Use of $v^2 - u^2 = 2as$	(M1)					
$s = 0.849 \text{ distance} = 0.849 + 3 = 0.11$	(A1) (A1✓)		ft their $(s + 3)$ total			
<b>Total</b>			<b>9</b>			

Q	Solution	Marks	Total	Comments
4 (a)	5cm	B1	1	
(b)	$\Sigma mx = (\Sigma m)\bar{x}$ About PS: $6(m)+6(m)+3(1)=(2m+1)\bar{x}$ $\bar{x} = \frac{12m+3}{2m+1}$	M1A1 A1	3	M1 one side correct AG
(c)	 $\tan 45^\circ = \frac{\bar{x}}{5}$ $\Rightarrow \frac{12m+3}{2m+1} = 5$ $12m+3 = 10m+5$ $2m = 2$ $m = 1$	M1 A1 A1 m1 A1	5	Principle applied Equation correct – use of part (a) Substitute and $\tan 45^\circ = 1$ Solving - dependent CAO
<b>Total</b>			<b>9</b>	

Q	Solution	Marks	Total	Comments
5 (a)				
(i)	<p>At P, PE = 70g 5 cos 30° KE = 0</p> <p>At Q, KE = <math>\frac{1}{2}(70)v^2</math> PE = 0</p> <p>Conservation of energy</p> $\frac{1}{2}(70)v^2 = 70g 5 \cos 30^\circ$ $v^2 = 10g \cos 30^\circ$ $\Rightarrow v \approx 9.21 \text{ms}^{-1}$	B1  M1A1  A1	4	PE or KE term seen correct (Non zero)  Forming equation  AG
(ii)	 <p>Force towards O = <math>T - 70g \cos 30^\circ</math></p> <p>For circular motion, <math>F = ma</math></p> $\Rightarrow T - 70g \cos 30^\circ = \frac{mv^2}{r}$ $\Rightarrow T - 70g \cos 30^\circ = \frac{70(9.21)^2}{5}$ $\Rightarrow T \approx 1782.28 \dots$ $\Rightarrow T = 1780 \text{N}$	B1  M1A1  ml A1	5	$\pm (T - 70g \cos 30^\circ)$ or $\frac{mv^2}{r}$ evaluated  Form equation – Res.force = $\frac{mv^2}{r}$  Dependent – substitute and rearrange  AWRT 1780

Q	Solution	Marks	Total	Comments
5 (b)	 <p>Vertically, <math>u = 9.21 \sin 30^\circ</math></p> <p><math>v = 0</math>  <math>a = -9.8</math>  <math>s = ?</math></p> <p>Using <math>v^2 - u^2 = 2as</math> :</p> $\left( \frac{9.21 \sin 30^\circ}{2(9.8)} \right) = 5$ <p><math>s \approx 1.08</math>  Approx 1 metre</p>	M1 A1		$v^2 - u^2 = 2as$ seen Initial vertical velocity component seen = $9.21 \sin 30^\circ$
(c)	Height of a man significant to length if rope/distances involved.			
	Air resistance would reduce speed/height.	B1	1	Must see 1. ... Comment that indicates effect of assumption
<b>Total</b>			<b>14</b>	
6	 <p>(a) Restitution: <math>v_1 - v_2 = eu</math></p> <p>Momentum: <math>mv_1 + mv_2 = mu</math></p> <p><math>v_1 - v_2 = eu</math> (1)</p> <p><math>v_1 + v_2 = u</math> (2)</p> <p>(1)+(2) <math>2v_1 = u(1+e)</math></p> <p><math>v_1 = \frac{u}{2}(1+e)</math></p> <p>(2)-(1) <math>2v_2 = u(1-e)</math></p> <p><math>v_2 = \frac{u}{2}(1-e)</math></p>	M1 M1 A1 M1 A1 B1 $\checkmark$		Attempt at restitution Attempt at momentum Both correct  AG

Q	Solution	Marks	Total	Comments
6 (b)	Speed = $\frac{2}{3} \times \frac{u}{2} (1+e) = \frac{u}{3} (1+e)$	B1	1	
(c)(i)	$\frac{u}{3}(1+e) = \frac{u}{2}(1-e)$ $2 + 2e = 3 - 3e$ $5e = 1$ $e = \frac{1}{5}$	M1   A1	2	Equating   Solving or showing
(ii)	Speed = $\frac{2u}{5}$	B1	1	
(d)	$B$ reached wall after $\frac{5d}{3u}$  In this time $A$ travels $\frac{5d}{3u} \times \frac{2u}{5} = \frac{2d}{3}$  $A$ and $B$ now have same speed so meet at half remaining distance = $\frac{1}{2} \left( \frac{d}{3} \right)$ $= \frac{d}{6}$  <b>Alternative :</b> Ratio of speeds after collision = 1.5 : 1 Ratio of distance after collision = 1 : $\frac{2}{3}$ Then $d - \frac{2}{3}d$ left = $\frac{d}{3}$ Same speed to meet half way = $\frac{1}{2} \left( \frac{d}{3} \right) = \frac{d}{6}$	M1A1✓  M1A1✓  M1  A1  (M1A1✓) (M1A1✓) (M1) (A1)	6	Attempt to find twice  Attempt to find distance  Attempt to find remaining distances  Special case $\frac{5d}{6} \Rightarrow 5$ marks
	<b>Total</b>		<b>16</b>	
	<b>Total</b>		<b>60</b>	