

GCE 2005

January Series



Mark Scheme

Mathematics A

(MAP4)

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website:
www.aqa.org.uk

Copyright © 2005 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

Dr Michael Cresswell Director General

Key to Mark Scheme

M	mark is for	method
m	mark is dependent on one or more M marks and is for	method
A	mark is dependent on M or m marks and is for	accuracy
B	mark is independent of M or m marks and is for	method and accuracy
E	mark is for	explanation
✓ or ft or F	follow through from previous incorrect result	
CAO	correct answer only	
AWFW	anything which falls within	
AWRT	anything which rounds to	
AG	answer given	
SC	special case	
OE	or equivalent	
A2,1	2 or 1 (or 0) accuracy marks	
-x EE	deduct x marks for each error	
NMS	no method shown	
PI	possibly implied	
SCA	substantially correct approach	
c	candidate	
SF	significant figure(s)	
DP	decimal place(s)	

Abbreviations used in Marking

MC – x	deducted x marks for mis-copy
MR – x	deducted x marks for mis-read
ISW	ignored subsequent working
BOD	given benefit of doubt
WR	work replaced by candidate
FB	formulae booklet

Application of Mark Scheme

No method shown:

Correct answer without working	mark as in scheme
Incorrect answer without working.....	zero marks unless specified otherwise

More than one method/choice of solution:

2 or more complete attempts, neither/none crossed out	mark both/all fully and award the mean mark rounded down
1 complete and 1 partial attempt, neither crossed out	award credit for the complete solution only

Crossed out work

do not mark unless it has not been replaced


Alternative solution using a correct or partially
correct method

award method and accuracy marks as
appropriate

MAP4

Q	Solution	Marks	Total	Comments
1(a)	$\beta = 1 - 2i$	B1	1	
(b)(i)	$\alpha\beta = (1 + 2i)(1 - 2i) = 5$	B1	1	
(ii)	$\alpha\beta\gamma = -30 \quad \gamma = -6$	M1A1F	2	
(c)	Method for either p or q $p = 4, \quad q = -7$	M1 A1FA1F	3	
Total			7	
2(a)	$LHS = r^2 (r^2 + 2r + 1 - (r^2 - 2r + 1))$ $= 4r^3$	M1 A1	2	AG
(b)	$4 \times 50^3 = 50^2 \times 51^2 - 49^2 \times 50^2$ $4 \times 51^3 = 51^2 \times 52^2 - 50^2 \times 51^2$ $4 \times 100^3 = 100^2 \times 101^2 - 99^2 \times 100^2$ $4S = 100^2 \times 101^2 - 49^2 \times 50^2$ $S = 24001875$	M1A1 m1 A1F	4	For $100^2 \times 101^2 - 50^2 \times 51^2$ M1A0m1A0 For $100^2 \times 99^2 - 49^2 \times 50^2$ M1A0m1A0 Clear cancellation shown. If $\sum r^3$ quoted mark M1A1 only
Total			6	
3(a)	$r = \sqrt{2}, \quad \theta = \frac{1}{4}\pi$	B1B1	2	
(b)	$(1+i)^{21} - (1-i)^{21}$ $= (\sqrt{2})^{21} e^{\frac{21\pi i}{4}} - (\sqrt{2})^{21} e^{\frac{-21\pi i}{4}}$ $(\sqrt{2})^{21} \left(\cos \frac{21\pi}{4} + i \sin \frac{21\pi}{4} \right.$ $\quad \left. - \cos \frac{21\pi}{4} + i \sin \frac{21\pi}{4} \right)$ $= (\sqrt{2})^{21} 2i \sin \frac{21\pi}{4}$ $= -2048i$	M1A1 A1F A1F A1F	5	If $\sqrt{2}$ not $(\sqrt{2})^{21}$ lose final A1 also provided of the correct form
Total			7	

MAP4 (cont)

Q	Solution	Marks	Total	Comments	
4(a)(i)	 <p>Circle Centre below x – axis, radius ≈ 5</p>	B1	2		
		B1			
		(ii) Half line with gradient ≈ 1 through $(-3, 0)$	B1		2
		(b)(i) Explanation from diagram	B1		1
		(ii) Verification that $ -7 - 4i + 3 + i = 5$ Verification that $\arg(-7 - 4i + 3) = -\frac{3\pi}{4}$	M1A1 M1A1		4
Total			9		
5(a)(i)	$P\left(\frac{e^x + e^{-x}}{2}\right) + q\left(\frac{e^x - e^{-x}}{2}\right) = r$ $(p + q)e^x + (p - q)e^{-x} = 2r$ $(p + q)e^{2x} - 2re^x + (p - q) = 0$	M1	3	AG	
		A1			
		A1			
		(ii) $e^x = \frac{2r \pm \sqrt{4r^2 - 4(p - q)(p + q)}}{2(p + q)}$ Use of $p^2 = q^2 + r^2$ to show that $e^x = \frac{r}{p + q}$ $e^x > 0 \Rightarrow$ one solution	M1A1 m1A1		5
		(b) $e^x = \frac{12}{18}$ $x = \ln\left(\frac{2}{3}\right)$	M1 A1		2
Total			10		

MAP4 (cont)

Q	Solution	Marks	Total	Comments
6(a)	$f(n+1) - f(n) = 4 \times 7^{n+1} + 3 \times 5^{n+1}$ $+ 5 - 4 \times 7^n - 3 \times 5^n - 5$ Grouping in powers of 7 and 5 $= 4 \times 7^n (7 - 1) + 3 \times 5^n (5 - 1)$ $= 24 \times 7^n + 12 \times 5^n$	M1 m1 A1 A1	4	AG
(b)	$f(1) = M(12)$ shown Assume result true for $n = k$ Then $f(k+1) = f(k) + M(12)$ $= M(12)$ $P(k) \Rightarrow P(k+1)$ and $P(1)$ true	B1 M1 A1 E1	4	Clear demonstration Provided M1 earned
	Total		8	

MAP4 (cont)

Q	Solution	Marks	Total	Comments
7(a)	$\frac{d}{dx}(\sinh^{-1} x) = \frac{1}{\sqrt{1+x^2}}$	B1		
	$\frac{d}{dx}(x\sqrt{1+x^2}) = \sqrt{1+x^2} + \frac{x^2}{\sqrt{1+x^2}}$	M1A1		Reasonable attempt at product rule for M1
	Result = $2\sqrt{1+x^2}$	A1	4	AG
(b)(i)	$S = 2\pi \int_{\ln(\frac{3}{4})}^{\ln(\frac{4}{3})} e^x \sqrt{1+(e^x)^2} dx$	M1		
	$= 2\pi \int_{\ln(\frac{3}{4})}^{\ln(\frac{4}{3})} e^x \sqrt{1+e^{2x}} dx$	A1	2	AG
(ii)	$u = e^x, \frac{du}{dx} = e^x$	M1		Use of formula possibly implied
	$S = 2\pi \int_{\frac{3}{4}}^{\frac{4}{3}} \sqrt{1+u^2} du$	A1		Must be of this form to score further marks ignore limits here
	$= \pi \left[\sinh^{-1} u + u \sqrt{1+u^2} \right]_{\frac{3}{4}}^{\frac{4}{3}}$	A1		
	$= \left[\sinh^{-1} \frac{4}{3} + \frac{4}{3} \sqrt{1+\left(\frac{4}{3}\right)^2} \right]$	A1F		
	$- \left[\sinh^{-1} \frac{3}{4} + \frac{3}{4} \sqrt{1+\left(\frac{3}{4}\right)^2} \right]$			
	$= \pi \left[\ln 3 + \frac{20}{9} - \ln 2 - \frac{15}{16} \right]$	m1A1F		
$= \pi \left[\ln \frac{3}{2} + \frac{185}{144} \right]$	A1F	7		
	Total		13	
	Total		60	