

GCE 2004
June Series



Mark Scheme

Mathematics and Statistics B *MBD2*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from:

Publications Department, Aldon House, 39, Heald Grove, Rusholme, Manchester, M14 4NA
Tel: 0161 953 1170

or

download from the AQA website: www.aqa.org.uk

Copyright © 2004 AQA and its licensors

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

Dr Michael Cresswell Director General

Key to Mark Scheme

M	mark is for	method
m	mark is dependent on one or more M marks and is for	method
A	mark is dependent on M or m marks and is for	accuracy
B	mark is independent of M or m marks and is for	accuracy
E	mark is for	explanation
✓ or ft or F		follow through from previous incorrect result
cao		correct answer only
cso		correct solution only
awfw		anything which falls within
awrt		anything which rounds to
acf		any correct form
ag		answer given
sc		special case
oe		or equivalent
sf		significant figure(s)
dp		decimal place(s)
A2,1		2 or 1 (or 0) accuracy marks
-x ee		deduct x marks for each error
pi		possibly implied
sca		substantially correct approach

Abbreviations used in Marking

MC – x	deducted x marks for mis-copy
MR – x	deducted x marks for mis-read
isw	ignored subsequent working
bod	given benefit of doubt
wr	work replaced by candidate
fb	formulae book

Application of Mark Scheme

No method shown:

Correct answer without working	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise

More than one method / choice of solution:

2 or more complete attempts, neither/none crossed out	mark both/all fully and award the mean mark rounded down
1 complete and 1 partial attempt, neither crossed out	award credit for the complete solution only

Crossed out work	do not mark unless it has not been replaced
------------------	--

Alternative solution using a correct or partially correct method	award method and accuracy marks as appropriate
---	---

Mathematics and Statistics B Discrete 2 MBD2 June 2004

Question Number and Part	Solution	Marks	Total	Comments
1(a)(i)	Nearest neighbour approach gives $A E B C F G H D A$	M1 A1 A1	3	
(ii)	It uses all the 1p links	B1	1	
(b)(i)	Odd vertices $A B C G$ Pairings $AB CG$; $2 + 2$ $AC BG$; $(2 + 1) + 2$ $AG BC$; $(2 + 2) + 1$ So repeat AB and CG	M1 A1 A1 A1	4	
(ii)	Repeat $BC (=1)$, with message starting at A and finishing at G .	B1 B1	2	
Total			10	
2 (a)	Auxiliary equation $m^2 - 5m + 6 = 0$ has roots 2 and 3. General solution $u_n = A.2^n + B.3^n$	M1 A1 A1 A1✓	4	ft
(b)	Try $u_n = k$ to give $k - 5k + 6k = 1$ and $k = \frac{1}{2}$ General solution $u_n = A.2^n + B.3^n + \frac{1}{2}$	M1 A1 A1	3	
Total			7	
3 (a)(i)	Hamming distance = 4	M1 A1	2	
(ii)	Can correct 1 error per word	B1	1	
(b)(i)	e.g. $\begin{bmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 1 & 1 \end{bmatrix}$	B1 M1 A1	3	(or M1 A1 A1 for 2 non-zero words)
(ii)	Matrix $\times (1\ 1\ 0\ 0\ 1\ 1)^T$ and $(0\ 1\ 1\ 0\ 0\ 1)^T$ gives $(0\ 0\ 0)^T$ and $(1\ 0\ 1)^T$ So 1st correct, 2nd has error in 4th place, correcting to 110011011101	M1 A1 M1 A1	4	sc B1 for answer only
Total			10	

MBD2 (cont)

Question Number and Part	Solution	Marks	Total	Comments
4(a)	$P = 3x + 2y + 2z$	B1	1	
(b)(i)	$ \begin{array}{ccccccc} P & x & y & z & s & t & u \\ 1 & 0 & 1 & -\frac{1}{2} & 0 & 1\frac{1}{2} & 0 & 225 \\ 0 & 0 & 0 & \frac{1}{2} & 1 & -\frac{1}{2} & 0 & 5 \\ 0 & \textcircled{1} & 1 & \frac{1}{2} & 0 & \frac{1}{2} & 0 & 75 \\ 0 & 0 & 1 & 2 & 0 & -1 & 1 & 30 \end{array} $	M1 A1 M1 A1 A1	5	Choice of pivot and pivot $\rightarrow 1$ Row deductions
(ii)	Still a negative in top row	B1	1	
4(c)	$ \begin{array}{ccccccc} P & x & y & z & s & t & u \\ 1 & 0 & 1 & 0 & 1 & 1 & 0 & 230 \\ 0 & 0 & 0 & \textcircled{1} & 2 & -1 & 0 & 10 \\ 0 & 1 & 1 & 0 & -1 & 1 & 0 & 70 \\ 0 & 0 & 1 & 0 & -4 & 1 & 1 & 10 \end{array} $	M1 A1 A1 A1	4	
(d)	Maximum of P is 230 at (70,0,10)	B1 \checkmark B1 \checkmark	2	ft near misses
(e)	Slack variable $u \neq 0$. Third inequality has slack; i.e. $2x + 3y + 3z \leq 180$	M1 A1	2	(or test each inequality)
	Total		15	
5 (a)	$2^9 = 512$	M1 A1	2	
(b)(i)	Half the number e.g. By symmetry: each code with an even number of blacks corresponds (by colour change) to one with an odd number of blacks.	B1 B1	2	(or direct count)
(ii)	Can detect one error per bar code	B1 B1	2	
(c)	9:256, 10:512, 11:1024, 12:2048 So increase to 12 strips (or more)	M1 A1	2	
(d)	Reverse of one code can equal a different code. e.g Add an additional black strip on the left and white strip on the right.	B1 M1 A1	3	
	Total		11	

MBD2 (cont)

Question Number and Part	Solution	Marks	Total	Comments
6 (a)	Vertices S and T Arcs SS_1, SS_2, T_1T, T_2T and T_3T Capacities 18, 15, 10, 13, 12 (or more) respectively	M1 A1	 2	
(b) (i)	$8 + 2 + 5 + 4 + 12 = 31$	B1	1	
(ii)	$AB AC DC DE$ (or $AB CB CT_2 CE DE$)	M1 A1	 2	
(c)	e.g. $SS_1ABT_1T: 8$ $SS_1DET_3T: 7$ $SS_2DCET_3T: 5$ $SS_2DACT_2T: 5$ $SS_1ACBT_2T: 2$ $SS_1DCET_2T: 3$	M1 A1 A1 A1 A1 A1	 6	
(d)	All flows \leq all cuts So, by (b)(ii), all flows ≤ 30 . Hence the flow of 30 is maximum possible.	M1 A1	 2	
(e)	e.g. For T_1 to get 10 BT_1 will have a flow of 10. Then, looking at vertex B , max inflow = 10. Hence BT_2 has 0 flow. So maximum arriving at T_2 is from CT_2 and ET_2 with a total capacity of 9.	M1 A1 A1	 3	
	Total		16	

MBD2 (cont)

Question Number and Part	Solution	Marks	Total	Comments
7 (a)(i)	Can take any 1 of the 6 vertical paths	M1 A1	2	(or draw the paths)
(ii)	$n + 1$	B1	1	
(b)(i)	Answer = no. of ways of proceeding from C to B = $n + 1$ from (a)(ii)	B1	1	
(ii)	From D same situation as from A but $n - 1$ wide	B1	1	
(iii)	From A can move to D or C ; R_{n-1} of first type, $n + 1$ of second. R_1 = no of routes with just two vertical squares (so three choices of horizontal route) = 3	M1 A1 B1	3	
(iv)	$R_n = R_{n-1} + (n + 1)$ = $R_{n-2} + n + (n + 1)$... = $R_1 + (3 + 4 + \dots + (n + 1))$ = $3 + (3 + 4 + \dots + (n + 1))$ = $1 + 2 + 3 + \dots + (n + 1)$	M1 A1 A1	3	(or formally solve the recurrence relation)
	Total		11	
	TOTAL		80	