

GCE 2004
June Series



Mark Scheme

Mathematics and Statistics B *MBP5*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available from:

Publications Department, Aldon House, 39, Heald Grove, Rusholme, Manchester, M14 4NA
Tel: 0161 953 1170

or

download from the AQA website: www.aqa.org.uk

Copyright © 2004 AQA and its licensors

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

Dr Michael Cresswell Director General

Key to Mark Scheme

M	mark is for	method
m	mark is dependent on one or more M marks and is for	method
A	mark is dependent on M or m marks and is for	accuracy
B	mark is independent of M or m marks and is for	accuracy
E	mark is for	explanation
✓ or ft or F		follow through from previous incorrect result
cao		correct answer only
cso		correct solution only
awfw		anything which falls within
awrt		anything which rounds to
acf		any correct form
ag		answer given
sc		special case
oe		or equivalent
sf		significant figure(s)
dp		decimal place(s)
A2,1		2 or 1 (or 0) accuracy marks
-x ee		deduct x marks for each error
pi		possibly implied
sca		substantially correct approach

Abbreviations used in Marking

MC – x	deducted x marks for mis-copy
MR – x	deducted x marks for mis-read
isw	ignored subsequent working
bod	given benefit of doubt
wr	work replaced by candidate
fb	formulae book


Application of Mark Scheme

No method shown:	
Correct answer without working	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise
More than one method / choice of solution:	
2 or more complete attempts, neither/none crossed out	mark both/all fully and award the mean mark rounded down
1 complete and 1 partial attempt, neither crossed out	award credit for the complete solution only
Crossed out work	do not mark unless it has not been replaced
Alternative solution using a correct or partially correct method	award method and accuracy marks as appropriate

Mathematics and Statistics B Pure 5 MBP5 June 2004

Question Number and Part	Solution	Marks	Total	Comments
1(a)(i)	$(1+x)^{-1} \approx 1-x \dots$ $+x^2 - x^3 \dots$	B1 B1	2	
(ii)	$(1+4x)^{\frac{1}{2}} = 1 +/ - kx \dots$ $\left(\begin{array}{l} \left(\frac{1}{2}\right)(4x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}(4x)^2 + \\ \left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right) \frac{(4x)^3}{3!} + \dots \end{array} \right)$ $= 1 + 2x - 2x^2 + 4x^3 \dots$	M1 A1 A2,1	4	Valid start to binomial expn. Unsimplified form condone 1 error (A1 if 3 simplified terms correct)
(b)	$\dots = 2(1+4x)^{\frac{1}{2}} + 4(1+x)^{-1} = \dots$ $= 6 + 4x^3$	M1 A1	2	Clear correct use of both expansions cso Must be convinced
Total			8	
2	Mid-ordinates are 1.25, and 1.75 $\int \approx 0.5[f(1.25) + f(1.75)]$ $\approx 0.5\{0.17885098.. + 0.0622753..\}$ $\approx 0.1205 \dots = 0.121$ (to 3 dp)	B1 M1 A1	3	oe must be 0.121
Total			3	
3(a)	$R \cos \alpha = 6$ or $R \sin \alpha = 8$ or $\tan \alpha = \frac{8}{6}$ $R^2 = 6^2 + 8^2$ $\Rightarrow 10 \cos(x + 53.1^\circ)$	M1 M1 A1	3	Accept seen; condone negative signs Alternatively use 2 results in line 1. For α accept awrt 53.1° but R must be 10.
(b)	$\cos(x + \alpha) = 3/R$ $x + \alpha = 360n \pm \dots$ $x + \alpha = 360n \pm 72.5(4\dots)$ $x = 360n \pm 72.5(4\dots) - 53.1(3\dots)$ $[x = 360n + \{19.3 \text{ to } 19.5 \text{ inclusive}\}]$ $x = 360n - \{125 \text{ to } 126 \text{ inclusive}\}]$	M1 m1 A1 A1	4	Possibly implied Accept degrees, rads., mix oe but right hand side must be in degrees Any equivalent forms in degrees sc if m0 then award B1 for either one general solution or 2 particular solutions. covering both branches
Total			7	

MBP5 (cont)

Question Number and Part	Solution	Marks	Total	Comments
4(a)(i) (ii) (b)	$\frac{dy}{dx} = \frac{1}{6}$ Gradient of normal = -6 Eqn of normal $y - 1 = -6(x - 1)$ $y - 1 = -6x + 6 \Rightarrow y + 6x = 7$ $2y \, dy = \frac{1}{x+2} \, dx$ $y^2 = \int \frac{1}{x+2} \, dx$ $y^2 = \ln x+2 + c$ When $x = 1, y = 1 \Rightarrow 1 = \ln 3 + c$ $y^2 = \ln(x+2) + 1 - \ln 3$	B1 M1 m1 A1 M1 A1 A1 m1 A1	1 3 5	ft on (i); used $m \times m' = -1$ ag Clear attempt to separate variables appropriately Condone absence of mod signs and '+c' Valid method for c oe
Total			9	
5(a) (b) (c)	Asymptote $x = -1$ $y = x - 1 + \frac{1}{x+1}$ Asymptote $y = x - 1$ Turning point (0,0) When $y = -4, x^2 + 4x + 4 = 0$ Turning point (-2, -4) 	B1 M1 A1 B1 M1 A1 B1 B1 B1	3 3 3	Full attempt to divide out Alternative Valid method to find $y'(x)$ and then puts $y'(x) = 0$ [M1] $x^2 + 2x = 0 \Rightarrow$ TPs (0,0) [A1] and (-2, -4) [A1] Single upper branch; shape and y not < 0 Single lower branch; shape and y not > -4 Dependent on previous two Bs. Asymptotic behaviour on both branches; through the origin
Total			9	

MBP5 (cont)

Question Number and Part	Solution	Marks	Total	Comments
6(a)	$\cos \theta = \frac{\begin{pmatrix} 1 \\ -1 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix}}{\sqrt{1^2 + (-1)^2 + 2^2} \cdot \sqrt{2^2 + 2^2 + 1^2}}$ $\cos \theta = \frac{2}{\sqrt{6} \cdot \sqrt{9}}$	M1 B1 A1	3	Use of a valid formula For either term in the denominator
(b)(i)	<p>Angle between the normals to the planes</p> $= \cos^{-1} \frac{2}{\sqrt{6} \cdot \sqrt{9}}$ <p>Acute angle between the planes</p> $= 74.2^\circ$	M1 A1✓	2	Only fit on one arithmetical slip and A0 in (i). Accept nearest degree
(ii)	<p>Subst. $(-5, 3, 4)$ into Π_1 gives</p> $-5 - 3 + 8 = 0$ <p>so $(-5, 3, 4)$ lies on Π_1</p> <p>Subst. $(-5, 3, 4)$ into Π_2 gives</p> $-10 + 6 + 4 = 0$ <p>so $(-5, 3, 4)$ lies on Π_2</p>	B1	1	ag
(iii)	$\mathbf{r} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda \overrightarrow{OA}$ $\mathbf{r} = \begin{pmatrix} 0 \\ 0 \\ 0 \end{pmatrix} + \lambda \begin{pmatrix} -5 \\ 3 \\ 4 \end{pmatrix}$	M1 A1	2	Clear understanding that the required line is OA , where A is $(-5, 3, 4)$. oe
	Total		8	

MBP5 (cont)

Question Number and Part	Solution	Marks	Total	Comments
7(a)	At P , $1 + \cos t = 0$ $\Rightarrow t = \pi \Rightarrow x$ -coordinate of P is π^2	M1 A1	2	ag
(b)	$\frac{dx}{dt} = 2t$, $\frac{dy}{dt} = -\sin t$ $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\sin t}{2t}$	M1 A1	2	Attempts both and at least one correct (possibly implied)
(c)(i)	$\frac{d}{dt} \left(\frac{dy}{dx} \right) = \frac{d}{dt} \left(\frac{-\sin t}{2t} \right)$ $= \left[\frac{2t(-\cos t) - (-\sin t)2}{4t^2} \right]$ $= \frac{\sin t - t \cos t}{2t^2}$	M1 A1	2	quotient rule oe used ag cso
(ii)	$\frac{d^2y}{dx^2} = \frac{dt}{dx} \frac{d}{dt} \left(\frac{dy}{dx} \right)$ $= \frac{1}{2t} \left[\frac{\sin t - t \cos t}{2t^2} \right]$	M1 A1	2	using valid formula
(iii)	Necessary condition for pt. of inflection $\frac{d^2y}{dx^2} = 0 \Rightarrow \sin t - t \cos t = 0$ $\Rightarrow \frac{\sin t}{\cos t} = t \Rightarrow \tan t = t$	E1	1	ag
(d)(i)	$\int t \cos t dt = t(\sin t) - \int (\sin t) dt$ $= t \sin t + \cos t + c$	M1 A1 A1✓	3	Condone sign errors only cao ft on previous result Condone absence of $+c$
(ii)	Shaded area $= \int_0^{\pi} y \frac{dx}{dt} dt$ $= \int_0^{\pi} (1 + \cos t) 2t dt$ Shaded area $= \pi^2 - 4$	M1 A1 B1 A1	4	Need attempt to write integrand in terms of t . (ignore limits) Ignore limits For correct limits seen Award for correct solution only
	Total		16	
	TOTAL		60	

