



## General Certificate of Education

# Mathematics and Statistics 6320 *Specification B*

*MBP5 Pure 5*

## Mark Scheme

*2005 examination – June series*

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

## Key to Mark Scheme

<b>M</b>	mark is for	method
<b>m</b>	mark is dependent on one or more M marks and is for	method
<b>A</b>	mark is dependent on M or m marks and is for	accuracy
<b>B</b>	mark is independent of M or m marks and is for	accuracy
<b>E</b>	mark is for	explanation
<b>✓ or ft or F</b>		follow through from previous incorrect result
<b>cao</b>		correct answer only
<b>cso</b>		correct solution only
<b>awfw</b>		anything which falls within
<b>awrt</b>		anything which rounds to
<b>acf</b>		any correct form
<b>ag</b>		answer given
<b>sc</b>		special case
<b>oe</b>		or equivalent
<b>sf</b>		significant figure(s)
<b>dp</b>		decimal place(s)
<b>A2,1</b>		2 or 1 (or 0) accuracy marks
<b>-x ee</b>		deduct x marks for each error
<b>pi</b>		possibly implied
<b>sca</b>		substantially correct approach

## Abbreviations used in Marking

<b>MC – x</b>		deducted x marks for mis-copy
<b>MR – x</b>		deducted x marks for mis-read
<b>isw</b>		ignored subsequent working
<b>bod</b>		given benefit of doubt
<b>wr</b>		work replaced by candidate
<b>fb</b>		formulae book

## Application of Mark Scheme

### **No method shown:**

Correct answer without working	mark as in scheme
Incorrect answer without working	zero marks unless specified otherwise

### **More than one method / choice of solution:**

2 or more complete attempts, neither/none crossed out	mark both/all fully and award the mean mark rounded down
1 complete and 1 partial attempt, neither crossed out	award credit for the complete solution only

### **Crossed out work**

do not mark unless it has not been replaced

### **Alternative solution** using a correct or partially correct method

award method and accuracy marks as appropriate

## Mathematics and Statistics B Pure 5 MBP5 June 2005

Q	Solution	Marks	Total	Comments
1	$h = 0.5$ Integral = $h/2 \{ \dots \}$ $\{ \dots \} = [f(1) + f(2.5) + 2(f(1.5) + f(2))]$ $= \{1 + (4\sqrt{2}-1) + 2[(2\sqrt{2}-1) + 3]\}$ $\{4\sqrt{2}-1=4.65685.. \} \quad \{2\sqrt{2}-1=1.82842.. \}$  Integral to 3sf = 3.83	B1 M1 A1 A1	4	Where $f(x) = 2^x - 1$ . All 4 terms correct. [accept 3 dp or better for each term or 15.31(37...) seen or 3.82(8...) seen if index or surd form not given] cao Must be 3.83
<b>Total</b>			<b>4</b>	
2(a)	$(1+8x)^{\frac{1}{2}} = 1 + kx + \dots$ $\left( \begin{aligned} &\left(\frac{1}{2}\right)(8x) + \frac{\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)}{2!}(8x)^2 + \\ &\left(\frac{1}{2}\right)\left(-\frac{1}{2}\right)\left(-\frac{3}{2}\right)(8x)^3 + \dots \end{aligned} \right)$ $= 1 + 4x$ $-8x^2$ $+32x^3 \dots$	M1 A1 A1 A1	4	Valid start to binomial expn.  Accept $a = 4, b = -8, c = 32$ sc if 0/3 give B1 for at least two unsimplified terms correct in $\{ \}$ above
(b)	Valid for $-\frac{1}{8} < x < \frac{1}{8}$	B1	1	oe Accept $ x  \leq \frac{1}{8}$ oe
<b>Total</b>			<b>5</b>	
3(a)	Asymptotes: $x = 0; y = -4$	B1 B1	2	If no contradiction, accept <b>equations</b> of asymptotes shown on the graph
(b)		B2 B1	3	Correct sketch [B1 if <b>either</b> (i) one correct branch <b>or</b> (ii) correct 2-branch shape translated <b>or</b> (iii) 2-branch curve with intended symmetry about y-axis.  Only pts of intersection with x-axis at -0.5 and 0.5
(c)	$\frac{dy}{dx} = 0 - 2x^{-3} = -2$ at $(1, -3)$  Gradient of normal = $\frac{1}{2}$  Eqn of normal $y + 3 = \frac{1}{2}(x - 1)$	M1 m1 A1	3	Attempts to find $y'$ at $(1, -3)$ having got at least one 'term' correct in $y'(x)$  Uses $m \times m' = -1$ , <u>numerical</u> $m$ 's. PI  Accept in any correct form <u>provided</u> cso
<b>Total</b>			<b>8</b>	

## MBP5 (cont)

Q	Solution	Marks	Total	Comments
4(a)	$R\cos \alpha = 1$ or $R\sin \alpha = 1$ or $\tan \alpha = 1$	M1	3	Accept seen; condone negative signs PI by correct value of $\alpha$ . Altn Use two of results in line1.
	$R^2 = 1^2 + 1^2$ $\Rightarrow R = \sqrt{2}, \alpha = \frac{\pi}{4}$	M1 A1		
(b)	$\sin(x + \alpha) = \frac{1}{R\sqrt{2}} \{=0.5\}$	M1	4	Using (a) to reach $\sin(x + \alpha) = k$ PI  Accept degrees, rads., mix but need both sets of gen. solns....(watch out for valid equivalents)  oe condone degrees or mix but need both sets.
	$x + \alpha = 2\pi n + \sin^{-1} (*)$ , also $x + \alpha = 2\pi n + [\pi - \sin^{-1} (*)]$ , (* = cand's $1/(R\sqrt{2})$ .)	m1		
	$x + \alpha = 2\pi n + \frac{\pi}{6}$ $x + \alpha = 2\pi n + \pi - \frac{\pi}{6}$	A1		
	$x = 2\pi n - \frac{\pi}{12}; 2\pi n + \frac{7\pi}{12}$ [ $x = 2\pi n - \{0.261 \text{ to } 0.262 \text{ inclusive}\}$ $x = 2\pi n + \{1.83 \text{ to } 1.84 \text{ inclusive}\}$ ]	A1		
(c)	$\int \dots = R \int x \sin(x + \alpha) dx$	M1	4	Any equivalent general forms for $x$ in radians. <b>sc</b> if m0 then award B1 for either one general soln. or 2 particular solns. covering both branches condone degrees or mix. <u>In (c) do NOT penalise wrong values for <math>R</math> and <math>\alpha</math>.</u> Use of part (a)
	$\int x \sin(x + \alpha) dx =$			
	$-x \cos(x + \alpha) + \int \cos(x + \alpha) dx$	m1 A1		
	$\int \dots = R \{-x \cos(x + \alpha) + \sin(x + \alpha)\} + c$	A1✓		
	<u>ALT</u> Attempts to integrate <u>both</u> $x \sin x$ and $x \cos x$ by parts	(M1)		
	$\int x \sin x = -x \cos x + \int \cos x$ $\int x \cos x = x \sin x - \int \sin x$ $\dots = -x \cos x + x \sin x + \sin x + \cos x + c$	(m1) (A1) (A1✓)		
<b>Total</b>			<b>11</b>	

## MBP5 (cont)

Q	Solution	Marks	Total	Comments
5(a)	At (0, -1), $\frac{dy}{dx} = \frac{0}{-1} = 0$ (so P is a stationary point)	B1	1	(so.....) not required
(b)(i)	$\frac{d^2y}{dx^2} = \frac{y(1-x)\frac{dy}{dx}}{y^2}$	M1		Clear use of quotient rule [or relevant product rule]
	$\frac{d^2y}{dx^2} = \frac{y(1-x)\left(\frac{x}{y}\right)}{y^2}$	m1		Subst. of $\frac{dy}{dx} = \frac{x}{y}$ oe
	$\frac{d^2y}{dx^2} = \frac{y^2 - x^2}{y^3}$	A1	3	<b>ag</b> cso
(ii)	At (0, -1), $\frac{d^2y}{dx^2} = \frac{1-0}{-1} = -1 < 0$ so P is a maximum point	B1	1	cso
(c)	$y dy = x dx$	M1		Clear attempt to separate variables
	$\frac{y^2}{2} = \frac{x^2}{2} + c$	A1		$\frac{y^2}{2} = \frac{x^2}{2}$
	$\frac{1}{2} = \frac{0}{2} + c$	m1		Use of boundary conditions
	$y^2 = x^2 + 1$	A1	4	
<b>Total</b>			<b>9</b>	
6(a)(i)	$\begin{pmatrix} 1 \\ 2 \\ 4 \end{pmatrix} \cdot \begin{pmatrix} 2 \\ 2 \\ 1 \end{pmatrix} = 2 + 4 + 4 = 10$	B1	1	
(ii)	Magnitude of direction vectors are $\sqrt{21}$ and $\sqrt{9}$ $10 = \sqrt{21} \times \sqrt{9} \cos \theta$ $\cos \theta = \frac{10}{\sqrt{189}} \{=0.72739\dots\}$ $\Rightarrow \theta = 43.3^\circ \{= 43^\circ \text{ to nearest degree}\}$	B1 M1 A1	3	Award for one correct. Use of dot product (ft on earlier values) <b>ag</b> Accept without seeing 3sf if clear
(b)	$(2+s) + (-1+2s) + (-2+4s) = 20$ $\Rightarrow s = 3$ $\Rightarrow Q$ has position vector $5\mathbf{i}+5\mathbf{j}+10\mathbf{k}$	M1 A1 A1✓	3	ft a slip in finding s <b>sc</b> (cand uses $l_2$ ): Mark as max M1A0A1 {8.6i + 6.6j + 4.8k} Condone (5,5,10) notation.
<b>Total</b>			<b>7</b>	

**MBP5 (cont)**

Q	Solution	Marks	Total	Comments
7(a)	When $t = \frac{\pi}{2}$ , $x = 4\sin \frac{\pi}{2} = 4$ and $y = \cos \frac{\pi}{2} = 0$ ie (4,0)	B1	1	<b>ag</b> Accept any valid method
(b)	$\frac{dx}{dt} = 4\cos t$ , $\frac{dy}{dt} = -\sin t$ $\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}} = \frac{-\sin t}{4\cos t} = -\frac{1}{4}\tan t$	M1 A1	2	Attempts both and at least one correct (possibly implied) <b>ag</b> obtained convincingly
(c)(i)	At P, $x = 4\sin \frac{\pi}{4}$ , $y = \cos \frac{\pi}{4}$ grad of tang. = $-\frac{1}{4}\tan \frac{\pi}{4} = -\frac{1}{4}$ Eq tang, $y - \cos \frac{\pi}{4} = -\frac{1}{4}\left(x - 4\sin \frac{\pi}{4}\right)$ $\Rightarrow y - \frac{1}{\sqrt{2}} = -\frac{1}{4}x + \frac{1}{\sqrt{2}}$ $\Rightarrow y = -\frac{1}{4}x + \sqrt{2}$	M1 B1 M1 A1	4	oe Accept $x = 2.82$ to $2.83$ inclusive Accept $y =$ awrt $0.71$ oe <b>ag</b> cso obtained convincingly
(ii)	When $y = 0$ , $x = 4\sqrt{2}$ ; When $x = 0$ , $y = \sqrt{2}$ Area of triangle = 4 Area shaded = area of $\Delta$ – area ‘under curve’ area ‘under curve’ = $\int_0^{\frac{\pi}{2}} y \frac{dx}{dt} dt$ $= \int \cos t (4\cos t) dt$ Area shaded = $4 - \int_0^{\frac{\pi}{2}} 4\cos^2 t dt =$ pr. ans	M1 A1 M1 M1 A1 A1	6	Attempts to find pts where tangent intersects axes Must be justified Need attempt to write integrand in terms of $t$ . Condone wrong/missing limits Ignore limits <b>ag</b> cso be convinced; correct limits should have appeared before the printed answer stage
(iii)	Area shaded = $4 - 2 \int_0^{\frac{\pi}{2}} (1 + \cos 2t) dt$ .... = $4 - 2 \left[ t + \frac{1}{2} \sin 2t \right]_0^{\frac{\pi}{2}}$ ... = $4 - \pi$	M1 A1 A1	3	Use of $2\cos^2 t = 1 + \cos 2t$ (condone sign errors) for [.....] cso
	<b>Total</b>		<b>16</b>	
	<b>TOTAL</b>		<b>60</b>	