

General Certificate of Education
June 2007
Advanced Subsidiary Examination



MATHEMATICS
Unit Mechanics 1A

MM1A/W

Tuesday 5 June 2007 1.30 pm to 2.45 pm

For this paper you must have:

- an 8-page answer book
 - the **blue** AQA booklet of formulae and statistical tables.
- You may use a graphics calculator.

Time allowed: 1 hour 15 minutes

Instructions

- Use blue or black ink or ball-point pen. Pencil should only be used for drawing.
- Write the information required on the front of your answer book. The *Examining Body* for this paper is AQA. The *Paper Reference* is MM1A/W.
- Answer **all** questions.
- Show all necessary working; otherwise marks for method may be lost.
- The **final** answer to questions requiring the use of calculators should be given to three significant figures, unless stated otherwise.
- Take $g = 9.8 \text{ m s}^{-2}$, unless stated otherwise.

Information

- The maximum mark for this paper is 60.
- The marks for questions are shown in brackets.
- Unit Mechanics 1A has a **written paper and coursework**.


Advice

- Unless stated otherwise, you may quote formulae, without proof, from the booklet.

Answer **all** questions.

- 1** A hot air balloon is at rest on the ground. When the balloon is released, it rises to a height of 320 metres in 80 seconds. The balloon moves under the action of its weight and a vertical lift force. Assume that the balloon has a constant acceleration during this motion.
- (a) Show that the acceleration of the balloon is 0.1 m s^{-2} . *(3 marks)*
- (b) Find the speed of the balloon when it reaches a height of 320 metres. *(2 marks)*
- (c) The mass of the balloon is 450 kg. Show that the magnitude of the vertical lift force is 4500 N, correct to two significant figures. *(3 marks)*
- (d) After a while, the vertical lift force is reduced so that the balloon rises at a constant speed. State the magnitude of the vertical lift force when this is the case. *(1 mark)*
- 2** Two particles, *A* and *B*, are moving on a smooth horizontal surface. Particle *A* has mass 2 kg and velocity $\begin{bmatrix} 3 \\ -2 \end{bmatrix} \text{ m s}^{-1}$. Particle *B* has mass 3 kg and velocity $\begin{bmatrix} -4 \\ 1 \end{bmatrix} \text{ m s}^{-1}$. The two particles collide, and they coalesce during the collision.
- (a) Find the velocity of the combined particles after the collision. *(3 marks)*
- (b) Find the speed of the combined particles after the collision. *(2 marks)*


- 3 A sign, of mass 2 kg, is suspended from the ceiling of a supermarket by two light strings. It hangs in equilibrium with each string making an angle of 35° to the vertical, as shown in the diagram. Model the sign as a particle.



- (a) By resolving forces horizontally, show that the tension is the same in each string. *(2 marks)*
- (b) Find the tension in each string. *(5 marks)*
- (c) If the tension in a string exceeds 40 N, the string will break.

Find the mass of the heaviest sign that could be suspended as shown in the diagram. *(3 marks)*


- 4 Two particles, of masses 3.45 kg and 1.45 kg, are connected by a light string that passes over a smooth peg. The particles are released from rest with the strings vertical, as shown in the diagram.



- (a) By forming an equation of motion for each particle, show that the magnitude of the acceleration of each particle is 4 m s^{-2} . *(5 marks)*
- (b) Find the tension in the string. *(2 marks)*
- (c) Initially the particles are at the same level.

Find the speed of the heavier particle when it is 1 metre lower than the lighter particle. Assume that neither particle hits the floor or the peg. *(3 marks)*

- 5 An aeroplane flies in air that is moving due east at a speed of $V \text{ m s}^{-1}$. The velocity of the aeroplane relative to the air is 150 m s^{-1} due north. The aeroplane actually travels on a bearing of 030° .
- (a) Show that $V = 86.6 \text{ m s}^{-1}$, correct to three significant figures. (2 marks)
- (b) Find the magnitude of the resultant velocity of the aeroplane. (3 marks)
- 6 A tennis ball is hit from a height of 2.45 metres above horizontal ground. Initially it travels horizontally at a speed of 20 m s^{-1} , as shown in the diagram.



- (a) Show that the time taken for the tennis ball to reach the ground is 0.707 seconds, correct to three significant figures. (3 marks)
- (b) Find the horizontal distance travelled by the ball when it hits the ground. (2 marks)
- (c) Find the angle between the velocity of the ball and the horizontal when the ball hits the ground. (4 marks)
- 7 A boat is initially at the origin, heading due east at 5 m s^{-1} . It then experiences a constant acceleration of $(-0.2\mathbf{i} + 0.25\mathbf{j}) \text{ m s}^{-2}$. The unit vectors \mathbf{i} and \mathbf{j} are directed east and north respectively.
- (a) State the initial velocity of the boat as a vector. (1 mark)
- (b) Find an expression for the velocity of the boat t seconds after it has started to accelerate. (2 marks)
- (c) Find the value of t when the boat is travelling due north. (3 marks)
- (d) Find the bearing of the boat from the origin when the boat is travelling due north. (6 marks)

END OF QUESTIONS