



General Certificate of Education

Mathematics 6360

MM2B Mechanics 2B

Mark Scheme

2006 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.


Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MM2B

Q	Solution	Marks	Total	Comments
1(a)	$\mathbf{v} = (6t^2 - 2t)\mathbf{i} + (1 - 12t^2)\mathbf{j}$	M1 A1 A1	3	differentiating both components one component correct second component correct
(b)(i)	$\mathbf{v}\left(\frac{1}{3}\right) = \left(\frac{6}{9} - \frac{2}{3}\right)\mathbf{i} + \left(1 - \frac{12}{9}\right)\mathbf{j} = -\frac{1}{3}\mathbf{j}$	M1 A1	2	substituting the value for t into their \mathbf{v} correct velocity
(ii)	Travelling due south	A1ft	1	correct description (Follow through from $\mathbf{v} = \pm k\mathbf{j}$)
(c)	$\mathbf{a} = (12t - 2)\mathbf{i} - 24t\mathbf{j}$ $\mathbf{a}(4) = 46\mathbf{i} - 96\mathbf{j}$	M1 A1 A1	3	differentiating their velocity correct acceleration at time t correct acceleration at $t = 4$
(d)	$\mathbf{F} = 6(46\mathbf{i} - 96\mathbf{j}) = 276\mathbf{i} - 576\mathbf{j}$ $F = \sqrt{276^2 + 576^2} = 639 \text{ N}$ or $a = \sqrt{46^2 + 96^2} = 106.45$ $F = 6 \times 106.45 = 639 \text{ N}$	M1 M1 A1	3	apply Newton's second law correctly finding magnitude correct magnitude
Total			12	
2(a)	$\text{KE} = \frac{1}{2} \times 0.6 \times 14^2 = 58.8 \text{ J}$	M1 A1	2	use of KE formula correct energy
(b)	$0.6 \times 9.8h = 58.8$ $h = \frac{58.8}{0.6 \times 9.8} = 10 \text{ m}$	M1 A1 A1	3	two term energy equation involving PE and previous energy correct equation correct height Note: Constant acceleration methods not accepted.
(c)(i)	WD against resistance $= 58.8 - 0.6 \times 9.8 \times 8$ $= 11.76 = 11.8 \text{ J (to 3 sf)}$	M1 A1 A1	3	three term energy equation correct equation correct value
(ii)	$8F = 11.76$ $F = 1.47 \text{ N}$	M1 A1ft	2	using work done = Fd with $d = 8$ correct force accept 1.48
(d)	The magnitude of the force would <u>vary</u> with the speed of the ball.	B1	1	appropriate explanation
Total			11	

MM2B (cont)

Q	Solution	Marks	Total	Comments
3(a)		B1	1	correct force diagram, with labels and arrows.
(b)	$2T = 0.5 \times 98$ $T = 24.5 \text{ N}$ <p style="text-align: center;">AG</p>	M1 A1 A1	3	moment equation correct equation correct positive value for the tension from correct working
(c)(i)	$2 \times 2 \times 24.5 = 3 \times 9.8 \times m + 0.5 \times 98$ $m = \frac{98 - 49}{3 \times 9.8} = \frac{5}{3} = 1.67 \text{ kg (to 3 sf)}$ <p>Or</p> $2 \times 24.5 = 3 \times 9.8m$ $m = \frac{49}{29.4} = \frac{5}{3} = 1.67 \text{ kg}$	B1 M1 A1 A1 (M1A1) (M1A1)	4	tension doubled moment equation correct equation correct mass for equation for finding m
(ii)	$R = 24.5 \times 2 + 98 + \frac{5}{3} \times 9.8 = 163 \text{ N}$	M1 A1 A1	3	considering vertical equilibrium with 3 terms correct equation correct reaction must be consistent with 3(c)(i) if awarding accuracy marks
(d)	This allows the centre of mass to be placed at the <u>centre of the rod</u> for the moment calculations.	B1	1	correct explanation
Total			12	

MM2B (cont)

Q	Solution	Marks	Total	Comments
4(a)	$\frac{1}{2}mU^2 = \frac{1}{2}mv^2 + mgl(1 - \cos 60^\circ)$ $U^2 = v^2 + gl$ $v = \sqrt{U^2 - gl}$	M1 A1 dM1 A1	4	three/four term energy equation with a trig term correct equation solving for v or v^2 correct v in a simplified form
(b)	$T - mg \cos 60^\circ = m \frac{v^2}{l}$ $T = m \left(\frac{U^2 - gl}{l} + \frac{g}{2} \right) = m \left(\frac{U^2}{l} - \frac{g}{2} \right)$	M1 dM1 A1 dM1 A1	5	resolving towards the centre of the circle with three terms substituting for v^2 correct equation making T the subject correct expression for T . Simplification not necessary.
(c)	$T - mg = m \frac{U^2}{l}$ $T = m \left(\frac{U^2}{l} + g \right)$	M1 A1	2	considering the vertical forces and using Newton's second law with $\frac{U^2}{l}$ correct T
Total			11	
5(a)	$F = 800 + \frac{1200}{20}t = 800 + 60t$ $1200a = 800 + 60t$ $a = \frac{800}{1200} + \frac{60}{1200}t = \frac{2}{3} + \frac{t}{20}$ <p style="text-align: right;">AG</p>	M1 A1 B1 dM1 A1	5	finding the gradient of the line correct gradient correct intercept using Newton's second law on two terms correct result from correct working
(b)	$v = \int \frac{2}{3} + \frac{t}{20} dt = \frac{2t}{3} + \frac{t^2}{40} + c$ $v = 0, t = 0 \Rightarrow c = 0$ $v = \frac{2t}{3} + \frac{t^2}{40}$	M1 A1 A1	3	integrating correct integral with or without c showing $c = 0$
(c)	$s = \int_0^{20} \frac{2t}{3} + \frac{t^2}{40} dt$ $= \left[\frac{t^2}{3} + \frac{t^3}{120} \right]_0^{20}$ $= 200 \text{ m}$	M1 A1 dM1 A1	4	integrating correct integral, with or without c . use of both limits or finding c correct distance
(d)	The $\frac{2t}{3}$ term would change, because only the constant term in the force would change. When integrated this becomes the t term in the velocity.	B1 B1	2	correct term correct explanation
Total			14	

MM2B (cont)

Q	Solution	Marks	Total	Comments
6(a)	$a = \frac{14^2}{50} = 3.92$ $F = 1200 \times 3.92$ AG $= 4704$ N	M1 A1 dM1 A1	4	finding acceleration correct acceleration use of $F = ma$ correct force from correct working
(b)	$R = 1200 \times 9.8 = 11760$ $4704 \leq \mu \times 11760$ $\mu \geq \frac{4704}{11760}$ AG $\mu \geq 0.4$	B1 M1 A1	3	normal reaction applying $F \leq \mu R$ or $F = \mu R$ correct result from correct working
Total			7	
7(a)	$20 \frac{dv}{dt} = -10\sqrt{v}$ $\frac{dv}{dt} = -\frac{\sqrt{v}}{2}$ $\int \frac{1}{\sqrt{v}} dv = \int -\frac{1}{2} dt$ AG $2\sqrt{v} = -\frac{t}{2} + c$ $t = 0, v = 25 \Rightarrow c = 10$ $v = \left(\frac{20-t}{4}\right)^2$	M1 A1 dM1 dM1 A1 dM1 A1	7	applying Newton's second law with $\frac{dv}{dt}$ correct differential equation separating variables integrating correct integrals with or without c finding the constant of integration correct final result from correct working
(b)	$t = 20$	B1	1	correct time
Total			8	
TOTAL			75	