



General Certificate of Education

Mathematics 6360

MS04 Statistics 4

Mark Scheme

2007 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website: www.aqa.org.uk

Copyright © 2007 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
√ or ft or F	follow through from previous incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	or equivalent	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks.

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MS04

Q	Solution	Marks	Total	Comments
1	$H_0 : \sigma = 10 \quad H_1 : \sigma \neq 10$ $\Sigma(x - \bar{x})^2 = 254$ Under $H_0, \sigma^2 = 100$ Hence $\chi^2_{\text{calc}} = \frac{254}{100} = 2.54$ $\nu = 9$ $\chi^2_{\text{crit}} (2.7, 19.0)$ Reject H_0 Evidence that headmaster's belief is incorrect	B1 M1A1 M1A1 B1 B1 A1✓	8	Both Or $s^2 = 28.2$; B1 for 25.4 $\frac{9 \times 28.2}{100} = 2.54$ Both required
Total			8	
2(a)	$E(X) = \frac{1}{p} \quad \text{Var}(X) = \frac{1-p}{p^2}$ $\frac{1}{p} = \frac{4(1-p)}{p^2}$ $\Rightarrow p = 4 - 4p$ $\Rightarrow 5p = 4$ $\Rightarrow p = 0.8$	B1 M1 A1	3	Both
(b)	$P(X > 4 X > 2) = \frac{P(X > 4)}{P(X > 2)}$ $= \frac{0.2^4}{0.2^2}$ $= 0.2^2$ $= 0.04$	M1 m1 A1✓ A1	4	Use of Or $\frac{1 - 0.8(1 + 0.2 + 0.2^2 + 0.2^3)}{1 - 0.8(1 + 0.2)}$ ✓ on (a) CAO
Total			7	

MS04 (cont)

Q	Solution	Marks	Total	Comments		
3(a)	Differences 22, 16, -7, 17, 30, -21, -2, 1, 6, -9	M1		Attempt at differences	2-sample t-test	
	$\bar{d} = \frac{53}{10} = 5.3$	B1				B1
	$H_0 : \mu_d = 0 \quad H_1 : \mu_d > 0$	B1, B1		\bar{d} for μ_d B1		B1, B1
	$t_{\text{calc}} = \frac{5.3 - 0}{\left(\frac{15.85}{\sqrt{10}}\right)}$	B1		s or σ		
	$= 1.06$	M1		Or $\left(\frac{15.03}{\sqrt{9}}\right)$		
		A1				
	$\nu = 9$	B1		PI		$\nu = 18$ B1
	$t_{\text{crit}} = 1.833$	B1				$t_{\text{crit}} = 1.734$ B1
	Retain H_0 - No evidence that mean mark is less on written examination	A1✓	10	OE		5/10 max
	(b) Random sample Differences are normally distributed	E1 E1	2	Differences required		E1 E0 1/2 max
Total			12			
4(a)	$E(X) = \int_0^{\infty} \lambda x e^{-\lambda x} dx$	M1		Use of		
	$= \left[-x e^{-\lambda x} \right]_0^{\infty} + \int_0^{\infty} e^{-\lambda x} dx$	A1		Integrate by parts		
	$= \left[\frac{-e^{-\lambda x}}{\lambda} \right]_0^{\infty}$	A1		Correctly		
	$= \frac{1}{\lambda}$	A1	4	AG		
	(b)(i) $\frac{1}{a} = \frac{1}{62.5} = 0.016$	M1A1	2			
	(ii) $F(t) = 1 - e^{-0.016t}$	M1A1		Or $\int_{80}^{\infty} 0.016 e^{-0.016t} dt$		
	$P(T > 80) = e^{-0.016 \times 80}$	M1		Award M1A1 for complement		
	$= 0.278$	A1	4			
	(iii) Either $e^{-0.016 \times 100} \div e^{-0.016 \times 80}$	M1A1				
	$= 0.726$	A1	3			
or $e^{-0.016 \times 20}$	(M2)					
$= 0.726$	(A1)					
Total			13			

MS04 (cont)

Q	Solution	Marks	Total	Comments																		
5(a)	Mean = $\frac{\sum fx}{\sum f} = \frac{270}{100} = 2.7$	M1A1	2	AG																		
(b)	<table style="margin-left: 20px;"> <tr> <td>O</td> <td>E</td> </tr> <tr> <td>7</td> <td>6.72</td> </tr> <tr> <td>15</td> <td>18.15</td> </tr> <tr> <td>27</td> <td>24.50</td> </tr> <tr> <td>25</td> <td>22.05</td> </tr> <tr> <td>11</td> <td>14.88</td> </tr> <tr> <td>10</td> <td>8.04</td> </tr> <tr> <td>3</td> <td>3.62</td> </tr> <tr> <td>2</td> <td>2.04</td> </tr> </table>	O	E	7	6.72	15	18.15	27	24.50	25	22.05	11	14.88	10	8.04	3	3.62	2	2.04	M1		Probabilities
O	E																					
7	6.72																					
15	18.15																					
27	24.50																					
25	22.05																					
11	14.88																					
10	8.04																					
3	3.62																					
2	2.04																					
		M1		× 100																		
		M1		≥ 7 Frequency = 2.04																		
		M1		Combine classes																		
	$H_0: X \sim \text{Po}$	B1																				
	$\chi^2_{\text{calc}} = \frac{0.28^2}{6.72} + \frac{3.15^2}{18.15} + \frac{2.50^2}{24.50} + \frac{2.95^2}{22.05}$	M1																				
	$+ \frac{3.88^2}{14.88} + \frac{1.96^2}{8.04} + \frac{0.66^2}{5.66}$	A1		≥ 4 terms correct																		
	= 2.77	A1		(0.0117 + 0.5467 + 0.2551 + 0.3947 + 1.012 + 0.4778 + 0.0770)																		
	$\nu = 7 - 2 = 5$	A1		AWFW (2.75, 2.85)																		
	$\chi^2_{\text{crit}} = 9.236$	B1																				
	$\chi^2_{\text{calc}} \ll \chi^2_{\text{crit}}$	B1✓		✓ on $\nu = 6$ only ($\chi^2_{\text{crit}} = 10.645$)																		
	\Rightarrow Accept $X \sim \text{Po}$																					
	ie no evidence that it is not a Poisson distribution	A1✓	11	✓ on χ^2_{calc} and upper $\chi^2_{5 \text{ or } 6}$																		
	Total		13																			

MS04 (cont)

Q	Solution	Marks	Total	Comments
6(a)	$E(aX_1 + bX_2 + cX_3)$ $= aE(X_1) + bE(X_2) + cE(X_3)$ $\Rightarrow \mu = a\mu + b\mu + c\mu$ $\Rightarrow a + b + c = 1$	M1 M1 A1	3	Can be implied by next line AG
(b)(i)	$\text{Var}(T_1) =$ $\frac{1}{9}\text{Var}(X_1) + \frac{1}{4}\text{Var}(X_2) + \frac{1}{36}\text{Var}(X_3)$ $= \frac{7\sigma^2}{18}$ $\text{Var}(T_2) =$ $\frac{4}{9}\text{Var}(X_1) + \frac{9}{16}\text{Var}(X_2) + \frac{25}{144}\text{Var}(X_3)$ $= \frac{85\sigma^2}{72}$ Hence $\text{RE}(T_1 \text{ wrt } T_2) = \frac{\text{Var}(T_2)}{\text{Var}(T_1)}$ $= \frac{85}{72} \times \frac{18}{7} = \frac{85}{28}$	M1 A1 M1 A1 \checkmark	5	Either T_1 or T_2 Accept any correct unreduced fraction or $0.389\sigma^2$ Any equivalent fraction or $1.18\sigma^2$ Use of AWFW [3.03,3.04]
(ii)	Since $\frac{85}{28} > 1$, T_1 is preferred	M1 A1	2	SC $0.39 < 1.18 \Rightarrow T_1$ more efficient B1 \checkmark
	Total		10	

MS04 (cont)

Q	Solution	Marks	Total	Comments
7(a)	$\sigma_x^2 = \frac{761.2}{11} = 69.2$	M1	2	Either
	$\sigma_y^2 = \frac{386.1}{9} = 42.9$	A1		Both correct
(b)(i)	$\frac{69.2}{42.9} = 1.613$	M1A1	8	Both
	$\nu_1 = 12 - 1 = 11 \quad \nu_2 = 10 - 1 = 9$	B1		
	$F_{11,9} = 3.102 \quad F_{9,11} = 2.896$	B1,B1		
	$\frac{1}{3.102} \leq \frac{\left(\frac{\sigma_x^2}{\sigma_y^2}\right)}{1.613} \leq 2.896$	M1 A1✓		
	$\therefore 0.520 \leq \frac{\sigma_x^2}{\sigma_y^2} \leq 4.67$	A1		B1 for limit without full working max 2/5
(ii)	Reject suggestion Since $1 \in \text{CI}$	B1✓ E1✓	2	
	Total		12	
	TOTAL		75	