

General Certificate of Education

Mathematics 6360

MS2A Statistics 2A

Mark Scheme

2006 examination - June series

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Key To Mark Scheme And Abbreviations Used In Marking

M	mark is for method					
m or dM	mark is dependent on one or more M marks and is for method					
A	mark is dependent on M or m marks and is for accuracy					
В	mark is independent of M or m marks an	d is for method	l and accuracy			
Е	mark is for explanation					
$\sqrt{\text{or ft or F}}$	follow through from previous					
	incorrect result	MC	mis-copy			
CAO	correct answer only	MR	mis-read			
CSO	correct solution only	RA	required accuracy			
AWFW	anything which falls within	FW	further work			
AWRT	anything which rounds to	ISW	ignore subsequent work			
ACF	any correct form FIW from incorrect work					
AG	answer given BOD given benefit of doubt					
SC	special case	WR	work replaced by candidate			
OE	or equivalent FB formulae book					
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme			
−x EE	deduct x marks for each error	G	graph			
NMS	no method shown	c	candidate			
PI	possibly implied	sf	significant figure(s)			
SCA	substantially correct approach	dp	decimal place(s)			

No Method Shown

Where the question specifically requires a particular method to be used, we must usually see evidence of use of this method for any marks to be awarded. However, there are situations in some units where part marks would be appropriate, particularly when similar techniques are involved. Your Principal Examiner will alert you to these and details will be provided on the mark scheme.

Where the answer can be reasonably obtained without showing working and it is very unlikely that the correct answer can be obtained by using an incorrect method, we must award **full marks**. However, the obvious penalty to candidates showing no working is that incorrect answers, however close, earn **no marks**.

Where a question asks the candidate to state or write down a result, no method need be shown for full marks

Where the permitted calculator has functions which reasonably allow the solution of the question directly, the correct answer without working earns **full marks**, unless it is given to less than the degree of accuracy accepted in the mark scheme, when it gains **no marks**.

Otherwise we require evidence of a correct method for any marks to be awarded.

MS2A

Q	Solution	Marks	Total	Comments
1(a)	For a 1-year period			
	The number of A grades $\sim P_o(3)$			
	For a 5-year period			
	Number of A grades $\sim P_o(15)$	B1		
	P(Total A-grades > 18)			
	$=1-(Total \le 18)$	M1		
	=1-0.8195	1411		
	= 0.1805			
	= 0.181	A1	3	AWFW 0.180 and 0.181
(b)(i)	$X + Y \sim P_o (10)$	B1		
	$P(X + Y \le 14) = 0.917$	M1A1	3	(0.916 to 0.917)
(ii)	<i>X</i> and <i>Y</i> are independent variables	E1	1	
	Total		7	

MS2A (cont)

O O	,		Marks	Total	Comments	
V	Solution			Wiaiks	Total	Comments
2(a)(i)	22-34 2 35-39 2 40-59 2	A B 21 32 72 36 27 12 20 80	Total 53 108 39 200	B1 B1	2	for A values for B values
(ii)	and age	on between a		B1		At least H ₀
	O_i	\mathbf{E}_{i}	$\frac{\left(O_{i}-E_{i}\right)^{2}}{E_{i}}$	M1 M1		attempt row & column totals attempt at E_i
	21 72 27 32	31.8 64.8 23.4 21.2	3.6679 0.8000 0.5538 5.5019	M1		attempt at $\frac{(O_i - E_i)^2}{E_i}$
	36 12	43.2	1.2000 0.8308	M1		attempt at χ^2
	$\sum O_{r} = 200$ $v = (3-1)(2-1)$	$\sum E_i = 200$	$X^2 = 12.554$	A1 B1		(AWFW 12.5 to 12.6) provided correct method used
	$v = (3-1)(2-1)$ $\chi_{1\%}^{2}(2) = 9.2$			B1 √		on their ν and χ^2
	Reject H ₀					
		ol is situated	t the area within seems to have e of the staff	E1√	9	on χ^2 and calculated value dep on H_0 correct, if stated
(b)	22 - 34 age g		taff employed in sected in school in school B	E1 E1	2	
			Total		13	

Q Solution Marks Total Comments 3(a) $\mu = E(X) = \sum_{\text{all } x} x P(X = x)$ M1 $\mu = (1 \times 0.09) + (2 \times 0.13) + (3 \times 0.50) + (4 \times 0.15) + (5 \times 0.12)$ M1 $\mu = 3.05$ A1 A1 $\sigma^2 = \text{Var}(X) = E(X^2) - \mu^2$ M1 $= 1.2075$ M1 $= 1.2075$ M1 $= 1.2075$ M1 $= P(1.95 < X < 4.15)$ M1 $= P(1.95 < X < 4.15)$ M1 $= 0.78$ A1 3 Total 7 4(a) $\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 1 (b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ B1 Var $(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 Standard deviation $= \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ M1 (or seen on diagram) $= 0.6 \times 1.25$ $= 0.75$ A1 2	MS2A (cont)							
$\mu = (1 \times 0.09) + (2 \times 0.13) + (3 \times 0.50) \\ + (4 \times 0.15) + (5 \times 0.12) $ $\mu = 3.05$ $\sigma^2 = \text{Var}(X) = \text{E}(X^2) - \mu^2$ $= 10.51 - 3.05^2 \\ = 1.2075$ $\therefore \sigma = \sqrt{1.2075} = 1.10$ M1 $= P(1.95 < X < 4.15) \\ = P(2 \le X \le 4) \\ = 0.78$ M1 $= 0.78$ Total $\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 I (b) $\text{E}(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ B1 $\text{Var}(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 $\text{Standard deviation} = \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $\text{P}(R < 3) = \text{P}(-0.3 < R < 0.3) \\ = 0.6 \times 1.25 \\ = 0.75$ A1 2 (or seen on diagram) $= 0.6 \times 1.25 \\ = 0.75$		Solution	Marks	Total	Comments			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$	3(a)	$\mu = E(X) = \sum_{\text{all } x} x P(X = x)$						
$\sigma^{2} = \operatorname{Var}(X) = \operatorname{E}(X^{2}) - \mu^{2}$ $= 10.51 - 3.05^{2}$ $= 1.2075$ $\therefore \sigma = \sqrt{1.2075} = 1.10$ A1 4 (1.099) (b) $P(\mu - \sigma < X < \mu + \sigma)$ $= P(1.95 < X < 4.15)$ $= P(2 \le X \le 4)$ $= 0.78$ A1 3 $\sigma = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 1 (b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ $Var(R) = \frac{1}{12}(0.8)^{2} = \frac{4}{75}$ M1 Standard deviation $= \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ A1 2			M1					
$= 10.51 - 3.05^{2} = 1.2075$ $= 1.2075 = 1.10$ $A1 $			A1					
(b) $P(\mu - \sigma < X < \mu + \sigma)$ = P(1.95 < X < 4.15) $= P(2 \le X \le 4)$ $= 0.78$ M1 $= 0.78$ Total 7 4(a) $\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 1 (b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ $Var(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 Standard deviation $= \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ A1 2 M1 (or seen on diagram)		$=10.51-3.05^2$	M1					
		$\therefore \sigma = \sqrt{1.2075} = 1.10$	A1	4	(1.099)			
$= P(2 \le X \le 4) $ $= 0.78$ Total $A1 = 3$ $A(a) = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ $Var(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 $Standard deviation = \frac{2}{5\sqrt{3}} = 0.231$ A1 $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ A1 2 (or seen on diagram)	(b)							
		,	M1					
4(a) $\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 1 (b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ $Var(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 Standard deviation $= \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ A1 2 M1 (or seen on diagram)		,						
4(a) $\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$ B1 1 (b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ B1 $Var(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$ M1 Standard deviation $= \frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ M1 (or seen on diagram) $= 0.6 \times 1.25$ $= 0.75$ A1 2			Al					
(b) $E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$ $Var(R) = \frac{1}{12}(0.8)^{2} = \frac{4}{75}$ $Standard deviation = \frac{2}{5\sqrt{3}} = 0.231$ $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ B1 M1 A1 3 (or seen on diagram)	4(9)			1				
Var(R) = $\frac{1}{12}$ (0.8) ² = $\frac{4}{75}$ M1 Standard deviation = $\frac{2}{5\sqrt{3}}$ = 0.231 A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ M1 (or seen on diagram) = 0.6 × 1.25	4 (a)	$\alpha = \frac{1}{0.4 - (-0.4)} = 1.25$	B1	1				
Standard deviation = $\frac{2}{5\sqrt{3}} = 0.231$ A1 3 (c) $P(R < 3) = P(-0.3 < R < 0.3)$ $= 0.6 \times 1.25$ $= 0.75$ A1 A1 3 (or seen on diagram)	(b)	$E(R) = \frac{1}{2}(-0.4 + 0.4) = 0$	B1					
(c) $P(R < 3) = P(-0.3 < R < 0.3)$ M1 (or seen on diagram) = 0.6×1.25 A1 2		$\operatorname{Var}(R) = \frac{1}{12}(0.8)^2 = \frac{4}{75}$	M1					
$= 0.6 \times 1.25$ $= 0.75$ A1 2		Standard deviation = $\frac{2}{5\sqrt{3}} = 0.231$	A1	3				
= 0.75 A1 2	(c)		M1		(or seen on diagram)			
			A1					

MS2A (cont)

Q Q	Solution	Marks	Total	Comments
5(a)	$\overline{x} = \frac{471}{5} = 94.2$	B1		Or $s^2 = 36.7$
	5 $s = 6.058$	B1		
	5 – 0.030	D1		
	v = 4 1-tailed test	B1		
	$t_{crit} = -2.132$	B1		Or seen on diagram
	$H_0: \mu = 100$			
	$H_1: \mu < 100$	B1		
	$t = \frac{94.2 - 100}{6.058 \sqrt{5}} = -2.14$	M1A1		$\frac{\text{their } \overline{x} - 100}{\text{their } s / \sqrt{5}}$
	Reject H ₀ at 5% level of significance	A1ft		on their <i>t</i> and critical value
	Evidence at the 5% level of significance to support the members' belief that the batteries last less than 100 hours.	E1√	9	(t values only)
(b)	$\overline{x} = \frac{8080}{80} = 101$ $s^2 = \frac{6399}{79} = 81 \text{or} s^2 = \frac{6399}{80} = 80.0$ $s = 9 s = 8.944$	B1		For s (or s^2) and \overline{x}
	$H_0: \mu = 100$ $H_1: \mu \neq 100$	B1		
	$\overline{X} \sim N\left(100, \frac{81}{80}\right)$ under H_0	B1		or 100, $9/\sqrt{80}$ used
	$z = \frac{101 - 100}{9\sqrt{80}} = 0.99$	M1A1		allow use of t method (AWFW 0.99 to 1.00)
	2-tailed test			
	$z_{crit} = \pm 1.96$	B1		on their z and critical value
	Accept H ₀ at 5% level of significance sufficient evidence at the 5% level of significance to support the manufacturer's	A1√		
	belief	E1√	8	
	Total		17	

MS2A (cont)

Q Q	Solution	Marks	Total	Comments
6(a)	0.5	B2	2	B1 for line segment (0, 0.2) to (1, 0.6) B1 for correct shaped curve (1, 0.6) to (4,0)
	for $0 \le x \le 1$			
	$F(x) = \int_{0}^{x} \frac{1}{5} (2x+1) dx$	M1		
	$= \frac{1}{5} \left(x^2 + x \right) \Big _0^x$ $= \frac{1}{5} x \left(x + 1 \right)$	A1		
	$=\frac{1}{5}x(x+1)$	A1	3	
(ii)	$P(X \ge 0.5) = 1 - F(0.5)$			
	$P(X \ge 0.5) = 1 - F(0.5)$ $= 1 - \frac{1}{5} \times 0.5 \times 1.5$	M1		
	=1-0.15 = 0.85	A1	2	(OE)
(iii)	$F(q_1) = 0.25$	B1		$\frac{1}{5}x(x+1) = 0.25$
	$F(q_1) = 0.25$ $F(0.5) = \frac{3}{20} = 0.15 < 0.25$			↓
	$F(0.75) = \frac{21}{80} = 0.2625 > 0.25$	M1		Attempt to solve
				$q_1 = 0.725$
	$0.5 < q_1 < 0.75$	A1	3	$ \downarrow \\ 0.5 < q_1 < 0.75 \qquad AG $
		otal	10	0.5 \ \q_1 \ \ 0.75 \ AU
	TOT		60	