

GCE 2005

January Series



ASSESSMENT and
QUALIFICATIONS
ALLIANCE

Mark Scheme

Mathematics/Statistics

MS/SS1B

Mark schemes are prepared by the Principal Examiner and considered, together with the relevant questions, by a panel of subject teachers. This mark scheme includes any amendments made at the standardisation meeting attended by all examiners and is the scheme which was used by them in this examination. The standardisation meeting ensures that the mark scheme covers the candidates' responses to questions and that every examiner understands and applies it in the same correct way. As preparation for the standardisation meeting each examiner analyses a number of candidates' scripts: alternative answers not already covered by the mark scheme are discussed at the meeting and legislated for. If, after this meeting, examiners encounter unusual answers which have not been discussed at the meeting they are required to refer these to the Principal Examiner.

It must be stressed that a mark scheme is a working document, in many cases further developed and expanded on the basis of candidates' reactions to a particular paper. Assumptions about future mark schemes on the basis of one year's document should be avoided; whilst the guiding principles of assessment remain constant, details will change, depending on the content of a particular examination paper.

Further copies of this Mark Scheme are available to download from the AQA Website:
www.aqa.org.uk

Copyright © 2005 AQA and its licensors. All rights reserved.

COPYRIGHT

AQA retains the copyright on all its publications. However, registered centres for AQA are permitted to copy material from this booklet for their own internal use, with the following important exception: AQA cannot give permission to centres to photocopy any material that is acknowledged to a third party even for internal use within the centre.

Set and published by the Assessment and Qualifications Alliance.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334. Registered address AQA, Devas Street, Manchester. M15 6EX.

Dr Michael Cresswell Director General

Key to mark scheme and abbreviations used in marking

M	mark is for method		
m or dM	mark is dependent on one or more M marks and is for method		
A	mark is dependent on M or m marks and is for accuracy		
B	mark is independent of M or m marks and is for method and accuracy		
E	mark is for explanation		
✓ or ft or F	follow through from previous		
	incorrect result	MC	mis-copy
CAO	correct answer only	MR	mis-read
CSO	correct solution only	RA	required accuracy
AWFW	anything which falls within	FW	further work
AWRT	anything which rounds to	ISW	ignore subsequent work
ACF	any correct form	FIW	from incorrect work
AG	answer given	BOD	given benefit of doubt
SC	special case	WR	work replaced by candidate
OE	OE	FB	formulae book
A2,1	2 or 1 (or 0) accuracy marks	NOS	not on scheme
-x EE	deduct x marks for each error	G	graph
NMS	no method shown	c	candidate
PI	possibly implied	sf	significant figure(s)
SCA	substantially correct approach	dp	decimal place(s)

MS/SS1B

Q	Solution	Marks	Total	Comments
1(a)	The takings appear to increase slightly as the air temperature increases Weak positive (linear) correlation between air temperature and takings One (or two) unusual results	B1 B1	 2	OE Comments on ranges of values of x and $y \Rightarrow$ B0 OE
(b)	Monday 10	B1	1	CAO; accept point (4, 312)
(c)	$r = 0.817$ to 0.818	B3	3	AWFW for attempts at Σx , $\Sigma x^2 \times 5$ or $S_{xx} \times 3$ M1 for attempted use of correct formula for r M1 for answer A1 If Monday 4 identified in (b), then: $r = 0.0156$ to 0.0157 scores M2 If no Monday removed, then: $r = 0.318$ to 0.319 scores M1
(d)	Temperature at another time Number of other/competing stalls Month/time of year Rainfall/snow Publicity	E1	1	Or a sensible alternative Number of customers \Rightarrow E0 Weather \Rightarrow E0 Population of town \Rightarrow E0
Total			7	


MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
2(a)	Volume $\sim N(\mu, 3.5^2)$			
	Mean, $\bar{x} = \frac{1830}{12} = 152.5$	B1		CAO ($s_{n-1} = 3.778, s_n = 3.617$)
	98% $\Rightarrow z = 2.3263$	B1		AWFW 2.32 to 2.33
	CI for μ is $\bar{x} \pm z \times \frac{(\sigma \text{ or } s)}{\sqrt{n}}$	M1		Use of Must have $(\pm\sqrt{n})$ with $n > 1$
	Thus $152.5 \pm 2.3263 \times \frac{3.5}{\sqrt{12}}$ (150.1 to 150.2, 154.8 to 154.9)	A1✓ A1	5	ft on \bar{x} and z only AWFW
(b)	Evidence, from CI, that mean volume is (above) 150 ml	B1✓		ft on CI in part (a); must be clear comparison of mean of 150 with CI
	In sample, some cans have volumes less than 150 ml	B1		Or reference to range of can volumes in sample
	Thus claim of 150 ml is not justified	B1dep	3	Dependent upon making some comment about mean volume and some comment about individual can volume or range of can volumes
(c)	Volume is normally distributed	E1	1	Accept 'population' or 'X' but not 'it' or ' \bar{X} ' etc ie must be clear statement sample too small \Rightarrow E0
	Total		9	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
3(a)	Scatter Diagram 8, 9 or 10 points plotted	B2	2	5, 6 or 7 points plotted B1
(b)	$b = 7.49$ to 7.51 $a = 14.1$ to 14.6 Regression Line (implied) ≥ 2 points calculated or use of point (\bar{x}, \bar{y}) eg $x = 0 \quad y = 14.3$ & $x = 25 \quad y = 201.9$ straight line drawn	B2 B2 M1 A1	6	AWFW; accept 7.5 AWFW for attempts at $\Sigma x, \Sigma x^2 \times 4$ or $S_{xx} \times 2$ M1 M1 for attempted use of correct formula for b A1 for answers
(c)(i)	$y_{15} = 126$ to 128 Reliable as 15 is within (observed) range	B1 E1		AWFW OE accept points close to line
(ii)	$y_{35} = 276$ to 278 Not reliable as 35 is outside (observed) range	B1 E1	4	AWFW OE accept $y > 4$ hrs so break needed point off graph \Rightarrow E0
(d)	a : time to travel to and from area from/to depot b : (average) time to deliver a /one parcel (within area)	E1 E1	2	OE Both correct but reversed \Rightarrow E1 OE Proportional to packages \Rightarrow E0
Total			14	

Question 3 (a) & (b)



MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
4(a)(i)	$X \sim N(\mu, 4^2)$ $\mu = 106$ $P(X < 110) = P\left(Z < \frac{110-106}{4}\right)$	M1	3	Standardising (109.5, 110 or 110.5) with 106 and ($\sqrt{4}$, 4 or 4^2) and/or $(106 - x)$
	$= P(Z < 1)$ $= 0.841$	A1 A1		
(ii)	$P(\text{underweight}) = P(X < 100)$ $= P(Z < -1.5) = 1 - \Phi(1.5)$ $= 1 - 0.93319 = 0.0668 \text{ to } 0.067$	M1 m1 A1	3	Use of AFWW 99 to 100 Area change AWFW (0.06681)
(b)	2% $\Rightarrow z = -2.0537$	B1	4	AWFW 2.05 to 2.06; ignore sign Standardising AFWW 99 to 100 with μ and 4 Equating z-term to z-value; not using 0.02, 0.98 or $ 1 - z $
	$z = \frac{100 - \mu}{4}$	M1		
	Thus $\frac{100 - \mu}{4} = -2.0537$ Thus $\mu = 108.2 \text{ to } 108.3$	m1 A1		
(c) (i)	$\mu = 108.5$ Mean, $\mu = 108.5$	B1	2	CAO CAO; OE
	Variance, $\frac{\sigma^2}{n} = \frac{4^2}{10} = 1.6$	B1		
(ii)	$P(\bar{X} > 110) = P\left(Z > \frac{110-108.5}{\sqrt{1.6}}\right)$	M1	3	Standardising (109.5, 110 or 110.5) with [μ from (i)] and $\left[\sqrt{\frac{\sigma^2}{10}} \text{ or } \frac{\sigma^2}{10} \text{ from (i)}\right]$ and/or $(\mu - x)$ Area change AWFW (0.11784)
	$= P(Z > 1.19) = 1 - \Phi(1.19)$	m1		
	$= 0.117 \text{ to } 0.119$	A1		
Total			15	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments
5(a)(i)	$p = 0.4$ Attempted use of B(7, 0.4) in (a) $P(X \leq 2) = 0.419$ to 0.421	M1 B1		AWFW (0.4199)
(ii)	$P(X > 1 \text{ and } X < 5) = P(2 \leq X \leq 4)$ $= P(X \leq 4)$ $- P(X \leq 1)$ $= 0.9037 - 0.1586 = 0.744$ to 0.746	M1 M1 A1	5	Identification of at least 2, 3 and 4 Identification of exactly 2, 3 and 4 AWFW (0.7451)
(b)	$P(Y = 7) = \binom{n}{7} (0.4)^7 (0.6)^{n-7}$ $= \binom{28}{7} (0.4)^7 (0.6)^{21}$ $= 0.0425$ to 0.0427	M1 A1 A1	3	Correct expression for B(7; n, 0.4) with $n \neq 7$ Fully correct expression may be implied AWFW (0.042556)
(c)	Mean = $np = 2.8$ SD = $\sqrt{np(1-p)} = \sqrt{1.68}$ $= 1.29$ to 1.31	B1 B1	2	CAO AWFW
(d) (i)	Mean = 2.8 SD = 2.24 to 2.27 $s_{n-1}^2 = 5.14$ to 5.15 and $s_{n-1}^2 = 5.04$	B1 B2	3	CAO $\Sigma fx = 140$ AWFW $\Sigma fx^2 = 644$ Substitution of values into correct formula for variance or SD or SD = 5.03 to 5.15 AFWW M1
(ii)	Means are the same SDs differ greatly Thus answers do not support Aaron's belief	B1✓ B1✓ B1	3	ft on (c) and (d)(i) ft on (c) and (d)(i); but must be s with σ or s^2 with σ^2 Dependent on B1 above CAO
Total			16	

MS/SS1B (cont)

Q	Solution	Marks	Total	Comments																				
6(a)(i)	<table border="1" style="width: 100%; border-collapse: collapse;"> <tr> <td style="width: 10%;"></td> <td style="width: 15%; text-align: center;"><i>M</i></td> <td style="width: 15%; text-align: center;"><i>A</i></td> <td style="width: 15%; text-align: center;"><i>S</i></td> <td style="width: 15%; text-align: center;"><i>T</i></td> </tr> <tr> <td><i>M</i></td> <td style="text-align: center;">38</td> <td style="text-align: center;">369</td> <td style="text-align: center;">303</td> <td style="text-align: center;">710</td> </tr> <tr> <td><i>F</i></td> <td style="text-align: center;">26</td> <td style="text-align: center;">275</td> <td style="text-align: center;">643</td> <td style="text-align: center;">944</td> </tr> <tr> <td><i>T</i></td> <td style="text-align: center;">64</td> <td style="text-align: center;">644</td> <td style="text-align: center;">946</td> <td style="text-align: center;">1654</td> </tr> </table>		<i>M</i>	<i>A</i>	<i>S</i>	<i>T</i>	<i>M</i>	38	369	303	710	<i>F</i>	26	275	643	944	<i>T</i>	64	644	946	1654			
		<i>M</i>	<i>A</i>	<i>S</i>	<i>T</i>																			
	<i>M</i>	38	369	303	710																			
	<i>F</i>	26	275	643	944																			
<i>T</i>	64	644	946	1654																				
	$P(F) = 944/1654 (= 0.571)$	M1	1	Use of																				
(ii)	$P(F \cap A) = 275/1654 (= 0.166)$	M1	1	Use of																				
(iii)	$P(F \cup A) = \frac{944 + 369}{1654}$ $= 1313/1654 \text{ or } 0.793 \text{ to } 0.795$	M1 A1	2	Use of; OE CAO/AFWW (0.7938)																				
(iv)	$P(F A) = \frac{\text{their (ii)}}{644/1654}$ $= 275/644 \text{ or } 0.426 \text{ to } 0.428$	M1 A1	2	Use of CAO/AFWW (0.4270)																				
(b)(i)	$P(MMM) = \frac{710 \times 709 \times 708}{1654 \times 1653 \times 1652}$ $= 0.0788 \text{ to } 0.0791$	M1 A1	2	Use of (without replacement) AWFW (no fraction) (0.07891)																				
(ii)	$P(MFF) = \frac{710 \times 944 \times 943 \times 3}{1654 \times 1653 \times 1652}$ $= 0.419 \text{ to } 0.421$	M1 M1 A1	3	Use of one combination of <i>MFF</i> (without replacement) Use of multiplier of 3 AWFW (no fraction) (0.4198)																				
(c)(i)	Female (and) Academic	B1	1	CAO																				
(ii)	Male OR Academic (or both)	B1 B1	2	Not female \Rightarrow B0 'OR' must be clearly stated or implied Addition of 'not both' \Rightarrow B0																				
	Total		14																					
	TOTAL		75																					