

ALLIANCE

# Mark scheme June 2003

## GCE

### Physics A

Unit PA01

Copyright © 2003 AQA and its licensors. All rights reserved.

The Assessment and Qualifications Alliance (AQA) is a company limited by guarantee registered in England and Wales 3644723 and a registered charity number 1073334 Registered address: Addleshaw Booth & Co., Sovereign House, PO Box 8, Sovereign Street, Leeds LS1 1HQ Kathleen Tattersall: *Director General* 

www.theallpapers.com

### Unit 1

### 1 (a) number of protons = number of electrons (e.g.14) $\checkmark$ number of protons + number of neutrons = 28 $\checkmark$ (b)(i) nuclei with the same number of protons $\checkmark$ but different number of neutrons/nucleons $\checkmark$ (b)(ii) $(137-55) = 82 \checkmark$ (b)(iii) $\frac{Q}{m} = \frac{92 \times 1.60 \times 10^{-19}}{236 \times 1.67 \times 10^{-27}} \checkmark$ $= 3.73 \times 10^7 (C \text{ kg}^{-1}) \checkmark$ (b)(iv) X (= 236 - 137 - 4) = 95 $\checkmark$

2 (a)(i) positron, neutron, neutrino, positive pion ✓✓(if all correct) (lose ✓ for each error)

(a)(ii) electron, proton, negative muon  $\checkmark \checkmark$  (if all correct) (lose  $\checkmark$  for each error) (4)

(b)(i) 
$$(\mu^{-}) \rightarrow e^{-} + \overline{v_{e}} + v_{\mu} \checkmark$$

(b)(ii) difference: mass or half-life or generation of lepton ✓ similarity: both leptons or both negatively charged ✓ (3)

(c)



 $\frac{(3)}{(10)}$ 

(2)

(<u>6</u>) (8)

3

(a) there must be a large distance between collisions to allow

electrons to gain enough energy  $\checkmark$ 

[or the vapour must not completely absorb the electrons] (1)

(b)	the mercury vapour emits ultra violet (radiation) $\checkmark$ the coating absorbs electromagnetic radiation/light from the mercury $\checkmark$ emits longer wavelengths/lower frequencies $\checkmark$ in the visible region $\checkmark$	<sub>max</sub> (3)
		<u>(4)</u>
<b>4</b> (a)	the minimum frequency (of radiation) $\checkmark$ required to eject photoelectrons $\checkmark$	(2)
(b)(i)	(use of $\phi = hf_0$ gives) $\phi = 6.63 \times 10^{-34} \times 4.85 \times 10^{14} \checkmark$ = 3.22 × 10 <sup>-19</sup> (J) $\checkmark$	
(b)(ii)	$\phi \left( = \frac{3.22 \times 10^{-19}}{1.60 \times 10^{-19}} \right) = 2.01 \text{ (eV) } \checkmark$	
	(allow C.E. for value of $\phi$ from (i))	(3)
(c)	line parallel to the given line $\checkmark$ with half the value of the <i>x</i> - intercept $\checkmark$	(2)
(d)	statement : increase the light intensity/brightness $\checkmark$	
	explanation : more incident photons (per second) one photon interacts with one electron more emitted electrons (per second) greater rate of <u>flow charge</u> (any two) $\checkmark \checkmark$	<u>(3)</u> (10)
5	$c_{1} = (300 \times 10^{8})$	
(a)(i)	(use of $n = \frac{1}{2}$ gives) $c_{\text{obsc}} = \frac{5.00 \times 10^{8}}{100} = 2.07 \times 10^{8} \text{ m s}^{-1} \checkmark$	

(a)(i) (use of 
$$n = \frac{c_1}{c_2}$$
 gives)  $c_{\text{glass}} \left( = \frac{3.00 \times 10^8}{1.45} \right) = 2.07 \times 10^8 \text{ m s}^{-1} \checkmark$ 

(a)(ii) use of 
$$\frac{\sin\theta_1}{\sin\theta_2} = \frac{c_1}{c_2} \checkmark$$
  
 $c_{\text{liquid}} = \frac{2.07 \times 10^8 \times \sin 29.2^\circ}{\sin 26.6^\circ} = 2.26 \times 10^8 \text{ m s}^{-1} \checkmark$  (3)  
(allow C.E. for values of  $c_{\text{glass}}$  from (i))

(b) use of 
$$_{1}n_{2} = \frac{c_{1}}{c_{2}}$$
 and  $_{1}n_{2} = \frac{n_{2}}{n_{1}} \checkmark$   
to give  $n_{\text{liquid}} = \frac{1.45 \times 2.07 \times 10^{8}}{2.26 \times 10^{8}} = 1.33 \checkmark$ 

$$\left[ \text{or } n_l = \frac{c_1}{c_{\text{liquid}}} = \frac{3 \times 10^8}{2.26 \times 10^8} = 1.33 \right] \text{ (allow C.E. for value of } c_{\text{liquid}} \text{ )}$$

[or use 
$$_1n_2 = \frac{\sin\theta_1}{\sin\theta_2}$$
 and  $_1n_2 = \frac{n_2}{n_1}$  to give correct answer] (2)

(c) diagram to show : total internal reflection on the vertical surface  $\checkmark$ refraction at bottom surface with angle in air greater than that in the liquid (29.2°)  $\checkmark$  (2)

<u>(7)</u>

6  
(a)(i) an electron moves up from one energy level to another 
$$\checkmark$$
  
(a)(ii) an electron is removed from an atom  $\checkmark$   
(b) (use of  $hf = E_2 - E_1$  gives)  $f = (2.56 - 1.92) \times 10^{-19} \checkmark / 6.63 \times 10^{-34}$   
 $= 9.65 \times 10^{13}$  Hz  $\checkmark$   
(allow C.E. for incorrect  $\Delta E$ )  
(2)  
(2)  
(2)  
(2)  
(4)

7

6

(a)(i) electrons behave as both particles and waves  $\checkmark$ 

(a)(ii) particle: deflection in an electromagnetic field  
or other suitable examples 
$$\checkmark$$
  
wave: electron diffraction  $\checkmark$  (3)

(b) (use of 
$$\lambda = \frac{h}{mv}$$
 gives)  $v \left( = \frac{h}{m\lambda} \right) = \frac{6.63 \times 10^{-34}}{9.11 \times 10^{-31} \times 1.7 \times 10^{-10}} \checkmark$   
= 4.28 × 10<sup>6</sup> m s<sup>-1</sup> ✓ (2)

Quality of Written Communication (Q3(b) and Q4(d) 
$$\checkmark \checkmark$$
(2)(2)(2)