

General Certificate of Education
January 2003
Advanced Level Examination



PHYSICS (SPECIFICATION A)
Unit 4 Waves, Fields and Nuclear Energy

PA04

Section A

Monday 27 January 2003 Morning Session

In addition to this paper you will require:

- an objective test answer sheet;
- a black or blue ball-point pen;
- a calculator;
- a question paper/answer book for Section B (enclosed).

Time allowed: The total time for Section A and Section B of this paper is 1 hour 30 minutes

Instructions

- Use a blue or black ball-point pen. Do **not** use pencil.
- Answer **all** questions in this section.
- For each question there are four responses. When you have selected the response which you think is the most appropriate answer to a question, mark this response on your answer sheet.
- Mark all responses as instructed on your answer sheet. If you wish to change your answer to a question, follow the instructions on your answer sheet.
- Do all rough work in this book **not** on the answer sheet.

Information

- The maximum mark for this section is 30.
- Section A and Section B of this paper together carry 15% of the total marks for Physics Advanced.
- All questions in Section A carry equal marks. No deductions will be made for incorrect answers.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- The question paper/answer book for Section B is enclosed within this question paper.

Data Sheet

- A perforated Data Sheet is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

Fundamental constants and values

Quantity	Symbol	Value	Units
speed of light in vacuo	c	3.00×10^8	m s^{-1}
permeability of free space	μ_0	$4\pi \times 10^{-7}$	H m^{-1}
permittivity of free space	ϵ_0	8.85×10^{-12}	F m^{-1}
charge of electron	e	1.60×10^{-19}	C
the Planck constant	h	6.63×10^{-34}	J s
gravitational constant	G	6.67×10^{-11}	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	N_A	6.02×10^{23}	mol^{-1}
molar gas constant	R	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	k	1.38×10^{-23}	J K^{-1}
the Stefan constant	σ	5.67×10^{-8}	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	α	2.90×10^{-3}	m K
electron rest mass	m_e	9.11×10^{-31}	kg
(equivalent to $5.5 \times 10^{-4}u$)			
electron charge/mass ratio	e/m_e	1.76×10^{11}	C kg^{-1}
proton rest mass	m_p	1.67×10^{-27}	kg
(equivalent to 1.00728u)			
proton charge/mass ratio	e/m_p	9.58×10^7	C kg^{-1}
neutron rest mass	m_n	1.67×10^{-27}	kg
(equivalent to 1.00867u)			
gravitational field strength	g	9.81	N kg^{-1}
acceleration due to gravity	g	9.81	m s^{-2}
atomic mass unit	u	1.661×10^{-27}	kg
(1u is equivalent to 931.3 MeV)			

Fundamental particles

Class	Name	Symbol	Rest energy /MeV
photon	photon	γ	0
lepton	neutrino	ν_e	0
		ν_μ	0
mesons	electron	e^\pm	0.510999
	muon	μ^\pm	105.659
	pion	π^\pm	139.576
		π^0	134.972
kaon	K^\pm	493.821	
	K^0	497.762	
	proton	p	938.257
baryons	neutron	n	939.551

Properties of quarks

Type	Charge	Baryon number	Strangeness
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1

Geometrical equations

- arc length = $r\theta$
- circumference of circle = $2\pi r$
- area of circle = πr^2
- area of cylinder = $2\pi rh$
- volume of cylinder = $\pi r^2 h$
- area of sphere = $4\pi r^2$
- volume of sphere = $\frac{4}{3}\pi r^3$

Mechanics and Applied Physics

- $v = u + at$
- $s = \left(\frac{u+v}{2}\right)t$
- $s = ut + \frac{at^2}{2}$
- $v^2 = u^2 + 2as$
- $F = \frac{\Delta(mv)}{\Delta t}$
- $P = Fv$
- efficiency = $\frac{\text{power output}}{\text{power input}}$
- $\omega = \frac{v}{r} = 2\pi f$
- $a = \frac{v^2}{r} = r\omega^2$
- $I = \sum mr^2$
- $E_k = \frac{1}{2} I\omega^2$
- $\omega_2 = \omega_1 + at$
- $\theta = \omega_1 t + \frac{1}{2} at^2$
- $\omega_2^2 = \omega_1^2 + 2a\theta$
- $\theta = \frac{1}{2} (\omega_1 + \omega_2)t$
- $T = I\alpha$
- angular momentum = $I\omega$
- $W = T\theta$
- $P = T\omega$
- angular impulse = change of angular momentum = Tt
- $\Delta Q = \Delta U + \Delta W$
- $\Delta W = p\Delta V$
- $pV^\gamma = \text{constant}$
- work done per cycle = area of loop
- input power = calorific value \times fuel flow rate
- indicated power as (area of $p-V$ loop) \times (no. of cycles/s) \times (no. of cylinders)
- friction power = indicated power - brake power
- efficiency = $\frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$
- maximum possible efficiency = $\frac{T_H - T_C}{T_H}$

Fields, Waves, Quantum Phenomena

- $g = \frac{F}{m}$
- $g = -\frac{GM}{r^2}$
- $g = -\frac{\Delta V}{\Delta x}$
- $V = -\frac{GM}{r}$
- $a = -(2\pi f)^2 x$
- $v = \pm 2\pi f \sqrt{A^2 - x^2}$
- $x = A \cos 2\pi ft$
- $T = 2\pi \sqrt{\frac{m}{k}}$
- $T = 2\pi \sqrt{\frac{l}{g}}$
- $\lambda = \frac{\omega s}{D}$
- $d \sin \theta = n\lambda$
- $\theta \approx \frac{\lambda}{D}$
- $n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$
- $n_2 = \frac{n_2}{n_1}$
- $\sin \theta_c = \frac{1}{n}$
- $E = hf$
- $hf = \phi + E_k$
- $hf = E_1 - E_2$
- $\lambda = \frac{h}{p} = \frac{h}{mv}$
- $c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$
- Electricity**
- $\epsilon = \frac{E}{Q}$
- $\epsilon = I(R+r)$
- $\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$
- $R_T = R_1 + R_2 + R_3 + \dots$
- $P = I^2 R$
- $E = \frac{F}{Q} = \frac{V}{d}$
- $E = \frac{1}{4\pi \epsilon_0} \frac{Q}{r^2}$
- $E = \frac{1}{2} QV$
- $F = BI$
- $F = BQv$
- $Q = Q_0 e^{-t/RC}$
- $\Phi = BA$

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	2.00×10^{30}	7.00×10^8
Earth	6.00×10^{24}	6.40×10^6

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

Electronics

Resistors

Preferred values for resistors (E24)
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2 2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2 6.8 7.5 8.2 9.1 ohms
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

Alternating Currents

$$f = \frac{1}{T}$$

Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left(\frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

SECTION A


In this section each item consists of a question followed by four suggested answers.

You are to select the most appropriate answer in each case.

You are advised to spend approximately 30 minutes on this section.

- 1 Which one of the following gives the phase difference between the particle velocity and the particle displacement in simple harmonic motion?
- A $\frac{\pi}{4}$ rad
- B $\frac{\pi}{2}$ rad
- C $\frac{3\pi}{4}$ rad
- D 2π rad
- 2 A particle oscillates with undamped simple harmonic motion. Which one of the following statements about the acceleration of the oscillating particle is true?
- A It is least when the speed is greatest.
- B It is always in the opposite direction to its velocity.
- C It is proportional to the frequency.
- D It decreases as the potential energy increases.

3




The graph shows, at a particular instant, the variation of the displacement of the particles in a transverse progressive water wave, of wavelength 4 cm, travelling from left to right. Which one of the following statements is **not** true?

- A The distance PS = 3 cm.
- B The particle velocity at Q is a maximum.
- C The particle at S is moving downwards.
- D Particles at P and R are in phase.

-
- 4 The audible range of a girl's hearing is 30 Hz to 16 500 Hz. If the speed of sound in air is 330 m s^{-1} , what is the shortest wavelength of sound in air which the girl can hear?
- A $\frac{30}{330} \text{ m}$
- B $\frac{16\,500}{330} \text{ m}$
- C $\frac{330}{16\,500} \text{ m}$
- D $\frac{330}{30} \text{ m}$
- 5 Which one of the following types of wave **cannot** be polarised?
- A radio
- B ultraviolet
- C microwave
- D ultrasonic
- 6 A uniform wire fixed at both ends is vibrating in its fundamental mode. Which one of the following statements is **not** correct for all the vibrating particles?
- A They vibrate in phase.
- B They vibrate with the same amplitude.
- C They vibrate with the same frequency.
- D They vibrate at right angles to the wire.
- 7 A $1 \mu\text{F}$ capacitor is charged using a constant current of $10 \mu\text{A}$ for 20 s. What is the energy finally stored by the capacitor?
- A $2 \times 10^{-3} \text{ J}$
- B $2 \times 10^{-2} \text{ J}$
- C $4 \times 10^{-2} \text{ J}$
- D $4 \times 10^{-1} \text{ J}$


8



A capacitor of capacitance $10\ \mu\text{F}$ is fully charged through a resistor R to a p.d. of $20\ \text{V}$ using the circuit shown. Which one of the following statements is **incorrect**?

- A The p.d. across the capacitor is $20\ \text{V}$.
- B The p.d. across the resistor is $0\ \text{V}$.
- C The energy stored by the capacitor is $2\ \text{mJ}$.
- D The total energy taken from the battery during the charging process is $2\ \text{mJ}$.

9



A model car moves in a circular path of radius $0.8\ \text{m}$ at an angular speed of $\frac{\pi}{2}\ \text{rad s}^{-1}$. What is its displacement from point P , $6\ \text{s}$ after passing P ?

- A zero
- B $1.6\ \text{m}$
- C $0.4\pi\ \text{m}$
- D $1.6\pi\ \text{m}$

- 10 A small mass is situated at a point on a line joining two large masses m_1 and m_2 such that it experiences no resultant gravitational force. If its distance from the mass m_1 is r_1 and its distance from the mass m_2 is r_2 , what is the value of the ratio $\frac{r_1}{r_2}$?

A $\frac{m_1^2}{m_2^2}$

B $\frac{m_2^2}{m_1^2}$

C $\sqrt{\frac{m_1}{m_2}}$

D $\sqrt{\frac{m_2}{m_1}}$

- 11 A planet of mass M and radius R rotates so rapidly that loose material at the equator just remains on the surface. What is the period of rotation of the planet?

G is the universal gravitational constant.

A $2\pi\sqrt{\frac{R}{GM}}$

B $2\pi\sqrt{\frac{R^2}{GM}}$

C $2\pi\sqrt{\frac{GM}{R^3}}$

D $2\pi\sqrt{\frac{R^3}{GM}}$

- 12 Which one of the following has different units to the other three?


- A gravitational potential
- B gravitational field strength
- C force per unit mass
- D gravitational potential gradient

- 13 Two horizontal parallel plate conductors are separated by a distance of 5.0 mm in air. The lower plate is earthed and the potential of the upper plate is + 50 V.

Which line, **A** to **D**, gives correctly the electric field strength, E , and the potential, V , at a point midway between the plates?

	<i>electric field strength $E/\text{V m}^{-1}$</i>	<i>potential V/V</i>
A	1×10^4 upwards	25
B	1×10^4 downwards	25
C	1×10^4 upwards	50
D	1×10^4 downwards	50

- 14 The diagram shows a vertical square coil whose plane is at right angles to a horizontal uniform magnetic field B . A current, I , flows in the coil, which can rotate about a vertical axis OO' .



Which one of the following statements is correct?

- A** The forces on the two vertical sides of the coil are equal and opposite.
B A couple acts on the coil.
C No forces act on the horizontal sides of the coil.
D If the coil is turned through a small angle about OO' , it will remain in position.
- 15 An α particle and a β^- particle both enter the same uniform magnetic field, which is perpendicular to their direction of motion. If the β^- particle has a speed 15 times that of the α particle, what is the value of the ratio

$$\frac{\text{magnitude of force on } \beta^- \text{ particle}}{\text{magnitude of force on } \alpha \text{ particle}} ?$$

- A** 3.7
B 7.5
C 60
D 112.5

END OF QUESTIONS

Surname		Other Names	
Centre Number		Candidate Number	
Candidate Signature			

General Certificate of Education
January 2003
Advanced Level Examination



PHYSICS (SPECIFICATION A)
Unit 4 Waves, Fields and Nuclear Energy

PA04

Section B

Monday 27 January 2003 Morning Session

In addition to this paper you will require:

- a calculator;
- a pencil and a ruler.

Time allowed: The total time for Section A and Section B of this paper is 1 hour 30 minutes

Instructions

- Use a blue or black ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions in the spaces provided. All working must be shown.
- Do all rough work in this book. Cross through any work you do not want marked.

Information


- The maximum mark for this Section is 30.
- Mark allocations are shown in brackets.
- Section A and Section B of this paper together carry 15% of the total marks for Physics Advanced.
- You are expected to use a calculator where appropriate.
- In questions requiring description and explanation you will be assessed on your ability to use an appropriate form and style of writing, to organise relevant information clearly and coherently, and to use specialist vocabulary where appropriate. The degree of legibility of your handwriting and the level of accuracy of your spelling, punctuation and grammar will also be taken into account.
- A *Data Sheet* is provided on pages 3 and 4 of Section A. You may wish to detach this perforated sheet at the start of the examination.

For Examiner's Use			
Number	Mark	Number	Mark
1			
2			
3			
4			
Total (Column 1)	→		
Total (Column 2)	→		
TOTAL			
Examiner's Initials			

Answer **all** questions.

You are advised to spend approximately **one** hour on this Section.

- 1 A simple pendulum consists of a 25 g mass tied to the end of a light string 800 mm long. The mass is drawn to one side until it is 20 mm above its rest position, as shown in the diagram. When released it swings with simple harmonic motion.



- (a) Calculate the period of the pendulum.

.....

.....

(2 marks)

- (b) Show that the initial amplitude of the oscillations is approximately 0.18 m, and that the maximum speed of the mass during the first oscillation is about 0.63 m s^{-1} .

.....

.....

.....

.....

.....

.....

.....

(4 marks)

- (c) Calculate the magnitude of the tension in the string when the mass passes through the lowest point of the first swing.

.....

.....

.....

.....

.....

(2 marks)



TURN OVER FOR THE NEXT QUESTION

2 A vertical screen is placed several metres beyond a vertical double slit arrangement illuminated by a laser. The diagram below shows a full-size tracing of the pattern of spots obtained on this screen. The black patches represent red light whilst the spaces between them are dark.



(a) Using the wave theory, explain how the pattern of bright and dark patches is formed. You may be awarded marks for the quality of written communication provided in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(3 marks)

(b) The slit separation was 0.90 mm and the distance between the slits and the screen was 4.2 m.

(i) Calculate the spacing of the bright fringes by taking measurements on the diagram of the tracing.

.....
.....

(ii) Hence determine the wavelength of the laser light used.

.....
.....
.....
.....


(4 marks)

$\frac{7}{7}$

TURN OVER FOR THE NEXT QUESTION

- 3 The mass of the nucleus of an isolated copper atom is 63 u and it carries a charge of $+29e$. The diameter of the atom is $2.3 \times 10^{-10} \text{ m}$.

P is a point at the outer edge of the atom.



(a) Calculate

- (i) the electric field strength at P due to the nucleus,

.....

.....

.....

- (ii) the gravitational potential at P due to the nucleus.

.....

.....

.....

(5 marks)

(b) Draw an arrow on the above diagram to show the direction of the electric field at the point P.

(1 mark)

6

