

General Certificate of Education  
June 2006  
Advanced Level Examination



**PHYSICS (SPECIFICATION A)**  
**Unit 4 Waves, Fields and Nuclear Energy**

**PA04**

**Section A**

Thursday 15 June 2006 9.00 am to 10.30 am

**For this paper you must have:**

- an objective test answer sheet
- a black ball-point pen
- a calculator
- a question paper/answer book for Section B (enclosed)

Time allowed: The total time for Section A and Section B of this paper is 1 hour 30 minutes

**Instructions**

- Use a black ball-point pen. Do **not** use pencil.
- Answer **all** questions in this section.
- For each question there are four responses. When you have selected the response which you think is the most appropriate answer to a question, mark this response on your answer sheet.
- Mark all responses as instructed on your answer sheet. If you wish to change your answer to a question, follow the instructions on your answer sheet.
- Do all rough work in this book **not** on the answer sheet.

**Information**

- The maximum mark for this section is 30.
- All questions in Section A carry equal marks. No deductions will be made for incorrect answers.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- The question paper/answer book for Section B is enclosed within this question paper.

**Data Sheet**

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

**Fundamental constants and values**

Quantity	Symbol	Value	Units
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$
charge of electron	$e$	$1.60 \times 10^{-19}$	C
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$
the Wien constant	$a$	$2.90 \times 10^{-3}$	$\text{m K}$
electron rest mass	$m_e$	$9.11 \times 10^{-31}$	kg
(equivalent to $5.5 \times 10^{-4} \text{u}$ )			
electron charge/mass ratio	$e/m_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$
proton rest mass	$m_p$	$1.67 \times 10^{-27}$	kg
(equivalent to 1.00728u)			
proton charge/mass ratio	$e/m_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$
neutron rest mass	$m_n$	$1.67 \times 10^{-27}$	kg
(equivalent to 1.00867u)			
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$
atomic mass unit	$u$	$1.661 \times 10^{-27}$	kg
(1u is equivalent to 931.3 MeV)			

**Fundamental particles**

Class	Name	Symbol	Rest energy /MeV
photon	photon	$\gamma$	0
lepton	neutrino	$\nu_e$	0
		$\nu_\mu$	0
	electron	$e^\pm$	0.510999
	muon	$\mu^\pm$	105.659
mesons	pion	$\pi^\pm$	139.576
		$\pi^0$	134.972
	kaon	$K^\pm$	493.821
		$K^0$	497.762
baryons	proton	p	938.257
	neutron	n	939.551

**Properties of quarks**

Type	Charge	Baryon number	Strangeness
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1

**Geometrical equations**

$$\text{arc length} = r\theta$$

$$\text{circumference of circle} = 2\pi r$$

$$\text{area of circle} = \pi r^2$$

$$\text{area of cylinder} = 2\pi rh$$

$$\text{volume of cylinder} = \pi r^2 h$$

$$\text{area of sphere} = 4\pi r^2$$

$$\text{volume of sphere} = \frac{4}{3}\pi r^3$$

**Mechanics and Applied Physics**

$$v = u + at$$

$$s = \left(\frac{u+v}{2}\right)t$$

$$s = ut + \frac{at^2}{2}$$

$$v^2 = u^2 + 2as$$

$$F = \frac{\Delta(mv)}{\Delta t}$$

$$P = Fv$$

$$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$$

$$\omega = \frac{v}{r} = 2\pi f$$

$$a = \frac{v^2}{r} = r\omega^2$$

$$I = \sum mr^2$$

$$E_k = \frac{1}{2}I\omega^2$$

$$\omega_2 = \omega_1 + at$$

$$\theta = \omega_1 t + \frac{1}{2}at^2$$

$$\omega_2^2 = \omega_1^2 + 2a\theta$$

$$\theta = \frac{1}{2}(\omega_1 + \omega_2)t$$

$$T = I\alpha$$

$$\text{angular momentum} = I\omega$$

$$W = T\theta$$

$$P = T\omega$$

$$\text{angular impulse} = \text{change of angular momentum} = Tt$$

$$\Delta Q = \Delta U + \Delta W$$

$$\Delta W = p\Delta V$$

$$pV^\gamma = \text{constant}$$

$$\text{work done per cycle} = \text{area of loop}$$

$$\text{input power} = \text{calorific value} \times \text{fuel flow rate}$$

$$\text{indicated power as (area of } p-V \text{ loop)} \times (\text{no. of cycles/s}) \times (\text{no. of cylinders})$$

$$\text{friction power} = \text{indicated power} - \text{brake power}$$

$$\text{efficiency} = \frac{W}{Q_{\text{in}}} = \frac{Q_{\text{in}} - Q_{\text{out}}}{Q_{\text{in}}}$$

$$\text{maximum possible}$$

$$\text{efficiency} = \frac{T_H - T_C}{T_H}$$

**Fields, Waves, Quantum Phenomena**

$$g = \frac{F}{m}$$

$$g = -\frac{GM}{r^2}$$

$$g = -\frac{\Delta V}{\Delta x}$$

$$V = -\frac{GM}{r}$$

$$a = -(2\pi f)^2 x$$

$$v = \pm 2\pi f \sqrt{A^2 - x^2}$$

$$x = A \cos 2\pi ft$$

$$T = 2\pi \sqrt{\frac{m}{k}}$$

$$T = 2\pi \sqrt{\frac{L}{g}}$$

$$\lambda = \frac{\omega s}{D}$$

$$d \sin \theta = n\lambda$$

$$\theta \approx \frac{\lambda}{D}$$

$${}^1n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$$

$${}^1n_2 = \frac{n_2}{n_1}$$

$$\sin \theta_c = \frac{1}{n}$$

$$E = hf$$

$$hf = \phi + E_k$$

$$hf = E_1 - E_2$$

$$\lambda = \frac{h}{p} = \frac{h}{mv}$$

$$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$$

**Electricity**

$$\epsilon = \frac{E}{Q}$$

$$\epsilon = I(R + r)$$

$$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$$

$$R_T = R_1 + R_2 + R_3 + \dots$$

$$P = I^2 R$$

$$E = \frac{F}{Q} = \frac{V}{d}$$

$$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$$

$$E = \frac{1}{2} QV$$

$$F = BI l$$

$$F = BQv$$

$$Q = Q_0 e^{-t/RC}$$

$$\text{magnitude of induced e.m.f.} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

### Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F l}{A e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

### Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

### Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ kms}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_c}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

### Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

### Electronics

#### Resistors

Preferred values for resistors (E24)  
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2  
2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2  
6.8 7.5 8.2 9.1 ohms  
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi f C}$$

### Alternating Currents

$$f = \frac{1}{T}$$

### Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left( \frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$



---

**SECTION A**

In this section each item consists of a question or an incomplete statement followed by four suggested answers or completions. You are to select the most appropriate answer in each case.

---


- 1 A mass  $M$  on a spring oscillates along a vertical line with the same period  $T$  as an object  $O$  in uniform circular motion in a vertical plane. When  $M$  is at its highest point,  $O$  is at its lowest point.



What is the least time interval between successive instants when the acceleration of  $M$  is exactly in the opposite direction to the acceleration of  $O$ ?

- A  $\frac{T}{4}$
- B  $\frac{T}{2}$
- C  $\frac{3T}{4}$
- D  $T$
- 2 A particle of mass  $m$  oscillates with amplitude  $A$  at frequency  $f$ . What is the maximum kinetic energy of the particle?
- A  $\frac{1}{2} \pi^2 m f^2 A^2$
- B  $\pi^2 m f^2 A^2$
- C  $2 \pi^2 m f^2 A^2$
- D  $4 \pi^2 m f^2 A^2$


- 3 The sound quality of a portable radio is improved by adjusting the orientation of the aerial. Which statement is a correct explanation of this improvement?
- A The radio waves from the transmitter are polarised.
  - B The radio waves from the transmitter are unpolarised.
  - C The radio waves become polarised as a result of adjusting the aerial.
  - D The radio waves become unpolarised as a result of adjusting the aerial.
- 4 A microwave transmitter is used to direct microwaves of wavelength 30 mm along a line XY. A metal plate is positioned at right angles to XY with its mid-point on the line, as shown.



When a detector is moved gradually along XY, its reading alternates between maxima and minima. Which one of the following statements is **not** correct?

- A The distance between two minima could be 15 mm.
- B The distance between two maxima could be 30 mm.
- C The distance between a minimum and a maximum could be 30 mm.
- D The distance between a minimum and a maximum could be 37.5 mm.


5







In a double slit system used to produce interference fringes, the separation of the slits is  $s$  and the width of each slit is  $x$ .  $L$  is a source of monochromatic light. Which one of the following changes would **decrease** the separation of the fringes seen on the screen?

- A moving the screen closer to the double slits
- B decreasing the width,  $x$ , of each slit, but keeping  $s$  constant
- C decreasing the separation,  $s$ , of the slits
- D exchanging  $L$  for a monochromatic source of longer wavelength

6




The diagram above shows the first four diffraction orders each side of the zero order when a beam of monochromatic light is incident normally on a diffraction grating of slit separation  $d$ . All the angles of diffraction are small. Which one of the patterns, **A** to **D**, drawn on the same scale, is obtained when the grating is exchanged for one with a slit separation  $\frac{d}{2}$ ?


**A****B****C****D**




- 7 A  $1000\ \mu\text{F}$  capacitor, initially uncharged, is charged by a steady current of  $50\ \mu\text{A}$ . How long will it take for the potential difference across the capacitor to reach  $2.5\ \text{V}$ ?
- A 20 s  
B 50 s  
C 100 s  
D 400 s
- 8 In experiments to pass a very high current through a gas, a bank of capacitors of total capacitance  $50\ \mu\text{F}$  is charged to  $30\ \text{kV}$ . If the bank of capacitors could be discharged completely in  $5.0\ \text{ms}$  what would be the mean power delivered?
- A 22 kW  
B 110 kW  
C 4.5 MW  
D 9.0 MW
- 9 For a particle moving in a circle with uniform speed, which **one** of the following statements is correct?
- A The displacement of the particle is in the direction of the force.  
B The force on the particle is in the same direction as the direction of motion of the particle.  
C The momentum of the particle is constant.  
D The kinetic energy of the particle is constant.
- 10 Which one of the following graphs correctly shows the relationship between the gravitational force,  $F$ , between two masses and their separation  $r$ .




A



B



C

D [www.theallpapers.com](http://www.theallpapers.com)

- 11 When at the surface of the Earth, a satellite has weight  $W$  and gravitational potential energy  $-U$ . It is projected into a circular orbit whose radius is equal to twice the radius of the Earth. Which line, **A** to **D**, in the table shows correctly what happens to the weight of the satellite and to its gravitational potential energy?

	weight	gravitational potential energy
<b>A</b>	becomes $\frac{W}{2}$	increases by $\frac{U}{2}$
<b>B</b>	becomes $\frac{W}{4}$	increases by $\frac{U}{2}$
<b>C</b>	remains $W$	increases by $U$
<b>D</b>	becomes $\frac{W}{4}$	increases by $U$

- 12 Two protons are  $1.0 \times 10^{-14}$  m apart. Approximately how many times is the electrostatic force between them greater than the gravitational force between them?

**A**  $10^{23}$

**B**  $10^{30}$

**C**  $10^{36}$

**D**  $10^{42}$

- 13 Particles of mass  $m$  carrying a charge  $Q$  travel in a circular path of radius  $r$  in a magnetic field of flux density  $B$  with a speed  $v$ . How many of the following quantities, if changed one at a time, would change the radius of the path?

- $m$
- $Q$
- $B$
- $v$


**A** one

**B** two

**C** three

**D** four

- 14 In the reaction shown, a proton and a deuterium nucleus,  ${}^2_1\text{H}$ , fuse together to form a helium nucleus,  ${}^3_2\text{He}$



What is the value of Q, the energy released in this reaction?

$$\text{mass of a proton} = 1.00728 \text{ u}$$

$$\text{mass of a } {}^2_1\text{H nucleus} = 2.01355 \text{ u}$$

$$\text{mass of a } {}^3_2\text{He nucleus} = 3.01493 \text{ u}$$

- A 5.0 MeV  
B 5.5 MeV  
C 6.0 MeV  
D 6.5 MeV
- 15 For a nuclear reactor in which the fission rate is constant, which one of the following statements is correct?
- A There is a critical mass of fuel in the reactor.  
B For every fission event, there is, on average, one further fission event.  
C A single neutron is released in every fission event.  
D No neutrons escape from the reactor.

**END OF SECTION A**

**There are no questions printed on this page**