



ASSESSMENT and
QUALIFICATIONS
ALLIANCE

Mark scheme

June 2003

GCE

Physics A


Unit PHA5/W

Copyright © 2003 AQA and its licensors. All rights reserved.

Units 5 - 9 : Section A

1

(a)(i)



correct arrows: A ✓

B ✓
C ✓

(a)(ii) $e^{-1} + {}^A_ZX \rightarrow {}^A_{Z-1}Y + v_e$ ✓ (4)

(b)(i) $((4.18 - 1.33) \times 10^{-13}) = 2.85 \times 10^{-13}$ (J) ✓

(b)(ii) 1.33×10^{-13} (J)
 0.30×10^{-13} (J) for 3 correct values ✓
 1.63×10^{-13} (J)

(b)(iii) (use of $\Delta E = hf$ gives) $f \left(= \frac{1.63 \times 10^{-13}}{6.63 \times 10^{-34}} \right) = 2.46 \times 10^{20}$ Hz ✓
 (allow C.E. from (b)(ii) if largest value taken) (3)

(c)(i) (✓ for each precaution with reason to $_{\max}2$)

handle with (long) (30 cm) tweezers
 because the radiation intensity decreases with distance

store in a lead box (immediately) when not in use
 to avoid unnecessary exposure to radiation

[or any sensible precaution with reason]

(b)(ii) γ rays are more penetrating and are therefore more hazardous
 (to the internal organs of the body)

β^- particles are more hazardous because they are more ionising ✓
 (✓ for any argued case for either radiation)

(3)
(10)

Unit 5 : Section B

2

- (a) ray diagram to show:
 rays reflected at concave mirror ✓
 rays reflected at convex mirror ✓
 rays crossing in front of eyepiece ✓ (3)
- (b) different focal points for rays at different distances from axis ✓
 shortest focal length for paraxial rays ✓ (2)
- (c) light of different wavelengths refracted to different foci ✓
 diagram showing refraction with blue focal length closest to lens ✓
 max (2)
 (7)

3

- (a)(i) $d = \frac{50 \times 10^6}{3.26} = 15.3 \times 10^6 \text{ (pc)} \checkmark$
- (a)(ii) (use of $v = Hd$ gives) $v = 65 \times 10^{-6} \text{ (km s}^{-1} \text{ pc}^{-1}) \times 15.3 \times 10^6 \checkmark$
 $\approx (1000 \text{ km s}^{-1})$
- (a)(iii) (use of $\frac{\Delta\lambda}{\lambda} = -\frac{v}{c}$ gives) $\Delta\lambda = \frac{1000 \times 10^3}{3 \times 10^8} \times 656.3 \text{ (nm)} = 2.19 \text{ (nm)} \checkmark$
 (allow C.E. for value of v from (ii))
 $\lambda_{\text{galaxy}} = 656.3 + 2.19 = 658.5 \text{ nm} \checkmark$ (4)
- (b) for the furthest point of the Universe, $d = \frac{c}{H} \checkmark$
 age of Universe = $\frac{d}{c} = \frac{1}{H} \checkmark$
 [or use of $v = Hd$ and $t = \frac{d}{v} \checkmark$
 if all started from same point
 $t = \text{age of Universe} = \frac{1}{H} \checkmark$]
 assumption: that H remains constant ✓ (3)
 (7)

4

(a) Hertzsprung -Russell diagram to show:
absolute magnitude scale from +15 to -10 ✓
temperature scale from 50 000 to 2500 (K) ✓
main sequence drawn correctly ✓
giants and dwarfs shown in correct areas ✓ (4)

(b) Alnitak : helium (absorption)
Sirius : hydrogen Balmer (absorption) lines 4 correct ✓✓
Sun : metals (absorption) 2 correct ✓
Antares : molecular bands (2)

(c) reference to $P = \sigma AT^4$ ✓
class M (Antares) cooler than class O (Alnitak) ✓
but same brightness, therefore cooler star bigger ✓
so Antares has larger surface area ✓ max.(3)
(9)

5

(a)(i) supernova: star whose luminosity increase enormously
due to it exploding ✓
(a)(ii) neutron star: star with the density of nuclear matter ✓
(a)(iii) black hole: an object whose escape velocity is greater
than speed of light ✓ (3)

(b) $\left(\text{use of } R = \frac{2GM}{c^2} \text{ gives} \right) R = \frac{2 \times 6.67 \times 10^{-11} \times 10 \times 2 \times 10^{30}}{(3 \times 10^8)^2} \checkmark$
 $= 2.96 \times 10^4 \text{ m } \checkmark$ (2)
(5)

Quality of Written Communication (Q1(c)(i) and Q4(c)) ✓✓ (2)
(2)