

Surname		Other Names	
Centre Number		Candidate Number	
Candidate Signature			

For Examiner's Use
--------------------

General Certificate of Education  
June 2007  
Advanced Level Examination



ASSESSMENT and  
QUALIFICATIONS  
ALLIANCE

**PHYSICS (SPECIFICATION A)**  
**Unit 5 Nuclear Instability: Astrophysics Option**

**PHA5/W**

Thursday 14 June 2007 9.00 am to 10.15 am

<p><b>For this paper you must have:</b></p> <ul style="list-style-type: none"> <li>• a calculator</li> <li>• a pencil and a ruler.</li> </ul>
---

For Examiner's Use			
Question	Mark	Question	Mark
1			
2			
3			
4			
5			
Total (Column 1) →			
Total (Column 2) →			
Quality of Written Communication			
TOTAL			
Examiner's Initials			

Time allowed: 1 hour 15 minutes

**Instructions**

- Use blue or black ink or ball-point pen.
- Fill in the boxes at the top of this page.
- Answer **all** questions.
- Answer the questions in the spaces provided.
- Show all your working.
- Do all rough work in this book. Cross through any work you do not want to be marked.

**Information**

- The maximum mark for this paper is 40.
- Two of these marks will be awarded for using good English, organising information clearly and using specialist vocabulary where appropriate.
- The marks for questions are shown in brackets.
- A *Data Sheet* is provided on pages 3 and 4. You may wish to detach this perforated sheet at the start of the examination.
- You are expected to use a calculator where appropriate.
- Questions 1(a) and 3(a) should be answered in continuous prose. In these questions you will be marked on your ability to use good English, to organise information clearly and to use specialist vocabulary where appropriate.

**Data Sheet**

- A perforated *Data Sheet* is provided as pages 3 and 4 of this question paper.
- This sheet may be useful for answering some of the questions in the examination.
- You may wish to detach this sheet before you begin work.

## Data Sheet

Fundamental constants and values				Mechanics and Applied Physics		Fields, Waves, Quantum Phenomena	
Quantity	Symbol	Value	Units				
speed of light in vacuo	$c$	$3.00 \times 10^8$	$\text{m s}^{-1}$	$v = u + at$		$g = \frac{F}{m}$	
permeability of free space	$\mu_0$	$4\pi \times 10^{-7}$	$\text{H m}^{-1}$	$s = \left(\frac{u+v}{2}\right)t$		$g = -\frac{GM}{r^2}$	
permittivity of free space	$\epsilon_0$	$8.85 \times 10^{-12}$	$\text{F m}^{-1}$	$s = ut + \frac{at^2}{2}$		$g = -\frac{\Delta V}{\Delta x}$	
charge of electron	$e$	$1.60 \times 10^{-19}$	C	$v^2 = u^2 + 2as$		$V = -\frac{GM}{r}$	
the Planck constant	$h$	$6.63 \times 10^{-34}$	J s	$F = \frac{\Delta(mv)}{\Delta t}$		$a = -(2\pi f)^2 x$	
gravitational constant	$G$	$6.67 \times 10^{-11}$	$\text{N m}^2 \text{kg}^{-2}$	$P = Fv$		$v = \pm 2\pi f \sqrt{A^2 - x^2}$	
the Avogadro constant	$N_A$	$6.02 \times 10^{23}$	$\text{mol}^{-1}$	$\text{efficiency} = \frac{\text{power output}}{\text{power input}}$		$x = A \cos 2\pi ft$	
molar gas constant	$R$	8.31	$\text{J K}^{-1} \text{mol}^{-1}$	$\omega = \frac{v}{r} = 2\pi f$		$T = 2\pi \sqrt{\frac{m}{k}}$	
the Boltzmann constant	$k$	$1.38 \times 10^{-23}$	$\text{J K}^{-1}$	$a = \frac{v^2}{r} = r\omega^2$		$T = 2\pi \sqrt{\frac{L}{g}}$	
the Stefan constant	$\sigma$	$5.67 \times 10^{-8}$	$\text{W m}^{-2} \text{K}^{-4}$	$I = \sum mr^2$		$\lambda = \frac{\omega s}{D}$	
the Wien constant	$a$	$2.90 \times 10^{-3}$	m K	$E_k = \frac{1}{2} I\omega^2$		$d \sin \theta = n\lambda$	
electron rest mass	$m_e$	$9.11 \times 10^{-31}$	kg	$\omega_2 = \omega_1 + at$		$\theta \approx \frac{\lambda}{D}$	
(equivalent to $5.5 \times 10^{-4} \text{u}$ )				$\theta = \omega_1 t + \frac{1}{2} at^2$		$n_2 = \frac{\sin \theta_1}{\sin \theta_2} = \frac{c_1}{c_2}$	
electron charge/mass ratio	$e/m_e$	$1.76 \times 10^{11}$	$\text{C kg}^{-1}$	$\omega_2^2 = \omega_1^2 + 2a\theta$		$n_2 = \frac{n_2}{n_1}$	
proton rest mass	$m_p$	$1.67 \times 10^{-27}$	kg	$\theta = \frac{1}{2} (\omega_1 + \omega_2)t$		$\sin \theta_c = \frac{1}{n}$	
(equivalent to 1.00728u)				$T = I\alpha$		$E = hf$	
proton charge/mass ratio	$e/m_p$	$9.58 \times 10^7$	$\text{C kg}^{-1}$	$\text{angular momentum} = I\omega$		$hf = \phi + E_k$	
neutron rest mass	$m_n$	$1.67 \times 10^{-27}$	kg	$W = T\theta$		$hf = E_1 - E_2$	
(equivalent to 1.00867u)				$P = T\omega$		$\lambda = \frac{h}{p} = \frac{h}{mv}$	
gravitational field strength	$g$	9.81	$\text{N kg}^{-1}$	$\text{angular impulse} = \text{change of angular momentum} = Tt$		$c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$	
acceleration due to gravity	$g$	9.81	$\text{m s}^{-2}$	$\Delta Q = \Delta U + \Delta W$		<b>Electricity</b>	
atomic mass unit	$u$	$1.661 \times 10^{-27}$	kg	$\Delta W = p\Delta V$		$\epsilon = \frac{E}{Q}$	
(1u is equivalent to 931.3 MeV)				$pV^\gamma = \text{constant}$		$\epsilon = I(R + r)$	
<b>Fundamental particles</b>				$\text{work done per cycle} = \text{area of loop}$		$\frac{1}{R_T} = \frac{1}{R_1} + \frac{1}{R_2} + \frac{1}{R_3} + \dots$	
<b>Class</b>	<b>Name</b>	<b>Symbol</b>	<b>Rest energy</b>	$\text{input power} = \text{calorific value} \times \text{fuel flow rate}$		$R_T = R_1 + R_2 + R_3 + \dots$	
			/MeV	$\text{indicated power as (area of } p-V \text{ loop)} \times (\text{no. of cycles/s}) \times (\text{no. of cylinders})$		$P = I^2 R$	
photon	photon	$\gamma$	0	$\text{friction power} = \text{indicated power} - \text{brake power}$		$E = \frac{F}{Q} = \frac{V}{d}$	
lepton	neutrino	$\nu_e$	0	$\text{efficiency} = \frac{W}{Q_{in}} = \frac{Q_{in} - Q_{out}}{Q_{in}}$		$E = \frac{1}{4\pi\epsilon_0} \frac{Q}{r^2}$	
		$\nu_\mu$	0	$\text{maximum possible efficiency} = \frac{T_H - T_C}{T_H}$		$E = \frac{1}{2} QV$	
mesons	electron	$e^\pm$	0.510999			$F = BIl$	
		$\mu^\pm$	105.659			$F = BQv$	
	pion	$\pi^\pm$	139.576			$Q = Q_0 e^{-t/RC}$	
		$\pi^0$	134.972			$Q = Q_0 e^{-t/RC}$	
kaon	$K^\pm$	493.821					
	$K^0$	497.762					
baryons	proton	p	938.257				
	neutron	n	939.551				
<b>Properties of quarks</b>							
<b>Type</b>	<b>Charge</b>	<b>Baryon number</b>	<b>Strangeness</b>				
u	$+\frac{2}{3}$	$+\frac{1}{3}$	0				
d	$-\frac{1}{3}$	$+\frac{1}{3}$	0				
s	$-\frac{1}{3}$	$+\frac{1}{3}$	-1				
<b>Geometrical equations</b>							
arc length = $r\theta$							
circumference of circle = $2\pi r$							
area of circle = $\pi r^2$							
area of cylinder = $2\pi rh$							
volume of cylinder = $\pi r^2 h$							
area of sphere = $4\pi r^2$							
volume of sphere = $\frac{4}{3}\pi r^3$							

$$\text{magnitude of induced emf} = N \frac{\Delta\Phi}{\Delta t}$$

$$I_{\text{rms}} = \frac{I_0}{\sqrt{2}}$$

$$V_{\text{rms}} = \frac{V_0}{\sqrt{2}}$$

### Mechanical and Thermal Properties

$$\text{the Young modulus} = \frac{\text{tensile stress}}{\text{tensile strain}} = \frac{F}{A} \frac{l}{e}$$

$$\text{energy stored} = \frac{1}{2} Fe$$

$$\Delta Q = mc \Delta\theta$$

$$\Delta Q = ml$$

$$pV = \frac{1}{3} Nmc^2$$

$$\frac{1}{2} mc^2 = \frac{3}{2} kT = \frac{3RT}{2N_A}$$

### Nuclear Physics and Turning Points in Physics

$$\text{force} = \frac{eV_p}{d}$$

$$\text{force} = Bev$$

$$\text{radius of curvature} = \frac{mv}{Be}$$

$$\frac{eV}{d} = mg$$

$$\text{work done} = eV$$

$$F = 6\pi\eta rv$$

$$I = k \frac{I_0}{x^2}$$

$$\frac{\Delta N}{\Delta t} = -\lambda N$$

$$\lambda = \frac{h}{\sqrt{2meV}}$$

$$N = N_0 e^{-\lambda t}$$

$$T_{\frac{1}{2}} = \frac{\ln 2}{\lambda}$$

$$R = r_0 A^{\frac{1}{3}}$$

$$E = mc^2 = \frac{m_0 c^2}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

$$l = l_0 \left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}$$

$$t = \frac{t_0}{\left(1 - \frac{v^2}{c^2}\right)^{\frac{1}{2}}}$$

### Astrophysics and Medical Physics

Body	Mass/kg	Mean radius/m
Sun	$2.00 \times 10^{30}$	$7.00 \times 10^8$
Earth	$6.00 \times 10^{24}$	$6.40 \times 10^6$

$$1 \text{ astronomical unit} = 1.50 \times 10^{11} \text{ m}$$

$$1 \text{ parsec} = 206265 \text{ AU} = 3.08 \times 10^{16} \text{ m} = 3.26 \text{ ly}$$

$$1 \text{ light year} = 9.45 \times 10^{15} \text{ m}$$

$$\text{Hubble constant } (H) = 65 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

$$M = \frac{\text{angle subtended by image at eye}}{\text{angle subtended by object at unaided eye}}$$

$$M = \frac{f_o}{f_e}$$

$$m - M = 5 \log \frac{d}{10}$$

$$\lambda_{\text{max}} T = \text{constant} = 0.0029 \text{ m K}$$

$$v = Hd$$

$$P = \sigma AT^4$$

$$\frac{\Delta f}{f} = \frac{v}{c}$$

$$\frac{\Delta \lambda}{\lambda} = -\frac{v}{c}$$

$$R_s \approx \frac{2GM}{c^2}$$

### Medical Physics

$$\text{power} = \frac{1}{f}$$

$$\frac{1}{u} + \frac{1}{v} = \frac{1}{f} \text{ and } m = \frac{v}{u}$$

$$\text{intensity level} = 10 \log \frac{I}{I_0}$$

$$I = I_0 e^{-\mu x}$$

$$\mu_m = \frac{\mu}{\rho}$$

### Electronics

#### Resistors

Preferred values for resistors (E24)  
Series: 1.0 1.1 1.2 1.3 1.5 1.6 1.8 2.0 2.2 2.4 2.7 3.0 3.3 3.6 3.9 4.3 4.7 5.1 5.6 6.2 6.8 7.5 8.2 9.1 ohms  
and multiples that are ten times greater

$$Z = \frac{V_{\text{rms}}}{I_{\text{rms}}}$$

$$\frac{1}{C_T} = \frac{1}{C_1} + \frac{1}{C_2} + \frac{1}{C_3} + \dots$$

$$C_T = C_1 + C_2 + C_3 + \dots$$

$$X_C = \frac{1}{2\pi fC}$$

### Alternating Currents

$$f = \frac{1}{T}$$

### Operational amplifier

$$G = \frac{V_{\text{out}}}{V_{\text{in}}} \quad \text{voltage gain}$$

$$G = -\frac{R_f}{R_1} \quad \text{inverting}$$

$$G = 1 + \frac{R_f}{R_1} \quad \text{non-inverting}$$

$$V_{\text{out}} = -R_f \left( \frac{V_1}{R_1} + \frac{V_2}{R_2} + \frac{V_3}{R_3} \right) \quad \text{summing}$$

**Turn over for the first question**

## SECTION A: NUCLEAR INSTABILITY

Answer **all** of this question.

- 1 (a) X and Y are two different  $\beta$  emitting sources. Initially they contain the same number of unstable nuclei. Both sources have their emissions recorded over a period of time. The *decay constant* of source X is greater than that of Y. State what is meant by decay constant and describe **two** differences in the recordings from the two sources.

You may be awarded additional marks to those shown in brackets for the quality of written communication in your answer.

.....

.....


.....

.....

.....

(3 marks)

- (b) The activity of a sample of radioactive iodine,  $^{131}_{53}\text{I}$ , is presented in the following graph.



- (i) Show that the decay constant of  $^{131}_{53}\text{I}$  is about  $1 \times 10^{-6} \text{ s}^{-1}$ .

.....

.....

.....

.....

.....

- (ii) Calculate the half-life of  $^{131}_{53}\text{I}$  in days.

.....

.....

.....

- (iii) Calculate the initial number of  $^{131}_{53}\text{I}$  atoms in the sample.

.....

.....

.....

.....

(7 marks)

**Turn over for the next question**

10
----

**SECTION B: ASTROPHYSICS**Answer **all** questions.

- 2 (a) A lens used by Galileo has a range of focal lengths from 0.98 m to 0.92 m, depending on the wavelength of the light passing through the lens.

- (i) Calculate the power of the lens for red light.

.....  
 .....

- (ii) Name the defect in the image which arises because a lens has different focal lengths for different wavelengths of light.

.....

*(3 marks)*

- (b) The telescope with which Galileo discovered Io, one of the satellites of Jupiter, had an angular magnification of 30. Calculate the maximum angular separation of the images of Io and Jupiter when viewed through this telescope.

$$\begin{aligned} \text{radius of the orbit of Io around Jupiter} &= 4.2 \times 10^5 \text{ km} \\ \text{least distance of Jupiter from the Earth} &= 6.0 \times 10^8 \text{ km} \end{aligned}$$

.....  
 .....

*(2 marks)*

- (c) A lens of focal length 0.95 m is used as the objective of an astronomical telescope. In normal adjustment, the telescope has an angular magnification of 30. Calculate the distance between the objective and eyepiece lenses.

.....  
 .....

*(2 marks)*



3 Stars of spectral classes A and B have strong hydrogen Balmer absorption lines in their spectra.

(a) Describe how Balmer absorption lines are produced.

You may be awarded additional marks to those shown in brackets for the quality of written communication in your answer.

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

(4 marks)

(b) (i) Why do the spectra of stars in classes F and G not have strong Balmer absorption lines?

.....

.....

.....

(ii) What is the prominent feature in the spectra of stars in classes F and G?

.....

(2 marks)

6
---

4 The data in the table gives some of the properties of the star Mu Cephei.

apparent magnitude	4.23
absolute magnitude	-6.81
surface temperature	3500 K

(a) (i) Calculate the wavelength of the peak in the black body radiation curve for Mu Cephei.

.....

.....

.....

.....

(ii) Sketch the black body radiation curve for Mu Cephei on the axes below. Label the wavelength axis with a suitable scale.



(3 marks)

(b) Calculate the distance from the Earth to Mu Cephei in light years.

.....

.....

.....

.....

.....

.....

(2 marks)

- (c) Mu Cephei is possibly the largest star yet discovered. Its radius is  $1.2 \times 10^9$  km, which is about the orbital radius of Saturn. Show that the power output of Mu Cephei is approximately 400 000 times that of the Sun.

$$\begin{aligned} \text{surface temperature of the Sun} &= 5800 \text{ K} \\ \text{radius of the Sun} &= 6.9 \times 10^5 \text{ km} \end{aligned}$$

.....

.....

.....

.....

.....

(2 marks)

7

**Turn over for the next question**

5 Tonantzintla 202 is a *quasar* with a red shift,  $\frac{\Delta f}{f}$ , of 0.366. When it was discovered in 1957 it was wrongly assumed to be a *white dwarf*.

(a) Explain what is meant by

(i) a white dwarf,

.....

.....

.....

.....

.....

(ii) a quasar.

.....

.....

.....

.....

.....

(5 marks)

(b) Ignoring relativistic effects, calculate for Tonantzintla 202,

(i) its recessional speed, relative to the Earth,

.....

.....

.....

(ii) its distance from the Earth.

.....

.....

.....

*(3 marks)*

<b>8</b>

**Quality of Written Communication** *(2 marks)*

<b>2</b>

**END OF QUESTIONS**

**There are no questions printed on this page**

**There are no questions printed on this page**

**There are no questions printed on this page**