

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education Advanced Subsidiary Level and Advanced Level

	CANDIDATE NAME							
	CENTRE NUMBER					CANDIDATE NUMBER		
*	BIOLOGY Advanced Prac	tical Skills	1				Мау	9700/35 //June 2013 2 hours
	Candidates ans	swer on th	e Question	Paper.				
	Additional Mate	erials:	As listed in	the Confidential	Instructions.			
*	READ THESE I	INSTRUC	TIONS FIR	ST				

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black ink. You may use a pencil for any diagrams, graphs or rough working.

Do **not** use red ink, staples, paper clips, highlighters, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions. Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use			
1			
2			
Total			

This document consists of **15** printed pages and **1** blank page.



UNIVERSITY of CAMBRIDGE International Examinations

[Turn over

www.theallpapers.com

You are reminded that you have **only one hour** for each question in the practical examination.

You should:

- read carefully through the whole of Question 1 and Question 2
- then plan your use of **the time** to make sure that you finish all the work that you would like to do.

You will gain marks for recording your results according to the instructions.

1 Glucose solutions change the colour of pink potassium manganate(VII) solution, **PM**.

Fig. 1.1 shows the colour change from pink to the colourless end-point.



Fig. 1.1

The rate of the colour change depends on the concentration of the glucose solution. The greater the concentration of glucose solution the faster the end-point is reached.

You are required to:

- make different concentrations of glucose solution
- find, for each glucose solution, the time taken for PM to change to colourless
- estimate the unknown concentrations of the glucose solutions, **U1** and **U2**.

You are provided with:

labelled	contents	hazard	volume/cm ³
G	20% glucose solution	none	100
W	distilled water	none	200
S	sulfuric acid	harmful	40
РМ	potassium manganate(VII) solution	harmful	20
U1	glucose solution	none	20
U2	glucose solution	none	20

Sulfuric acid and potassium manganate(VII) solution are harmful.

If any comes into contact with your skin, wash immediately under cold water. It is recommended that you wear safety goggles/glasses.

Proceed as follows:

- Using the 20% glucose solution, G, as a starting concentration you are required to make up 20 cm³ of each of four different concentrations of glucose solutions, 6%, 8%, 10%, 12%.
- (a) (i) Complete Table 1.1 to show how you will make the four glucose solutions 6%, 8%, 10% and 20%.

volume of 20% glucose solution/cm ³	volume of distilled water/cm ³	final percentage concentration of glucose
		6
		8
		10
12	8	12

Table 1.1

[2]

- 2. Make all the glucose solutions as in Table 1.1, in the containers provided.
- 3. Put 10 cm³ of each glucose solution into four separate test-tubes.
- 4. Using the syringe labelled **S**, put 5 cm³ of **S** into each test-tube. Insert the bung and, with your finger holding the bung in place, gently mix the solution in each test-tube. Do **not** turn the test-tube upside-down.

[Turn over www.theallpapers.com

When adding **PM** to the first glucose solution, you must **not** stop the timer, **just record** the time. When adding PM to the other glucose solutions, or at any of the end-points, Examiner's do **not** stop the timer, just record the time.

From your timer readings you will be required to calculate the time taken to reach the end-point in each test-tube.

(ii) Consider the units of the values recorded on your timer or clock.


State the:

- smallest value which your timer or clock shows
- smallest unit of time you have decided to record

Read up to step 11 before proceeding.

Proceed as follows:

Using the syringe labelled **PM**, put 2 cm³ of **PM** into the test-tube containing the 5. lowest concentration of glucose solution as shown in Fig. 1.2.





- Start timing and record the start time from your timer on Fig. 1.3 on page 5. 6.
- Immediately, put 2 cm³ of **PM** into the test-tube containing next highest concentration 7. of glucose solution.
- 8. Record start time from your timer on Fig. 1.3 on page 5.
- 9. Immediately, repeat steps 7 and 8 for the remaining concentrations of glucose solution.

For

Use

[1]

10. Observe the four test-tubes and record the time on Fig. 1.3 when each end-point is reached.






Fig. 1.3


[Turn over www.theallpapers.com

You are required to estimate the glucose concentration of solutions, U1 and U2 using the same procedure. Examiner's


For

Use

State one variable, which you will standardise when setting up the test-tubes to find (iii) the end-points for U1 and U2.



11. Use the same procedure to obtain the end-points for the solutions, U1 and U2 and record your times on Fig. 1.4.



6

Depending on the timer or clock you have used, you may find the following examples helpful so that you can process your results for **(v)** and for Step 12 to find the time taken to reach the end-point.

For Examiner's Use

Example 1: using stop-clock or stopwatch

start time end-point time	minutes:seconds 1:24 2:55	= 84 seconds = 175 seconds
time	taken to reach end-point	= 91 seconds

Example 2: using clock times

start time end-point time	hours:minutes:seconds 9:10:00 9:11:31	difference in time 1 minute 31 seconds			
time taken to reach end-point = 91 seconds					

(v) Using your results from Fig. 1.3, complete Table 1.2 to show the calculation to find the time taken for 6% glucose solution to reach the end-point.



6% solution start time 6% solution end-point time		
ti	me taken to reach end-point	=

12. Use the space on Fig. 1.3 and Fig. 1.4 for processing your readings to find the time taken to reach the end-point for the other glucose solutions, and for **U1** and **U2**.

(vi) Using **only** these processed results, prepare the space below to record the time taken to reach the end-point for **all six** solutions.

For Examiner's Use

[4]

Glucose solutions may be used for different purposes, for example:
Glucose tolerance test solutions, containing 25% glucose.
Sports drink solutions, containing 8% glucose.
Oral Rehydration solutions, containing 2% glucose.
(b) Suggest which of the above solutions is U2.

(c) (i) Identify one significant source of error in your investigation.
 [1]
 (ii) Describe two modifications to this investigation which would improve the confidence in your results.
 [1]
 (ii) Describe two modifications to this investigation which would improve the confidence in your results.

Question 2 starts on page 10

2 Fig. 2.1 shows a photomicrograph of a transverse section through part of a plant stem showing an eyepiece graticule scale as seen using a microscope.

For Examiner's Use




Fig. 2.1

An eyepiece graticule scale can be used to measure the layers of tissues and to help draw a plan diagram with the correct shape and proportions of the tissues, without needing to calibrate the eyepiece graticule scale.

(a) (i) The length of the vascular bundle (from K to R) in Fig. 2.1 was measured using the eyepiece graticule scale and recorded in Table 2.1.

layer	J	М	Ν	Р	length from K to R
number of eyepiece graticule scale divisions					43

Complete Table 2.1 by finding the thickness of the different layers L, M, N and P, labelled in Fig. 2.1, using the line between R and K and the eyepiece graticule scale. [2]

The length (from K to R) of the vascular bundle in eyepiece graticule divisions was used to make a scale drawing of the outline of the vascular bundle as shown in Examiner's Fig. 2.2.

R-Κ



- (ii) Complete the plan diagram of this part of the vascular bundle to show the proportions and shape of each of the tissues. Use the values in Table 2.1 to help you. [3]
- (iii) Using Fig. 2.2, count the total number of 1 cm by 1 cm squares occupied by the vascular bundle and count the total number of 1 cm by 1 cm squares occupied by the xylem tissue.

Count any 'half square' or 'more than half' as one square.

State the ratio of the area occupied by the vascular bundle to that of the xylem tissue.

You will lose marks if you do not show all the steps in finding the ratio, including indicating counted squares on Fig. 2.2.

[Turn over

ratio[2]

11

Use

For

L1 is a slide of a transverse section through the same plant stem as in Fig. 2.1. This plant grows mainly in Europe and Asia.

This stem shows a stained tissue, close to the epidermis, in the four corners of the stem. Near to the centre of the stem is a different tissue.

(b) Make a large drawing of one group of three whole, adjacent (touching) cells

• from the tissue in one corner, as observed on the specimen on L1.

Make a large drawing of one group of three whole, adjacent cells

• from the tissue near to the centre of the stem, as observed on the specimen on L1.

The drawings should show any difference in size (linear magnification) observed between each group of cells.

On your drawing, use a ruled label line and label to show one cell wall.

cells from the tissue in one corner

cells from the tissue near to the centre of the stem

Fig. 2.3 is a photomicrograph of a transverse section through a stem of a different plant species.

13

For Examiner's Use




Fig. 2.3

- (c) Prepare the space below so that it is suitable for you to record observable differences between the specimens on slide L1 and in Fig. 2.3 to include:
 - the vascular tissue
 - at least two other tissues.

[4]

Some scientists carried out an investigation into the **uptake** of glucose by five different types of plant tissues during the course of 25 minutes.

A piece of each type of plant tissue was placed in a solution of glucose.


The starting concentration of this glucose solution was 0.8 arbitrary units which was lower than the concentration inside the plant cells in each tissue.

The results after 25 minutes are shown in Table 2.2.

type of plant tissue	concentration of glucose in the cells/arbitrary units
А	2.0
В	6.5
С	4.2
D	5.6
E	3.2

Table 2.2

(d) (i) Plot a chart of the data shown in Table 2.2.



[4]

(ii)	Describe and explain the results shown in the chart you have drawn.	For Examiner's
		Use
	[3]	
	[Total: 23]	

BLANK PAGE

16

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.