

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS General Certificate of Education

Advanced Subsidiary Level and Advanced Level

CANDIDATE NAME		
CENTRE NUMBER	CANDIDATE NUMBER	

CHEMISTRY 9701/33

Advanced Practical Skills 1

May/June 2013

2 hours

Candidates answer on the Question Paper.

Additional Materials: As listed

As listed in the Confidential Instructions

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Give details of the practical session and laboratory where appropriate, in the boxes provided.

Write in dark blue or black pen.

You may use a soft pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Electronic calculators may be used.

You may lose marks if you do not show your working or if you do not use appropriate units.

Use of a Data Booklet is unnecessary.

Qualitative Analysis Notes are printed on pages 12 and 13.

A Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

;	Session
La	aboratory

For Exam	iner's Use
1	
2	
3	
Total	

This document consists of 14 printed pages and 2 blank pages.



1 You are to determine the enthalpy change of the reaction between hydrochloric acid and sodium hydroxide by adding various volumes of acid and alkali and measuring the change in temperature.

**FA 1** is 0.950 mol dm<sup>-3</sup> hydrochloric acid, HC*l*.

FA 2 is aqueous sodium hydroxide, NaOH.

#### (a) Method

- Support the plastic cup in a 250 cm³ beaker.
- Using a measuring cylinder, transfer 25 cm³ of **FA 1** into the cup and measure the temperature of the acid. Tilt the cup if necessary to cover the bulb of the thermometer.
- Record this initial temperature.

initial temperature of **FA 1** = .....°C

- Use a second measuring cylinder to transfer 10 cm³ of **FA 2** and 25 cm³ of water into a 100 cm³ beaker.
- Add this mixture to the plastic cup and stir.
- Measure the maximum temperature reached and record this maximum temperature in the table below.
- Rinse out the plastic cup and shake it to remove excess water.
- Repeat the experiment, using the volumes of **FA 1**, **FA 2** and water shown in the table. Record the maximum temperature for each experiment.

volume <b>FA 1</b> /cm³	volume <b>FA 2</b> /cm <sup>3</sup>	volume water/cm <sup>3</sup>	maximum temperature/°C
25	10	25	
25	15	20	
25	20	15	
25	25	10	
25	30	5	
25	35	0	


You are going to plot a graph using these results to find the volume of **FA 2** that gives the greatest maximum temperature.

**Before** you plot the graph, choose two further volumes of **FA 2** that will allow you to find more precisely the volume that gives the greatest maximum temperature.

Record the volumes you choose, carry out the experiments and record the corresponding maximum temperatures, in the table. [2]

**(b) (i)** On the grid below, plot the maximum temperature on the *y*-axis against the volume of **FA 2** on the *x*-axis.

For Examiner's Use



- (ii) Draw two straight lines of best fit on your graph, one to show where the temperature was increasing and the other after the greatest maximum temperature had been reached.
- (iii) Using your graph and the initial temperature recorded in (a), determine the maximum temperature change that could occur when 25 cm³ of FA 1 react with FA 2.

maximum temperature **change** = ......°C

[5]

		_		
(c)	Cal	cul	lati	on

(i)	Calculate the energy needed to produce the temperature change in <b>(b)(iii)</b> . (Assume that 4.3 J of heat energy changes the temperature of $1.0\mathrm{cm^3}$ of solution by $1.0\mathrm{^\circ C.}$ )
	energy needed = J
(ii)	Calculate the number of moles of HC1 used in each experiment.
	moles of $HCl = \dots mol$
(iii)	Calculate the enthalpy change, in $kJ  mol^{-1}$ , when 1 mole of $HC\it{l}$ reacts with NaOH.
	enthalpy change = $kJ mol^{-1}$ (sign) (value) [3]
	[Total: 10]

2 The identity of a metal, M, can be found by titrating a solution of its carbonate with hydrochloric acid of known concentration.

For Examiner's Use

**FA 3** is a solution of the metal carbonate, M<sub>2</sub>CO<sub>3</sub>, of concentration 6.90 g dm<sup>-3</sup>.

You are to dilute the hydrochloric acid that you used in **Question 1** and then titrate the carbonate solution with this acid.

### (a) Method

#### Dilution of the acid

- Pipette 25.0 cm³ of FA 1 into the 250 cm³ volumetric (graduated) flask labelled FA 4.
- Add distilled water to make the total volume 250 cm<sup>3</sup>.
- Stopper the flask and mix the contents thoroughly.

#### **Titration**

- Fill the burette with diluted hydrochloric acid, FA 4.
- Use a clean pipette to transfer 25.0 cm³ of **FA 3** into a conical flask.
- Titrate **FA 3** with **FA 4** using the indicator provided.
- Perform a rough titration and record your burette readings in the space below.

The rough	titre is		cm <sup>3</sup> .
-----------	----------	--	-------------------

- Carry out as many accurate titrations as you think necessary to obtain consistent results.
- Make certain any recorded results show the precision of your practical work.
- Record, in a suitable form below, all of your burette readings and the volume of FA 4 added in each accurate titration.

I	
II	
III	
IV	
V	
VI	
VII	

[7]

(b)		m your accurate titration results, obtain a suitable value to be used in your calculations.  by clearly how you obtained this value.
		25.0 cm³ of <b>FA 3</b> required cm³ of <b>FA 4</b>
(0)	Cal	[1]
(c)	Cai	culation
	The belo	e equation for the reaction between hydrochloric acid and the metal carbonate is given ow.
		$M_2CO_3 + 2HCl \rightarrow 2MCl + CO_2 + H_2O$
	(i)	Calculate the number of moles of hydrochloric acid present in the volume in <b>(b)</b> .
		moles of HCl = mol
	(ii)	Hence, calculate the number of moles of $\rm M_2CO_3$ present in 25.0 cm $^3$ of <b>FA 3</b> .
		moles of $M_2CO_3 = \dots mol$
	(iii)	Calculate the concentration of M <sub>2</sub> CO <sub>3</sub> in <b>FA 3</b> in mol dm <sup>-3</sup> .
		concentration of M <sub>2</sub> CO <sub>3</sub> = mol dm <sup>-3</sup>
	(iv)	Use your answer to (iii), and the fact that <b>FA 3</b> contains 6.90 g dm <sup>-3</sup> , to determine the relative atomic mass, $A_r$ , of M.
		$A_{r}$ of M =
	(v)	Use your answer to (iv) and the Periodic Table on page 16 to suggest the identity of
		M.
		M is[5]

(d)		concentration of a carbonate solution could be found using either the method in estion 1 or that in Question 2.
	(i)	Suggest, and explain, which of the methods is more accurate.
	(ii)	For the method that you think is less accurate, suggest an improvement to the practical procedure that could be made to improve the accuracy.
		[2]
		[Total: 15]

#### 3 Qualitative Analysis

At each stage of any test you are to record details of the following.

- colour changes seen
- the formation of any precipitate
- the solubility of such precipitates in an excess of the reagent added

Where gases are released they should be identified by a test, **described in the appropriate place in your observations**.

You should indicate clearly at what stage in a test a change occurs.

Marks are **not** given for chemical equations.

No additional tests for ions present should be attempted.

If any solution is warmed, a boiling tube MUST be used.

Rinse and reuse test-tubes and boiling tubes where possible.

Where reagents are selected for use in a test, the name or correct formula of the element or compound must be given.

(a) You are provided with a solid, FA 5. FA 5 is a mixture that contains two anions and two cations.

To all your sample of **FA 5** in a boiling tube add 3 cm depth of distilled water. Shake the tube and filter the contents. Keep the solid residue for tests in **(b)** and the filtered solution for tests in **(c)**.

(a)	(1)	acid, HNO <sub>3</sub> , using a dropping pipette until the solid <b>just</b> disappears. Record your observations and keep the solution for tests in (ii).
		observations
	(ii)	Divide the solution from test (i) equally into three test-tubes.
		To the first test-tube add aqueous sodium hydroxide, NaOH, until in excess. Record your observations.
		ah aaw ati an a
		observations
		Which cations, from those listed in the Qualitative Analysis Notes on page 12, would give these observations?

iii)	You are to devise tests that will positively identify which one of the cations you have suggested in (ii) is present. For each of the possible ions you should indicate the test and the expected result for each test in a suitable table in the space below.
	Use the solutions in the second and third test-tubes to carry out these tests and record your observations in the space below.
	Identify the cation present.  The cation present is

(c)	To 1 cm depth of filtered solution from (a) in a test-tube add 1 cm depth of dilute nitric act followed by a few drops of aqueous silver nitrate. Record your observation.	bid
	observation	
	Which further reagent could be added to this test-tube to help you to confirm the natural of the anion present?	ıre
	reagent	
	Carry out a test using this additional reagent. Record your observation and conclusi about the anion present.	on
	observation	
	The anion present is	[2]
(d)	Using your observation in <b>(b)(i)</b> state which other anion is present in <b>FA 5</b> .	
	The anion present is	[1]

For Examiner's

(e) Solutions FA 6 and FA 7 each contain one of the ions sulfite,  $SO_3^{2-}$ , sulfate,  $SO_4^{2-}$ , nitrite,  $NO_2^{-}$ , or nitrate,  $NO_3^{-}$ .

(i) Carry out the tests in the table below to identify which ion is present in each solution.

11	observ	/ations
test	FA 6	FA 7
To 1 cm depth of solution in a <b>boiling</b> tube, add a small piece of aluminium foil and 1 cm depth of aqueous sodium hydroxide. Warm the mixture with care.		
To 1 cm depth of solution in a test-tube, add a few drops of aqueous barium chloride or barium nitrate, then		
add dilute hydrochloric acid.		
To 1 cm depth of solution in a test-tube, add 1 cm depth of dilute hydrochloric acid.		

	FA 6 contains
	FA 7 contains
(iii)	What type of reaction takes place when a positive observation is seen with aluminium foil and aqueous sodium hydroxide in (i)?
	[5]

(ii) From your observations, identify the anion present in each solution.

[Total: 15]

# **Qualitative Analysis Notes**

Key: [ppt. = precipitate]

# 1 Reactions of aqueous cations

i	reac	tion with
ion	NaOH(aq)	NH <sub>3</sub> (aq)
aluminium, Al³+(aq)	white ppt. soluble in excess	white ppt. insoluble in excess
ammonium, NH <sub>4</sub> +(aq)	no ppt. ammonia produced on heating	_
barium, Ba <sup>2+</sup> (aq)	no ppt. (if reagents are pure)	no ppt.
calcium, Ca <sup>2+</sup> (aq)	white ppt. with high [Ca <sup>2+</sup> (aq)]	no ppt.
chromium(III), Cr³+(aq)	grey-green ppt. soluble in excess giving dark green solution	grey-green ppt. insoluble in excess
copper(II), Cu²+(aq)	pale blue ppt. insoluble in excess	blue ppt. soluble in excess giving dark blue solution
iron(II), Fe <sup>2+</sup> (aq)	green ppt. turning brown on contact with air insoluble in excess	green ppt. turning brown on contact with air insoluble in excess
iron(III), Fe³+(aq)	red-brown ppt. insoluble in excess	red-brown ppt. insoluble in excess
lead(II), Pb <sup>2+</sup> (aq)	white ppt. soluble in excess	white ppt. insoluble in excess
magnesium, Mg²+(aq)	white ppt. insoluble in excess	white ppt. insoluble in excess
manganese(II), Mn <sup>2+</sup> (aq)	off-white ppt. rapidly turning brown on contact with air insoluble in excess	off-white ppt. rapidly turning brown on contact with air insoluble in excess
zinc, Zn²+(aq)	white ppt. soluble in excess	white ppt. soluble in excess

[Lead(II) ions can be distinguished from aluminium ions by the insolubility of lead(II) chloride.]

### 2 Reactions of anions

ion	reaction
carbonate, CO <sub>3</sub> <sup>2-</sup>	CO <sub>2</sub> liberated by dilute acids
chromate(VI), CrO <sub>4</sub> <sup>2-</sup> (aq)	yellow solution turns orange with H <sup>+</sup> (aq); gives yellow ppt. with Ba <sup>2+</sup> (aq); gives bright yellow ppt. with Pb <sup>2+</sup> (aq)
chloride, Cl <sup>-</sup> (aq)	gives white ppt. with Ag <sup>+</sup> (aq) (soluble in NH <sub>3</sub> (aq)); gives white ppt. with Pb <sup>2+</sup> (aq)
bromide, Br <sup>-</sup> (aq)	gives cream ppt. with Ag <sup>+</sup> (aq) (partially soluble in NH <sub>3</sub> (aq)); gives white ppt. with Pb <sup>2+</sup> (aq)
iodide, I <sup>-</sup> (aq)	gives yellow ppt. with Ag <sup>+</sup> (aq) (insoluble in NH <sub>3</sub> (aq)); gives yellow ppt. with Pb <sup>2+</sup> (aq)
nitrate, NO <sub>3</sub> -(aq)	NH <sub>3</sub> liberated on heating with OH <sup>-</sup> (aq) and A <i>l</i> foil
nitrite, NO <sub>2</sub> -(aq)	$NH_3$ liberated on heating with $OH^-(aq)$ and $Al$ foil; NO liberated by dilute acids (colourless $NO \rightarrow$ (pale) brown $NO_2$ in air)
sulfate, SO <sub>4</sub> <sup>2-</sup> (aq)	gives white ppt. with Ba <sup>2+</sup> (aq) or with Pb <sup>2+</sup> (aq) (insoluble in excess dilute strong acids)
sulfite, SO <sub>3</sub> <sup>2-</sup> (aq)	SO <sub>2</sub> liberated with dilute acids; gives white ppt. with Ba <sup>2+</sup> (aq) (soluble in excess dilute strong acids)

## 3 Tests for gases

gas	test and test result
ammonia, NH <sub>3</sub>	turns damp red litmus paper blue
carbon dioxide, CO <sub>2</sub>	gives a white ppt. with limewater (ppt. dissolves with excess CO <sub>2</sub> )
chlorine, Cl <sub>2</sub>	bleaches damp litmus paper
hydrogen, H <sub>2</sub>	"pops" with a lighted splint
oxygen, O <sub>2</sub>	relights a glowing splint
sulfur dioxide, SO <sub>2</sub>	turns acidified aqueous potassium dichromate(VI) from orange to green

## **BLANK PAGE**

## **BLANK PAGE**

The Periodic Table of the Elements

	0	4.0 <b>He</b> Helium	20.2 <b>Ne</b> on 10	39.9 <b>Ar</b> Argon	83.8 <b>Kr</b> Krypton 36	131 <b>Xe</b> Xenon 54	Rn Radon 86	Ununoctium
	=>		19.0 <b>F</b> luorine	35.5 <b>C1</b> Chlorine	79.9 <b>Br</b> Bromine 35	127	At Astatine 85	
	5		16.0 <b>O</b> Oxygen	32.1 <b>S</b> Sulfur	79.0 Se Selenium	128 <b>Te</b> Tellurium 52	<b>Po</b> Polonium 84	<b>Uuh</b> Unuhexium 116
	>		14.0 <b>N</b> Nitrogen 7	31.0 Phosphorus	74.9 <b>AS</b> Arsenic 33	122 <b>Sb</b> Antimony 51	209 <b>Bismuth</b> 83	
	≥		12.0 <b>C</b> Carbon 6	28.1 <b>Si</b> Silicon	72.6 <b>Ge</b> Germanium 32	<b>Sn</b> 50	207 <b>Pb</b> Lead 82	Ununquadium
	≡		10.8 Boron 5	27.0 <b>A 1</b> Aluminium 13	69.7 <b>Ga</b> Gallium 31	115   n   n   1   1   1   1   1   1   1   1	204 <b>T 1</b> Thallium	
					65.4 <b>Zn</b> Zinc 30	Cd Cadmium 48	Hg Mercury	Uub Ununbium 112
					63.5 <b>Cu</b> Copper 29	108 <b>Ag</b> Silver 47	197 <b>Au</b> Gold	Ununuium 111
Group					58.7 <b>Nickel</b> Nickel	106 <b>Pd</b> Palladium 46	195 <b>Pt</b> Platinum 78	Uun Ununnilium 110
Gr			1		58.9 <b>Co</b> Cobalt 27	103 <b>Rh</b> Rhodium 45	192	Meitnerium 109
		1.0 Hydrogen			55.8 <b>Fe</b> Iron	Ruthenium 44	190 <b>Os</b> Osmium 76	Hassium 108
					Manganese 25	Tc Technetium 43	186 <b>Re</b> Rhenium 75	Bh Bohrium 107
					52.0 <b>Cr</b> Chromium 24	95.9 Molybdenum 42	184 <b>W</b> Tungsten 74	Seaborgium
					50.9 <b>V</b> Vanadium	92.9 <b>Nb</b> Niobium 41	181 <b>Ta</b> Tananam 73	Db Dubnium 105
					47.9 <b>Ti</b> Titanium	91.2 <b>Zr</b> Zirconium 40	178 <b>#</b> Hafnium 72	Rutherfordium
					<b>Sc</b> Scandium 21	88.9 <b>×</b>	139 <b>La</b> Lanthanum 57 *	Actinium teges
	=		9.0 <b>Be</b> Beryllium	24.3 <b>Mg</b> Magnesium	40.1 <b>Ca</b> Calcium 20	87.6 <b>St</b> rontium	137 <b>Ba</b> Barium 56	Radium 88
	_		6.9 <b>L.i</b> Lithium	23.0 <b>Na</b> Sodium	39.1 <b>K</b> Potassium	85.5 <b>Rb</b> Rubidium	133 <b>Cs</b> Caesium 55	Francium 87

*58-71 190-10	58-71 Lanthanides 90-103 Actinides	* ides tes	140 Cerium	141 <b>Pr</b> Praseodymium 59	Neodymium	Pm Promethium	Samarium	152 <b>Eu</b> Europium	157 <b>Gd</b> Gadolinium 64	159 <b>Tb</b> Terbium	163 Dy Dysprosium	165 <b>Ho</b> Holmium	167 <b>Er</b> Erbium	169 <b>Tm</b> Thulium	Yb Ytterbium	175 <b>Lu</b> Lutetium
	æ	a = relative atomic mass +	8	3	3	5		3		3	8		3		2	
Key	×	X = atomic symbol	두	Ра	>	dN	Pu		Cm	BK	ర	Es	Fm	Md	No	ئ
	۵	b = proton (atomic) number	Thorium 90	Protactinium 91	Uranium 92	Neptunium 93	Plutonium 94	Americium 95	Curium 96	Berkelium 97	Californium 98	٠, ١	Fermium 100	Mendelevium 101	Nobelium 102	Lawrendum 103

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.