

Candidate Name _____

Centre Number

Candidate

Number

--	--

CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Advanced Level

CHEMISTRY

PAPER 5 Practical Test

9701/5

OCTOBER/NOVEMBER SESSION 2002

1 hour 30 minutes

Candidates answer on the question paper.

Additional materials:

As listed in Instructions to Supervisors

TIME 1 hour 30 minutes

INSTRUCTIONS TO CANDIDATES

Write your name, Centre number and candidate number in the spaces at the top of this page.

Answer **all** questions.

Write your answers in the spaces provided on the question paper.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets [] at the end of each question or part question.

You are advised to show all working in calculations.

Use of a Data Booklet is unnecessary.

FOR EXAMINER'S USE

1	
2	
TOTAL	

This question paper consists of 5 printed pages and 3 blank pages.



- 1 You are to investigate the reaction between substance **X**, iodine and hydrogen ions.

FA 1 is 1.00 mol dm⁻³ sulphuric acid.

FA 2 is an aqueous solution of substance **X**.

FA 3 is 0.0038 mol dm⁻³ iodine, I₂.

Fill a burette with solution **FA 3**.

- (a) Using the measuring cylinder provided, measure out 20.0 cm³ of **FA 1** and 20.0 cm³ of **FA 2**, as shown in column 1 of *Table 1.1*, into a 250 cm³ conical flask. It is not necessary to rinse the measuring cylinder between solutions.

Measure out 4.0 cm³ of **FA 3** from the burette into a test-tube.

Start the reaction by tipping the **FA 3** from the test-tube into the conical flask. Start the stop-clock with one hand and swirl the contents of the flask with the other. Place the flask on a white tile and stop the clock as soon as the colour disappears.

Record the time (in seconds, to the nearest second) in *Table 1.1*.

Repeat the experiment using the different volumes of **FA 1**, **FA 2** and **FA 3** as shown in *Table 1.1*. Where water is required, use the measuring cylinder to add the water to the other solutions in the conical flask.

Experiment 2 is the same as experiment 1 to give you the opportunity of practising the technique.

The 'rate of reaction' can be calculated by using the relationship:

$$\text{'rate'} = \frac{\text{volume of FA 3 in cm}^3}{\text{time in seconds for colour to disappear}}$$

Table 1.1

	1	2	3	4
volume of FA 1 / cm ³	20.0	20.0	10.0	20.0
volume of FA 2 / cm ³	20.0	20.0	20.0	10.0
volume of water / cm ³	0.0	0.0	10.0	10.0
volume of FA 3 / cm ³	4.0	4.0	4.0	4.0
time for colour to disappear / s				
'rate' of reaction				

Calculate each 'rate' and complete *Table 1.1*.

[10]

As the total volume of liquid is the same in each experiment, the volume of any reagent can be used as a measure of its concentration.

(b) Compare experiments 2 and 3.

(i) Which reagents have the same concentration in both experiments?

[1]

(ii) Which reagent has a different concentration?

[1]

(iii) How is the rate of reaction affected by the change of concentration of the reagent named in (ii)?

[3]

(c) Compare experiments 2 and 4.

(i) Which reagents have the same concentration in both experiments?

[1]

(ii) Which reagent has a different concentration?

[1]

(iii) How is the rate of reaction affected by the change of concentration of the reagent named in (ii)?

[3]

(d) A text-book states that the reaction is zero order with respect to iodine. What volumes of reagents, compared with experiment 2, would you mix to investigate this statement?

FA 1 cm³ **FA 2** cm³

water cm³ **FA 3** cm³

[1]

[Total : 21]

