

**MARK SCHEME for the October/November 2006 question paper**

**9701 CHEMISTRY**

**9701/03**

Paper 3 (Practical Test), maximum raw mark 25

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

The grade thresholds for various grades are published in the report on the examination for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2006 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



<b>Page 1</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – Oct/Nov 2006</b>	<b>9701</b>	<b>3</b>

**Question 1**

**(b) Candidate's results**

Check (and correct if necessary) all subtractions in Tables 1.1 and 1.2. The subtraction of titration results labelled as rough need not be checked unless the candidate has included them in the volumes used to calculate the average.

Tick (if accepting the candidate's value) or correct this value, recording it by titration Table 1.2 on the script.

Calculate for each candidate:

$$\text{Candidate's Titre} \times \frac{\text{Candidate's volume of FA 1 diluted}}{\text{Supervisor's volume of FA 1 diluted}}$$

Record this titre, **correct to two decimal places**, against Table 1.2 and calculate the difference between the calculated value and the Supervisor's titre.

Award accuracy marks on the difference as shown on the next page.

<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – Oct/Nov 2006</b>	<b>9701</b>	<b>3</b>

Award **accuracy marks** for differences as follows:

<b>Accuracy mark</b>		<b>Spread Penalty</b>	
Mark	Difference to Supervisor / cm <sup>3</sup>	Range used / cm <sup>3</sup>	Deduction
8	Up to 0.20	0.20+ to 0.25	1
7	0.20+ to 0.25	0.25 to 0.30	2
6	0.25+ to 0.30	0.30+ to 0.35	3
5	0.30+ to 0.40	0.35+ to 0.40	4
4	0.40+ to 0.60	0.40+ to 0.50	5
3	0.60+ to 0.80	0.50+ to 0.60	6
2	0.80+ to 1.00	0.60+ to 0.80	7
1	1.00+ to 1.50	Greater than 0.80	8
0	Greater than 1.50		

[8]

**Deduct** from the accuracy mark **one mark** for **each** of the following errors:

- (i) Volume diluted, as recorded by the candidate in the final line of Table 1.1 is outside of the range 34.00 cm<sup>3</sup> to 35.00 cm<sup>3</sup>.  
Ignore any subtraction error that would take the volume out of range
- (ii) Any Initial and Final Burette reading transposed or 50 used as initial burette reading in either table **or**  
Final burette readings in either table (except for any titration recorded as Rough) not recorded to 2 decimal places, **or**  
"impossible" burette readings (e.g. 23.47 cm<sup>3</sup>) recorded at any point in either of the tables.  
All final burette readings = 50 cm<sup>3</sup>.  
(When assessing titration volumes to use you will need to consider this case carefully. Is there a general inversion of initial and final burette readings with 50 used instead of 0 or is this a genuine claim that all titres finished on 50 cm<sup>3</sup>. The volume run out will be the best evidence to use)
- (iii) No two recorded (uncorrected) titres within 0.1 cm<sup>3</sup> in Table 1.2.
- (iv) An incorrect average calculated **or**  
No selection of at least two titres for the calculation of the average shown (selected titres may be ticked or used in a calculation of the average.) **or**  
Error in subtraction in any accurate titre (any rough titre if included in selection of titres to calculate the average) or error in subtraction in Table 1.1

**THERE IS A MAXIMUM DEDUCTION OF TWO MARKS FROM THE ACCURACY MARKS**

Page 3	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – Oct/Nov 2006	9701	3

If 34.5 cm<sup>3</sup> of FA 1 are diluted, the expected titre is around 25 cm<sup>3</sup>.  
Evaluations for these values are given, in boxes, as a guide to expected values

- (c) Give **one mark** for calculating the correct concentration in mol dm<sup>-3</sup>  $\frac{4.93}{214.1}$  or **0.023(0)**

Do not give this mark if K = 39.0 has been used.

Give **one mark** for  $\frac{25}{1000}$  x concentration in mol dm<sup>-3</sup> calculated by candidate

or

Give **one mark** for  $\frac{4.93}{1000}$  x 25 and **one mark** for this expression x  $\frac{1}{214.1}$   
(one) (one)

The answer 5.75(7) x 10<sup>-4</sup> with no working gains both marks

5.75(7) x 10<sup>-4</sup>

[2]

- (d) Give **one mark** for ans (c) x 6 or 3.45(4) x 10<sup>-3</sup>

3.45(4) x 10<sup>-3</sup>

[1]

- (e) Give **one mark** for ans (d) x  $\frac{1000}{\text{Candidate's Titre}}$

0.138

[1]

- (f) Give **two marks** for ans (e) x  $\frac{250}{\text{Candidate's volume of FA 1 diluted}}$  x 36.5

one

one

Give **one** of these two marks for any answer where only (e) x 36.5 is seen.

and

1.001

36.54

Give **one mark** for a fully correct answer within 1% of the value calculated by the examiner. Do not give this mark if there are 'chemical' errors in (c) to (f) even if these errors are self-cancelling. The mark may be given if 214 rather than 214.1 has been used for M<sub>r</sub> of KIO<sub>3</sub>, providing the final answer is within 1% of the examiner calculated value

The correct answer is:  $\frac{31517.69}{\text{Candidate's Titre} \times \text{Candidate's volume of FA 1 diluted}}$

[3]

[Total for Qn 1 15]

Page 4	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – Oct/Nov 2006	9701	3

- 2 **FA 6** is a mixture of soluble **FA 7** (ammonium chloride) and insoluble **FA 8** (manganese(II) carbonate)

Some centres were unable to obtain manganese(II) carbonate and have been instructed to use magnesium zinc or copper carbonate. Magnesium carbonate was the recommended choice.

RETROSPECTIVE OBSERVATION MARKS MAY BE EARNED FROM THE CONCLUSIONS WHERE INCOMPLETE INFORMATION RECORDED, IN THE OBSERVATION BOX, AGAINST THE TEST IS COMPLETED.

**DO NOT GIVE RETROSPECTIVE MARKS FOR CONTRARY INFORMATION**

**Tests on the Filtrate FA 7**

Test	Observations		
(b) To 1 cm depth of the filtrate from (a) in a boiling-tube, add 2 cm depth of aqueous sodium hydroxide.	Give <b>one mark</b> for  No ppt. / colourless solution / no reaction (allow slightly brown – as $MnCO_3$ might not be fresh) <b>No yellow or orange colours</b>	1	
	Give <b>one mark</b> for  <b>Gas</b> evolved – <b>turns</b> red litmus blue (or other appropriate <b>test</b> for ammonia <b>gas</b> )  <i>Ignore any precipitate in this section</i>	1	
(c) To 1 cm depth of the filtrate from (a) in a test-tube, add 1 cm depth of aqueous lead nitrate, then	Give <b>one mark</b> for  White precipitate <b>which is</b>		
	heat the mixture to boiling point, then	soluble/partially soluble in hot water	1
	cool the tube by standing in a beaker of cold water	Give <b>one mark</b> if the (white) <u>precipitate reforms</u> or <b>crystals</b> form	1
(d) To 1 cm depth of the filtrate from (a) in a test-tube, add 1 cm depth of aqueous silver nitrate.	Give <b>one mark</b> for  White precipitate .....		
	Then add 5 cm depth of dilute aqueous ammonia.	Soluble in aqueous ammonia	1

**TO GAIN ANY DEDUCTION MARK THERE MUST BE REFERENCE TO THE REAGENTS. REFERENCE TO THE TEST ALONE IS SUFFICIENT PROVIDING THE OBSERVATION MARK HAS BEEN GIVEN FOR THAT TEST.**

Give **one mark** if the cation in **FA 7** is identified as  $NH_4^+$  or ammonium **and**

there is reference to ammonia evolved / alkaline gas / gas turning red litmus blue (on heating with sodium hydroxide.) **There must be a clear or unambiguous link to test (b) part (ii).**

[1]

Give **one mark** if the anion is identified as  $Cl^-$  **and**

There is reference to white precipitate with  $Pb^{2+}$  or white ppt, soluble in ammonia with  $Ag^+$ .

[1]

Page 5	Mark Scheme	Syllabus	Paper
	GCE A LEVEL – Oct/Nov 2006	9701	3

### Tests on the residue FA 8

Test	Observations [3]	
(e) Transfer the residue to a boiling-tube and add 4 cm depth of dilute hydrochloric acid. Retain the solution for tests (f) and (g).	Give <b>one mark</b> for Gas evolved turns lime-water milky / cloudy / chalky or gives a white ppt with Ca(OH) <sub>2</sub>  Details of the test may be given in the conclusion	1
(f) To 1 cm depth of the solution from (e) in a test-tube add, a little at a time, 5 cm depth of aqueous sodium hydroxide.	Give <b>one mark</b> for Off-white / buff / pale-brown ppt. Insoluble in excess (NaOH)  [For Mg <sup>2+</sup> - white ppt insol in excess For Zn <sup>2+</sup> - white ppt sol in excess For Cu <sup>2+</sup> - blue ppt insol in excess]	1
(g) To 1 cm depth of the solution from (e) in a test-tube add, a little at a time, 5 cm depth of dilute aqueous ammonia.	Give <b>one mark</b> for Off-white / buff / pale-brown ppt. Insoluble in excess (NH <sub>3</sub> (aq))  [For Mg <sup>2+</sup> - white ppt insol in excess For Zn <sup>2+</sup> - white ppt sol in excess For Cu <sup>2+</sup> - <u>pale</u> blue ppt sol in excess <b>or</b> blue ppt sol in excess to form a deep blue soln]	1

Give **one mark** if the cation in **FA 8** is identified as Mn<sup>2+</sup> or manganese(II) [Mg<sup>2+</sup>, Zn<sup>2+</sup>, Cu<sup>2+</sup>] **and**

There is reference to the colour of the precipitate and insolubility of precipitate in excess of sodium hydroxide and aqueous ammonia **or**

reference to an **off-white precipitate** in each test. [1]

**Allow deductions of other cations, given from incorrect observations, providing all observations – including the solubility of any precipitate in excess NaOH and ammonia for the ion given.**

**e.g. deduction of Fe<sup>3+</sup> from brown or red-brown precipitate, insoluble in excess aqueous NaOH and ammonia.**

Give **one mark** if the anion is identified as CO<sub>3</sub><sup>2-</sup> or carbonate **and**

there is reference to effervescence with hydrochloric acid **or** [1]

carbon dioxide is named (without a confirmatory test) as the gas given off **and** effervescence is an observation in (e) **or**

there is reference back to a positive test for carbon dioxide in (e).

There are 12 scoring points. If a candidate scores in excess of 10 points record the raw score, crossed through and replaced by 10 max.

[Total for Qn 2 10]

[Total for Paper 25]