UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced/Advanced Subsidiary Level

MARK SCHEME for the May/June 2006 question paper

9709 MATHEMATICS

9709/02

Paper 2

Maximum raw mark 50

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the Report on the Examination.

The minimum marks in these components needed for various grades were previously published with these mark schemes, but are now instead included in the Report on the Examination for this session.

• CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the May/June 2006 question papers for most IGCSE and GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.


Mark Scheme Notes

Marks are of the following three types:

- M Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.
- A Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).
- B Mark for a correct result or statement independent of method marks.
- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol √ implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.
 B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking g equal to 9.8 or 9.81 instead of 10.



The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Anv	Equivalent	Form (c	of answer is	egually	/ acceptable)

- AG Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
- BOD Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
- CAO Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
- CWO Correct Working Only often written by a 'fortuitous' answer
- ISW Ignore Subsequent Working
- MR Misread
- PA Premature Approximation (resulting in basically correct work that is insufficiently accurate)
- SOS See Other Solution (the candidate makes a better attempt at the same question)
- SR Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

Penalties

- MR -1 A penalty of MR -1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through √" marks. MR is not applied when the candidate misreads his own figures this is regarded as an error in accuracy. An MR-2 penalty may be applied in particular cases if agreed at the coordination meeting.
- PA -1 This is deducted from A or B marks in the case of premature approximation. The PA -1 penalty is usually discussed at the meeting.

	_	GCE A/AS LEVEL – May/June 2006 9709		02	
	ETTHER	State or imply non-modular inequality $(2x-7)^2 > 3^2$, or corresponding equation			
		Obtain critical values 2 and 5	AL		
0	CON.	State correct answer $x \le 2$, $x \ge 5$. State one critical value, e.g. $x = 5$, by solving a linear equation (or inequality) or from a	AI		
	OR.	graphical method ar by inspection	B1		
		State the other critical value correctly	B1		
		State correct answer $x < 2$, $x > 5$	В1	3	
2	(i) Use trig formulae to express LHS in terms of cos x and sin x		MI		
		errect exact values of cos 60°, sin 60°, etc	MI.		
70		n given answer	A1	3	
	(ii) State	or imply answer is $\cos^{-1}(1/\sqrt{3})$	M		
	Obtai	n answer 54 7°	.Al	1	
3	State con	rect derivative 1 – 2sin x	В		
	Equate de	grivative to zero and solve for x	MJ		
	Obtain ar	iswer x 1/2 m	All		
	Carry out	an appropriate method for determining the nature of a stationary point	MI		
	Show that $x = \frac{1}{6}\pi$ is a maximum with no errors seen				
	Obtain second answer $x = \frac{\pi}{6} \pm in \text{ range}$				
	Show the	Show this is a minimum point $(f.r.$ is on the incorrect derivative $1 + 2\sin x \cdot 1$			
	(i) Subs	titule $x = 1$ or $x = -2$ and equate to zero	MI		
7		in a correct equation, e.g. $a = b - 5 = 0$	Al		
	Obta	an a second correct equation, e.g. $-8a + 4b + 4 = 0$	AL		
		e a relevant pair of equations for n or for h	MI AI	- 5	
		$\sin a = 2$ and $b = 3$ stitute for a and h and either divide by $(x-1)(x+2)$ or attempt third factor by inspection	MI	-	
		ain answer $2r + 1$	Al	2	
5	(i) Shite	$2y\frac{dy}{dx}$ as the derivative of y^2	Bi		
	State	$2v + 2x \frac{dy}{dx}$ or equivalent, as derivative of $2xy$	BI		
	Equal	te attempted derivative of LHS to zero and set dv equal to zero	ML		
	Obtair	n given relation y = -3x correctly M1 is dependent on at least one B1 being carned cariter.	AI	4	
	(II) Carry	out complete method for finding x^2 or μ^2	MI		
		$n e^2 = 1 \text{ or } v^2 = 9.$	Al		
		n point (1, -3)	A1		
		n second point (-1, 3)	AD	- 4	

Mark Scheme

Syllabus

Paper

Page 1

	Page	2 Mark Scheme	Syllabus	Paper	
		GCE A/AS LEVEL – May/June 2006	9709	02	
	(i) N	Make recognizable sketch of an appropriate exponential curve, e.g.)	$= 9e^{-2x}$	BI	
		Sketch the appropriate second curve, e.g. $y = x$ correctly and justify t		131	
		Consider sign of $x - 9e^{-2x}$ at $x = 1$ and $x = 2$, or equivalent Complete the argument correctly with appropriate calculations		M1 A1	ģ
	(iii) S	State or imply the exputation $r = \frac{1}{2} (\ln 9 - \ln x)$		B1	
	(iv) L	Rearrange this in the form given in part (i), or work vice versal fee the iterative formula correctly at feast once Obtain final answer $x = 1.07$	and a state of family facility	MI Al	R
		Show sufficient iterations to justify its accuracy to 2 ft.p. or show the interval (1.065, 1.075)	ac is a sign change or the	Al	1
		Obtain derivative of the form $\frac{k}{2x+3}$, where $k=2$ or $k=1$		MI AI	
		Obtain correct derivative $\frac{2}{2x+3}$		141	
	 (ii) State Indefinite integral of the form m In(2x +3) Use limits correctly. Obtain given answer 			M1/de	p*
	(ili)	Carry out division method reaching a linear quotient and constant. Obtain quotient 2x + 1 Obtain remainder -3	remainder	MI AI	
		Attempt integration of an integrand of the form $ax + b + \frac{c}{2x + 3}$		MI	
		Obtain indefinite integral $\kappa^2 + \kappa - \frac{3}{2} \ln(2\kappa + 3)$		AIA	
	- 3	Substitute limits and obtain given answer [The f.t. mark is also available if the indefinite integral of the third integral is stated to be c in 3.]	term is omitted but its definite	AL	