

**MARK SCHEME for the October/November 2010 question paper  
for the guidance of teachers**

**9709 MATHEMATICS**

**9709/62**

Paper 6, maximum raw mark 50

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2010 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2010	9709	62

## Mark Scheme Notes

Marks are of the following three types:

**M** Method mark, awarded for a valid method applied to the problem. Method marks are not lost for numerical errors, algebraic slips or errors in units. However, it is not usually sufficient for a candidate just to indicate an intention of using some method or just to quote a formula; the formula or idea must be applied to the specific problem in hand, e.g. by substituting the relevant quantities into the formula. Correct application of a formula without the formula being quoted obviously earns the M mark and in some cases an M mark can be implied from a correct answer.

**A** Accuracy mark, awarded for a correct answer or intermediate step correctly obtained. Accuracy marks cannot be given unless the associated method mark is earned (or implied).

**B** Mark for a correct result or statement independent of method marks.

- When a part of a question has two or more "method" steps, the M marks are generally independent unless the scheme specifically says otherwise; and similarly when there are several B marks allocated. The notation DM or DB (or dep\*) is used to indicate that a particular M or B mark is dependent on an earlier M or B (asterisked) mark in the scheme. When two or more steps are run together by the candidate, the earlier marks are implied and full credit is given.
- The symbol  $\surd$  implies that the A or B mark indicated is allowed for work correctly following on from previously incorrect results. Otherwise, A or B marks are given for correct work only. A and B marks are not given for fortuitously "correct" answers or results obtained from incorrect working.
- Note: B2 or A2 means that the candidate can earn 2 or 0.  
B2/1/0 means that the candidate can earn anything from 0 to 2.

The marks indicated in the scheme may not be subdivided. If there is genuine doubt whether a candidate has earned a mark, allow the candidate the benefit of the doubt. Unless otherwise indicated, marks once gained cannot subsequently be lost, e.g. wrong working following a correct form of answer is ignored.

- Wrong or missing units in an answer should not lead to the loss of a mark unless the scheme specifically indicates otherwise.
- For a numerical answer, allow the A or B mark if a value is obtained which is correct to 3 s.f., or which would be correct to 3 s.f. if rounded (1 d.p. in the case of an angle). As stated above, an A or B mark is not given if a correct numerical answer arises fortuitously from incorrect working. For Mechanics questions, allow A or B marks for correct answers which arise from taking  $g$  equal to 9.8 or 9.81 instead of 10.

<b>Page 3</b>	<b>Mark Scheme: Teachers' version</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>GCE A LEVEL – October/November 2010</b>	<b>9709</b>	<b>62</b>

The following abbreviations may be used in a mark scheme or used on the scripts:

AEF	Any Equivalent Form (of answer is equally acceptable)
AG	Answer Given on the question paper (so extra checking is needed to ensure that the detailed working leading to the result is valid)
BOD	Benefit of Doubt (allowed when the validity of a solution may not be absolutely clear)
CAO	Correct Answer Only (emphasising that no "follow through" from a previous error is allowed)
CWO	Correct Working Only – often written by a 'fortuitous' answer
ISW	Ignore Subsequent Working
MR	Misread
PA	Premature Approximation (resulting in basically correct work that is insufficiently accurate)
SOS	See Other Solution (the candidate makes a better attempt at the same question)
SR	Special Ruling (detailing the mark to be given for a specific wrong solution, or a case where some standard marking practice is to be varied in the light of a particular circumstance)

### **Penalties**

MR –1	A penalty of MR –1 is deducted from A or B marks when the data of a question or part question are genuinely misread and the object and difficulty of the question remain unaltered. In this case all A and B marks then become "follow through $\sqrt{}$ " marks. MR is not applied when the candidate misreads his own figures – this is regarded as an error in accuracy. An MR–2 penalty may be applied in particular cases if agreed at the coordination meeting.
PA –1	This is deducted from A or B marks in the case of premature approximation. The PA –1 penalty is usually discussed at the meeting.

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2010	9709	62

<p><b>1</b> <math>4p + 5p^2 + 1.5p + 2.5p + 1.5p = 1</math>  <math>10p^2 + 19p - 2 = 0</math></p> <p><math>p = 0.1</math> or <math>-2</math></p> <p><math>p = 0.1</math></p>	<p>M1</p> <p>A1</p> <p>A1</p> <p>[3]</p>	<p>Summing 5 probs to = 1 can be implied</p> <p>For 0.1 seen with or without <math>-2</math></p> <p>Choosing 0.1 must be by rejecting <math>-2</math></p>
<p><b>2 (i)</b> <math>\Sigma(x - 50) = 824 - 16 \times 50 = 24</math></p> $\frac{\Sigma(x - 50)^2}{16} - \left(\frac{\Sigma(x - 50)}{16}\right)^2 = 6.5^2$ <p><math>\Sigma(x - 50)^2 = 712</math></p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>Correct answer</p> <p>Consistent substituting in the correct coded variance formula OR valid method for <math>\Sigma x^2</math> then expanding <math>\Sigma(x - 50)^2</math>, 3 terms at least 2 correct</p> <p>Correct answer</p>
<p><b>(ii)</b> new mean = <math>896/17 (= 52.7)</math></p> $\text{new var} = \frac{712 + 22^2}{17} - \left(\frac{24 + (72 - 50)}{17}\right)^2$ <p>new sd = 7.94</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>Correct answer</p> <p>Using the correct coded variance formula with <math>n = 17</math> and new coded mean<sup>2</sup> OR their <math>(\Sigma x^2 + 72^2)/17 - \text{their new mean}^2</math></p> <p>Rounding to correct answer, accept 7.95 or 7.98 or 7.91</p>

<p><b>3</b> <math>P(E \text{ and } 12) = \frac{2}{5} \times \frac{4}{36} = \frac{8}{180} (2/45)</math></p> <p><math>P(12) = \frac{3}{5} \times \frac{1}{36} + \frac{8}{180} = \frac{11}{180} (0.0611)</math></p> <p><math>P(E   12) = \frac{P(E \text{ and } 12)}{P(12)}</math></p> <p><math>= \frac{8}{11} (0.727)</math></p> <p>OR list  Even: 2 and (4,3) or (3,4) or (2,6) or (6,2)  4 and ditto  Gives 8 options</p> <p>Odd: 1 and (6,6) or 3 and (6,6) or 5 and (6,6)  Gives 3 options</p> <p><math>\text{Prob}(E   12) = 8/11</math></p>	<p>M1</p> <p>A1</p> <p>M1</p> <p>A1ft</p> <p>M1dep</p> <p>A1</p> <p>[6]</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p> <p>M1</p> <p>A1</p>	<p>2/5 or 3/5 mult by dice-related probability seen anywhere</p> <p><math>\frac{2}{5} \times \frac{4}{36}</math> seen oe</p> <p>Summing two 2-factor probs involving 2/5 and 3/5  3/5 <math>\times</math> 1/36 + their P(E and 12), ft their P(E 12)</p> <p>Subst in condit prob formula, must have a fraction</p> <p>Correct answer</p> <p>List attempt evens</p> <p>8 options</p> <p>List attempt odds</p> <p>3 options</p> <p>(Their even)/(their total)</p> <p>Correct answer</p>																														
<p><b>4 (i)</b></p> <table border="1" style="margin-left: 40px;"> <thead> <tr> <th style="border-right: 1px solid black;">sugar</th> <th></th> <th>flour</th> </tr> </thead> <tbody> <tr><td></td><td>194</td><td>1 5 9</td></tr> <tr><td></td><td>195</td><td>3</td></tr> <tr><td>8 1</td><td>196</td><td>2 4</td></tr> <tr><td>7</td><td>197</td><td>7</td></tr> <tr><td>9 4 3</td><td>198</td><td></td></tr> <tr><td>4</td><td>199</td><td></td></tr> <tr><td>8</td><td>200</td><td></td></tr> <tr><td>7 4 1</td><td>201</td><td></td></tr> </tbody> </table> <p>key</p> <table border="1" style="margin-left: 40px;"> <tr> <td style="border-right: 1px solid black;">1</td> <td>196</td> <td style="border-left: 1px solid black;">2</td> </tr> </table> <p>means 1.961 kg for sugar and 1.962 kg for flour</p>	sugar		flour		194	1 5 9		195	3	8 1	196	2 4	7	197	7	9 4 3	198		4	199		8	200		7 4 1	201		1	196	2	<p>B1</p> <p>B1</p> <p>B1</p> <p>B1ft</p> <p>[4]</p>	<p>Correct stem must be integers. (stem and leaves can be in reverse order)</p> <p>Correct leaves flour must be single and ordered</p> <p>Correct leaves sugar must be single and ordered</p> <p>Correct key needs all this, ft if single leaves and 1.96 etc in stem</p>
sugar		flour																														
	194	1 5 9																														
	195	3																														
8 1	196	2 4																														
7	197	7																														
9 4 3	198																															
4	199																															
8	200																															
7 4 1	201																															
1	196	2																														
<p><b>(ii)</b> med = 1.989 kg</p> <p>IQ range = 2.011 – 1.977</p> <p>= 0.034 kg</p>	<p>B1</p> <p>M1</p> <p>A1</p> <p>[3]</p>	<p>correct median</p> <p>subt their LQ from their UQ, UQ &gt; med, LQ &lt; med</p> <p>Correct answer</p>																														

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2010	9709	62

<p><b>5 (i)</b> Zotoc: <math>z = \frac{367 - 320}{21.6} = 2.176</math>  Ganmor: <math>z = \frac{367 - 350}{7.5} = 2.267</math></p> <p><math>P(\text{Zotoc}) = 0.985</math>  <math>P(\text{Ganmor}) = 0.988</math></p>	<p>M1  A1 A1 [3]</p>	<p>Standardising either car's fuel, no cc, no sq, no <math>\sqrt{\quad}</math></p> <p>Correct answer  Correct answer</p>
<p><b>(ii)</b> <math>z = 0.23</math>  <math>0.23 = \frac{x - 320}{21.6}</math></p> <p><math>x = 324.968</math>  <math>d = 4.97</math></p>	<p>B1 M1 M1ind A1 [4]</p>	<p><math>\pm 0.23</math> seen</p> <p>Standardising either car, no cc, no sq rt, no sq</p> <p><math>320 + d - 320</math> i.e. just <math>d</math> on num</p> <p>Correct answer, <math>-4.97</math> gets A0</p>
<p><b>6 (i)</b> constant/given prob, independent trials, fixed/given no. of trials, only two outcomes</p>	<p>B1 B1 [2]</p>	<p>One option correct  Three options correct</p>
<p><b>(ii)</b> <math>P(8, 9, 0, 1) =</math>  <math>{}^9C_8(0.3)^8(0.7) + (0.3)^9 + (0.7)^9 + {}^9C_1(0.3)(0.7)^8</math>  <math>= 0.196</math></p>	<p>M1 A1 A1 [3]</p>	<p>One term seen involving <math>(0.3)^x(0.7)^{9-x}({}^9C_x)</math></p> <p>Correct unsimplified expression  Correct answer</p>
<p><b>(iii)</b> mean = <math>90 \times 0.3 = 27</math>  var = 18.9  <math>P(X &gt; 35) = 1 - \Phi\left(\frac{35.5 - 27}{\sqrt{18.9}}\right)</math>  <math>= 1 - \Phi(1.955) = 0.0253</math>  <math>P(X &lt; 27) = \Phi\left(\frac{26.5 - 27}{\sqrt{18.9}}\right) = 1 - \Phi(0.115)</math>  <math>= 0.4542</math>  Total prob = 0.480 accept 0.48</p>	<p>B1 M1 M1 M1 A1 [5]</p>	<p>Expressions for 27 and 18.9 (4.347) seen</p> <p>Standardising one expression, must have sq rt in denom, cc not necessary</p> <p>Continuity correction applied at least once  <math>(1 - \Phi_1) + (1 - \Phi_2)</math> accept <math>(0.0329 + 0.5)</math> if no cc</p> <p>Rounding to correct answer</p>

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE A LEVEL – October/November 2010	9709	62

<p>7 (i) 4M 2W or 5M 1W</p> <p>chosen in <math>{}^{10}C_4 \times {}^9C_2 + {}^{10}C_5 \times {}^9C_1</math>  <math>= 9828</math></p>	<p>M1 A1 A1 [3]</p>	<p>At least 1 of <math>{}^{10}C_4 \times {}^9C_2</math> and <math>{}^{10}C_5 \times {}^9C_1</math> seen  Correct unsimplified expression  Correct answer</p>
<p>(ii) <math>{}^9C_3 \times {}^8C_1 + {}^9C_4 = 798</math></p> <p>Prob = <math>798/9828 = 0.0812</math></p>	<p>M1 A1 [2]</p>	<p>One of <math>{}^9C_3 \times {}^8C_1</math> and <math>{}^9C_4 \times ({}^8C_0)</math> seen  Correct answer</p>
<p>(iii) Albert + not T... <math>{}^9C_3 \times {}^8C_2 + {}^9C_4 \times {}^8C_1</math>  <math>= 3360</math>  Tracey + not A... <math>{}^9C_4 \times {}^8C_1 + {}^9C_5</math>  <math>= 1134</math></p> <p>Number of ways = 4494</p>	<p>M1 A1 A1 [3]</p>	<p>One of <math>{}^9C_3 \times {}^8C_2</math> or <math>{}^9C_4 \times {}^8C_1</math> or <math>{}^9C_5 \times ({}^8C_0)</math> seen  Unsimplified 3360 or 1134 seen  Correct final answer</p>
<p>(iv) <math>6! - 4! \times 5 \times 2</math> or <math>6! - 5! \times 2 (= 480)</math>  OR <math>4! \times 5 \times 4</math> or <math>4! \times {}^5P_2 (= 480)</math></p> <p>prob = <math>480/6! = 2/3 (0.667)</math></p> <p>OR using probabilities...as above</p> <p>OR Women together <math>5!/4! (= 5)</math>  Women not together = <math>15 - 5 = 10</math>  total ways MMMMW = <math>6!/4!2! = 15</math>  prob = <math>2/3</math></p>	<p>B1 M1 A1 [3]</p> <p>B1 M1 A1</p>	<p><math>6! - 4! \times 5 \times 2</math> or <math>6! - 5! \times 2</math>  or <math>4! \times 5 \times 4</math> or <math>4! \times {}^5P_2</math>  dividing by 6!  correct answer</p> <p>5 or 10 seen  Dividing by 15  Correct answer</p>