UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

GCE Advanced Subsidiary Level and GCE Advanced Level

MARK SCHEME for the May/June 2012 question paper for the guidance of teachers

9702 PHYSICS

9702/33

Paper 3 (Advanced Practical Skills 1), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2012 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2		Mark Scheme: Teachers' version	Syllabus	Paper		
		GCE AS/A LEVEL – May/June 2012	9702	33		
(a) (ii)	(a) (ii) Value of h_0 in range 0.70 m > h_0 > 0.50 m. Consistent with unit. [1]					
(b) (iii)	Valu	te of h , less than h_0 in (a)(ii) , with unit.		[1]		
		s of readings of h and m scores 5 marks, four sets scorelp from Supervisor –2 (setting up apparatus). Minor he		sor –1. [5]		
	nge o	f <i>m</i> : de 0.350 kg.		[1]		
	Column headings: [1]					
The	Each column heading must contain a quantity and a unit. The unit must conform to accepted scientific convention e.g. m / kg , $m(kg)$ or m in kg , $(h_0 - h)/m / m kg^{-1}$, $1/m / kg^{-1}$					
	nsiste	•		[1]		
		s of <i>h</i> must be given to the nearest mm.				
Sig		nt figures: nt figures for every row of values of 1/ <i>m</i> same as or or ole.	ne greater than	[1] m as recorded		
	lculati lues o	on: $f(h_0 - h)/m$ calculated correctly.		[1]		
(d) (i)	Sens Scal both Scal	s: sible scales must be used, no awkward scales (e.g. 3:1 les must be chosen so that the plotted points occupy x and y directions. les must be labelled with the quantity that is being plotte le markings must be no more than 3 large squares apa	at least half the	[1] e graph grid in		
	All o Dian	ting of points: bservations in the table must be plotted. neter of plots must be ≤ half a small square (no 'blobs') k to an accuracy of half a small square.		[1]		
		lity: points in the table must be plotted (at least 4) for this material ts must be less than 0.5 kg ⁻¹ (0.0005 g ⁻¹) of $1/m$ of a state of the contract o		[1] ded. Scatter of		
(ii)	Judg Thei Allov	of best fit: ge by balance of all points on the grid about the candidate me must be an even distribution of points either side of to whom one anomalous point only if clearly indicated by the community and the community of the community in the community and the community of the	he line along th andidate.			
(iii)	The	dient: hypotenuse of the triangle must be at least half the len read-offs must be accurate to half a small square in be	•			

1

Do not allow $\Delta x / \Delta y$.

			GCE AS/A LEVEL – May/June 2012	9702	33
		Read Or:	•		
	(e) Val	ue of F	P = candidate's gradient. Value of Q = candidate's inte	ercept.	[1]
	Uni	it for P	(e.g. m) consistent with value, and Q (m kg ⁻¹)		[1]
					[Total: 20]
2	(b) (ii)	Value	$ heta$ of $ heta_0$ to the nearest degree or 0.5° in range 70° # $ heta$	# 80°	[1]
	(iii)	Value	e of $ heta$ with unit, $ heta< heta_0$		[1]
	(iv)	Corre	ect calculation of $(\theta_0 - \theta)$		[1]
	(c) (i)	Value	of raw <i>d</i> with unit to nearest mm.		[1]
	(ii)	If rep	tute uncertainty in $2 \text{mm} < d < 5 \text{mm}$. eated readings have been taken, then the absolute. Correct method shown to find the percentage uncertainty.	•	[1] an be half the
	Sed Sed	cond va	alue of θ_0 within 1 °C of first value of θ_0 . alue of θ . alue of $\Delta \theta$ > first value of $\Delta \theta$ (check second value of of repeat readings of d here or in (c)(i) .	d > first value of	[1] [1] <i>d</i>). [1]
	(e) (i)	Two v	values of <i>k</i> calculated correctly.		[1]
	(ii)	Justifi	ication of s.f. in k linked to significant figures in d and	$\Delta \theta$.	[1]
	(iii)	Sensi	ible comment relating to the calculated values of	k, testing agai	nst a criterion

Mark Scheme: Teachers' version

Syllabus

Paper

Page 3

specified by the candidate.

[1]

Page 4	Mark Scheme: Teachers' version	Syllabus	Paper
	GCE AS/A LEVEL – May/June 2012	9702	33

(f)

	(i) Limitations 4 max.	(ii) Improvements 4 max.	No credit/not enough
A	two results not enough	take more readings <u>and plot a</u> <u>graph/</u> calculate more <i>k</i> values and <u>compare</u>	'repeat readings' on its own/ few readings/ take more readings and (calculate) average k/ only one reading
В	heat lost through sides and /or bottom	method to reduce heat loss/ lag/ insulate/ polystyrene container	use of lid/ heat loss in warming bowl/cup/ draughts/ heat loss to surroundings
С	temperature change is small/ $\Delta \theta$ values too close	time for longer/ higher starting temperature/ greater range of surface areas	
D	large (percentage) uncertainty in $\Delta \theta$	use thermometer with greater sensitivity or precision/ use thermometer that can read to 0.1°C	use more accurate thermometer/ thermometer not precise enough/ not just 'digital thermometer'
E	water in bowl barely covers (bulb of) thermometer	use larger volume of water/ use of thermocouple/ other <u>small</u> temperature <u>sensor</u> (e.g. probe)	not just 'digital thermometer' any reference to stirrer/ non-uniform temperature/ thermometer touching base
F	parallax error in measuring <u>d</u> / reason for difficulty in access in measuring <u>d</u>	use dividers/calipers	string measurements to measure <i>d</i>
G	difficult to mark level with reason	method of making mark stay e.g. depth gauge/ calibrated marks/ marker on outside	

Do not allow: use of coloured ink/reaction time/fans/draughts/water left behind/beakers not accurate/ helpers.

[Total: 20]