UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CHEMISTRY 0620/03

Paper 3

May/June 2004

1 hour 15 minutes

Candidates answer on the Question Paper. No Additional Materials required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen in the spaces provided on the Question Paper. You may use a pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, highlighters, glue or correction fluid. You may use a calculator.

Answer all questions.

The number of marks is given in brackets [] at the end of each question or part question. A copy of the Periodic Table is printed on page 12.

If you have been given a label, look at the details. If any details are incorrect or missing, please fill in your correct details in the space given at the top of this page.

Stick your personal label here, if provided.

For Examin	ier's Use
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of 12 printed pages.

IB04 06_0620_03/4RP © UCLES 2004



bat		reported from America that a turbine engine, the size of a button, might replas. The engine would be built from silicon which has suitable properties for tel.	
(a)	(i)	Why are batteries a convenient source of energy?	
			[1]
	(ii)	The engine will run on a small pack of jet fuel. What other chemical is needed burn this fuel?	to
			[1]
(b)	Sili	con has the same type of macromolecular structure as diamond.	
	(i)	Explain why one atom of either element can form four covalent bonds.	
			[2]
	(ii)	Predict two physical properties of silicon.	
			[2]
	(iii)	Name a different element that has a similar structure and properties to silicon.	
			[1]
(c)	Sili	con is made by the carbon reduction of the macromolecular compound, silicon(l	[V)
	(i)	Balance the equation for the reduction of silicon(IV) oxide.	
		SiO_2 + C \rightarrow Si + CO	[1]
	(ii)	Explain why the silicon(IV) oxide is said to be reduced.	[4]
	 (iii)	Describe the structure of silicon(IV) oxide. You may use a diagram.	[1]
			[2]

1

- 2 Sulphur is used to make sulphuric acid. In the UK, the annual production of the acid is about 2.5 million tonnes.
 - (a) The reactions in the manufacture of sulphuric acid by the Contact Process are shown below.


Sulphur			Sulphur dioxide	
S		reaction 1	SO ₂	
S	ulphur dioxide + oxygen	,	Sulphur trioxide	
	2SO ₂ + O ₂	reaction 2	2SO ₃	
	Sulphur trioxide		Oleum	
	SO ₃	reaction 3	$H_2S_2O_7$	
	Oleum + water		Sulphuric acid	
	$H_2S_2O_7$	reaction 4	H ₂ SO ₄	
(i)	Give a large scale source of the	element sulph	nur.	
			[1]	
(ii)	State another use of sulphur dio	xide.		
			[1]	
(iii)	How is sulphur changed into sul	phur dioxide?		
			[1]	
(iv)	Name the catalyst used in react	ion 2 .		
			[1]	
(v)	Reaction 2 is exothermic. Why i to increase the rate of this rever		ther than a higher temperature, used	
			[2]	
(vi)	Write a word equation for reaction	on 3 .		
			[1]	
(vii)	Write a symbol equation for read	ction 4.		
			[1]	

	4		
(b) About one third of this production of acid is used to make nitrogen and phosphorus-containing fertilisers.			
(i)	Name the third element that is essential for plant growth and is present in most fertilisers.		
	[1]		
(ii)	Name a nitrogen-containing fertiliser that is manufactured from sulphuric acid.		
	[1]		
(iii)	Rock phosphate (calcium phosphate) is obtained by mining. It reacts with concentrated sulphuric acid to form the fertiliser, superphosphate. Predict the formula of each of these phosphates.		
	fertiliser ions formula		
	calcium phosphate Ca ²⁺ and PO ₄ ³⁻		
	calcium superphosphate Ca ²⁺ and H ₂ PO ₄ ⁻ [2]		
(iv)	The ionic equation for the reaction between the phosphate ion and sulphuric acid is shown below.		
	PO_4^{3-} + $2H_2SO_4$ \rightarrow $H_2PO_4^-$ + $2HSO_4^-$		
	Explain why the phosphate ion is described as acting as a base in this reaction.		
	[2]		
An orga	anic compound decomposes to form nitrogen.		
С	$C_6H_5N_2Cl(aq)$ \rightarrow $C_6H_5Cl(I)$ + $N_2(g)$		
(a) Ex	xplain the state symbols.		
aq			
I			
g	[2]		


(b) Draw a diagram to show the arrangement of the valency electrons in **one** molecule of nitrogen.

3

(c) The rate of this reaction can be measured using the following apparatus.



The results of this experiment are shown on the graph below.



	(i)	How c	does the	rate o	of this	reaction	vary with	⊦time?
--	---	----	-------	----------	--------	---------	----------	-----------	--------

	[1]

(ii) Why does the rate vary?

	[2]

- (iii) The reaction is catalysed by copper powder. Sketch the graph for the catalysed reaction on the same grid. [2]
- (iv) Why is copper powder more effective as a catalyst than a single piece of copper?

- 4 (a) Insoluble compounds are made by precipitation.
 - (i) Complete the word equation for the preparation of zinc carbonate.


+	sodium carbonate →	zinc carbonate	+	
	odi 2011dio	oa. Dorraco		[2]

(ii) Complete the following symbol equation.

$$Pb(NO_3)_2$$
 + $NaCl \rightarrow$ + [2]

(iii) Write an ionic equation for the precipitation of the insoluble salt, silver(I) chloride.

(b) 2.0 cm³ portions of aqueous sodium hydroxide were added to 4.0 cm³ of aqueous iron(III) chloride. Both solutions had a concentration of 1.0 mol/dm³. After each addition, the mixture was stirred, centrifuged and the height of the precipitate of iron(III) hydroxide was measured. The results are shown on the following graph.



(i) Complete the ionic equation for the reaction.

$$Fe^{3+}$$
 +OH⁻ \rightarrow [1]

(ii) On the same grid, sketch the graph that would have been obtained if iron(II) chloride had been used instead of iron(III) chloride? [2]

	(III)	graph would be different. How are the shapes of these two graphs different and why?
		difference in shape
		reason for difference
		[2]
5		pper has the structure of a typical metal. It has a lattice of positive ions and a "sea" mobile electrons. The lattice can accommodate ions of a different metal.
	Giv	re a different use of copper that depends on each of the following.
	(i)	the ability of the ions in the lattice to move past each other
	()	
		[1]
	(ii)	the presence of mobile electrons
		[1]
	(iii)	the ability to accommodate ions of a different metal in the lattice
		[1]
	. , .	ueous copper(II) sulphate solution can be electrolysed using carbon electrodes. The s present in the solution are as follows.
		$Cu^{2+}(aq)$, $SO_4^{2-}(aq)$, $H^+(aq)$, $OH^-(aq)$
	(i)	Write an ionic equation for the reaction at the negative electrode (cathode).
		[1]
	(ii)	A colourless gas was given off at the positive electrode (anode) and the solution changes from blue to colourless.
		Explain these observations.
		[2]

(c)	re	queous copper(II) sulphate can be electrolysed using copper electrodes. T action at the negative electrode is the same but the positive electrode becom naller and the solution remains blue.	
	(i)	Write a word equation for the reaction at the positive electrode.	
			[1]
	(ii)	Explain why the colour of the solution does not change.	
			[2]
((iii)	What is the large scale use of this electrolysis?	
			[1]

6 In 2002, Swedish scientists found high levels of acrylamide in starchy foods that had been cooked above 120 °C. Acrylamide, which is thought to be a risk to human health, has the following structure.

$$C = C$$
 $CONH_2$

(a) (i) It readily polymerises to polyacrylamide. Draw the structure of this polymer.

[2]

(ii) Starch is formed by polymerisation. It has a structure of the type shown below. Name the monomer.



[1]

(iii) What are the differences between these two polymerisation reactions, one forming polyacrylamide and the other starch?

[2]

- **(b)** Acrylamide hydrolyses to form acrylic acid and ammonium ions.
 - (i) Describe the test for the ammonium ion.

test

result [2]

(ii) Given an aqueous solution, concentration 0.1 mol / dm³, how could you show that acrylic acid is a weak acid.

[2

(c) The structural formula of acrylic acid is shown below. It forms compounds called acrylates.

$$H$$
 $C = C$ H

(i) Acrylic acid reacts with ethanol to form the following compound.

$$\begin{array}{c} H \\ \\ \\ H \end{array} C = C \\ \begin{array}{c} COOCH_2CH_3 \\ \\ H \end{array}$$

Deduce the name of this compound. What type of organic compound is it?		
name		
type of compound		[2]

(ii) Acrylic acid is an unsaturated compound. It will react with bromine. Describe the colour change and draw the structural formula of the product of this addition reaction.

colour change	

structural formula of product

Chen react	nists use the concept of the mole to calculate the amounts of chemicals involved in a on.	l
(a) [Define mole.	
	[1]]
(b) 3	.0 g of magnesium was added to 12.0 g of ethanoic acid.	
N	$Mg + 2CH_3COOH \rightarrow (CH_3COO)_2Mg + H_2$	
T	he mass of one mole of Mg is 24 g.	
T	he mass of one mole of CH₃COOH is 60 g.	
(i) Which one, magnesium or ethanoic acid, is in excess? You must show your reasoning.	
	[3]]
(i	i) How many moles of hydrogen were formed?	
	[1]]
(ii	i) Calculate the volume of hydrogen formed, measured at r.t.p.	
	[2]
(c) li	n an experiment, 25.0cm^3 of aqueous sodium hydroxide, 0.4mol/dm^3 , was neutralised y 20.0cm^3 of aqueous oxalic acid, $H_2C_2O_4$.	ı
	$2NaOH + H2C2O4 \rightarrow Na2C2O4 + 2H2O$	
C	Calculate the concentration of the oxalic acid in mol/dm ³ .	
(Calculate the number of moles of NaOH in 25.0 cm ³ of 0.4 mol/dm ³ solution.	
	[1]]
(i	Use your answer to (i) and the mole ratio in the equation to find out the number of moles of H ₂ C ₂ O ₄ in 20 cm ³ of solution.	:
	[1]
(ii	i) Calculate the concentration, mol/dm³, of the aqueous oxalic acid.	
]

7

The Periodic Table of the Elements DATA SHEET

	_			_	_	_		
0	He Helium	20 Neon 10	40 Ar Argon	84 K rypton 36	131 Xe Xenon	Radon 86		175 Lu
₹		19 H Fluorine	35.5 C1 Chlorine	80 Br Bromine	127 I lodine 53	At Astatine 85		Yb
5		16 Oxygen			128 Te Tellurium			169 Tm
>		14 N Nitrogen 7	31 Phosphorus	75 AS Arsenic	122 Sb Antimony 51	209 Bi Bismuth		167 Er
≥		12 Carbon	28 Si Silicon	73 Ge Germanium	119 Sn Tin	207 Pb Lead 82		165 Ho
=		11 Boron 5	27 A1 Aluminium	70 Ga Gallium 31	115 In Indium	204 T 1 Thallium		162 Dy
				65 Zn Zinc 30	Cd Cadmium 48	201 Hg Mercury		159 Tb
				64 Cu Copper	108 Ag Silver 47	197 Au Gold		157 Gd
<u>.</u>				59 X Nickel 28	106 Pd Palladium 46	195 Pt Platinum 78		152 Eu
5				59 Co Cobalt	Rhodium 45	192 Ir Irdium		150 Sm
	T Hydrogen			56 Fe Iron	Ruthenium 44	190 Os Osmium 76		Pm
				55 Wn Manganese 25	Tc Technetium 43	186 Re Rhenium 75		144 Nd
				52 Cr Chromium	96 Mo Molybdenum 42	184 W Tungsten 74		141 Pr
				51 V Vanadium 23	93 Nb Nobium	181 Ta Tantalum		140 Ge
				48 T Titanium	91 Zr Zirconium	178 Hf Hafnium		
				Scandium 21		139 La Lanthanum 57 *	227 Ac Actinium 89	series
=		9 Be Beryllium	24 Mg Magnesium	Calcium	Strontium	137 Ba Barium 56	226 Ra Radium	*58-71 Lanthanoid series 90-103 Actinoid series
_		7 Li thium	23 Na Sodium	39 ★ Potassium 19	85 Rb Rubidium 37	133 CS Caesium 55	Fr Francium 87	*58-71 La
		1 III IV V VII VII Hydrogen 1 Hydro	III IV V VI VII VI		III IIV VV VVI VVII VVII			

90														
series bioc	140	141	144		150	152		159	162	165		169	173	175
TOID SELICES	ဝီ	Ŗ	PZ	Pm	Sm	Ш	gg	Д	٥	웃	ш	Ę	Υb	Ľ
מסוומס	Cerium 58	Praseodymium 59	Neodymium 60	Promethium 61	Samarium 62	Europium 63	9	Terbium 65	Dysprosium 66	Holmium 67	98	Thulium 69	Ytterbium 70	Lutetium 71
a = relative atomic mass	232		238											
X = atomic symbol	T	Ра)	d N	Pu		Cm		ర	Es	Fm	Md		בֿ
b = proton (atomic) number	Thorium 90	Protactinium 91	Uranium 92	Neptunium 93	Plutonium 94	Americium 95	96	Berkelium 97	Californium 98	0,	Fermium 100	Mendelevium 101	Nobelium 102	Lawrencium 103

The volume of one mole of any gas is 24 dm 3 at room temperature and pressure (r.t.p.).

Key