#### UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

International General Certificate of Secondary Education

## MARK SCHEME for the November 2004 question paper

#### 0620 CHEMISTRY

0620/03

Paper 3 (Extended Theory), maximum mark 150

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were initially instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began. Any substantial changes to the mark scheme that arose from these discussions will be recorded in the published *Report on the Examination*.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the *Report on the Examination*.

• CIE will not enter into discussion or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the November 2004 question papers for most IGCSE and GCE Advanced Level syllabuses.

**Grade thresholds** taken for Syllabus 0620 (Chemistry) in the November 2004 examination.

	maximum	minimum mark required for grade:				
	mark available	А	С	E	F	
Component 3	150	52	34	25	19	

The threshold (minimum mark) for B is set halfway between those for Grades A and C. The threshold (minimum mark) for D is set halfway between those for Grades C and E. The threshold (minimum mark) for G is set as many marks below the F threshold as the E threshold is above it.

Grade A\* does not exist at the level of an individual component.

### November 2004

## **INTERNATIONAL GCSE**

# MARK SCHEME

**MAXIMUM MARK: 150** 

**SYLLABUS/COMPONENT: 0620/03** 

CHEMISTRY Extended Theory

Page 1		Mark Scheme	Syllabus	Paper
		IGCSE – November 2004	0620	3

	(a)	carbon dioxide, water vapour, noble gases <b>or</b> a named noble gas Any <b>TWO</b>	[2]
	(b)	burning fossil fuels  COND that contain sulphur	[1] [1]
		acid rain <b>or</b> any effect of acid rain - deforestation, effect on stone work, corrosion of metals, acidity in lakes, health etc	[1]
		motor vehicles <b>or</b> petrol <b>or</b> car exhausts health, if specified then brain, nervous system, development of children etc do not select from list illnesses	[1] [1]
		OR lead in old paint harmful effect as above	[1] [1]
	(c) (i)	combustion <b>or</b> burning <b>NOT</b> dissolving in the ocean	[1]
	(ii)	6CO <sub>2</sub> + 6H <sub>2</sub> O exothermic	[1] [1]
	(d) (i)	glowing splint burst into flame or rekindled Must have glowing or equivalent idea	[1]
		<b>OR</b> any similar description that includes the two points glowing and relights.	
	(ii)	measure volume <b>or</b> count bubbles time	[1] [1]
		NOT units	
	(iii)	rate slows down  Because the reaction is photochemical <b>or</b> rate depends on intensity of light <b>or</b> light less bright or less light falling on plant <b>or</b> light provides energy for	[1]
		photosynthesis etc.	[1]
		TOTAL	= 15
		dilute filter	
		saturated cool	
		blue sulphate	[6]
		TOTAL	L = 6
}	(a) (i)	no change in concentration of reagents <b>or</b> rates equal Accept no change in amounts or it is as if the reaction has Stopped	[1]
	(ii)	back reaction is endothermic <b>or</b> the forward reaction is exothermic	[1]
		Increase in temperature favours the endothermic reaction which is the back reaction or vice versa.  NB look for correct conclusion re thermicity and comment re position of equilibrium.	[1]

Pag	ge 2	Mark Scheme	Syllabus	Paper
		IGCSE – November 2004	0620	3
	(iii) increased rate because molecules collide more frequently <b>or</b> concentration of molecules is			
		increased <b>or</b> molecules are closer <b>NOT</b> they have more KE		[1]
		increased yield high pressure favours side with few molecules <b>or</b> smaller	volume <b>or</b> m	
		to reduce the pressure this is product side this can be implied		[1] [1]
(b)	(i)	$CO_2$ and $H_2O$ balanced $2CH_3OH + 3O_2 = 2CO_2 + 4H_2O$		[1] [1]
	(ii)	methyl ethanoate		[1]
	(,	water		[1]
	(iii)	Methanoic (acid) accept formic acid		[1]
			-	ΓΟΤΑL = 13
4 (a)	(i)	Correct equation with a more reactive metal		[1]
	(ii)	Electron loss		[1]
	(iii)	Because they can accept electrons or take electrons aw from	ay	[1]
	(iv)	Silver or silver(I)		[1]
(b)	(i)	increase		[1]
	(ii)	zinc  COND and a correct reason - such as it loses electrons not it is more reactive  Need both zinc and reason for the mark.	nore easily <b>c</b>	r [1]
		(iii)from the more reactive to the less reactive NOT just	from zinc to I	ead [1]
				TOTAL = 7
5 (a)		Group II metals will lose 2e Group VI elements will gain 2e		[1] [1]
(b)		$\mathrm{SC}\mathit{l}_2$ <b>COND</b> 8e around both chlorine atoms 8e around sulphur with 2nbp and 2bp If x and o reversed ignore if this is the only error		[1] [1] [1]
(c)	(i)	lons cannot move in solid <b>or</b> can move in liquid		[1]
	(ii)	No ions in sulphur chloride <b>or</b> it is covalent <b>or</b> only molestrontium chloride has ions	cules <b>or</b> only	[1]
				TOTAL = 7

Mark Scheme

Page 2

Syllabus

Paper

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – November 2004	0620	3

correct structure $CH_2=CCl_2$	[1]
because it has a lower $M_r$ or density or its molecules move faster it is lighter ONLY [1] only comment - smaller molecules [0] answer implies or states sieve idea then [0]	[2]
ester linkage COND polymer chain showing different monomers and continuation -OOC-C <sub>6</sub> H <sub>4</sub> -COOCH <sub>2</sub> CH <sub>2</sub> O-	[1] [1]
fats or lipids	[1]
does not decompose easily when heated accept similar statements	[1]
does not decompose <b>or</b> non-biodegradable shortage of landfill sites <b>or</b> o space visual pollution poisonous/toxic/harmful gases when <u>burnt</u> <b>NOT</b> carbon monoxide, sulphur dioxide. If gas named has to be a correct one eg HC <i>l</i> , HCN dangerous to animals	f
Any <b>TWO</b>	[2]
conserve petroleum or save energy <b>NOT</b> cheaper	[1]
•	AL = 10
$Zn(OH)_2 = ZnO + H_2O$ reactant [1] products [1]	[2]
it would melt <b>or</b> it does not decompose <b>or</b> it does not react <b>NOT</b> no change	[1]
blue (solid) to black (solid) brown gas	[1] [1] [1]
Mark consequentially to any error <b>but not involving simple integers</b> There has to be some evidence that the candidate has attempted to work through the calculation and not merely inserted whole numbers. For example 2, 1, 160 or 1, 0.5, 80 number of moles of $Fe_2(SO_4)_3 = 1/40$ or $0.025$ number of moles of $Fe_2O_3$ formed = $1/40$ or $0.025$ mass of iron(III) oxide formed = $0.025 \times 160 = 4g$ number of moles of $SO_3$ produced = $3/40$ or $0.075$ volume of sulphur trioxide at r.t.p. = $0.075 \times 25$ = $1.8 \text{dm}^3$	[5]
)	because it has a lower <i>M<sub>r</sub></i> or density or its molecules move faster it is lighter ONLY [1] only comment - smaller molecules [0] answer implies or states sieve idea then [0] ester linkage COND polymer chain showing different monomers and continuation -OOC-C <sub>6</sub> H <sub>4</sub> -COOCH <sub>2</sub> CH <sub>2</sub> O-fats or lipids  does not decompose easily when heated accept similar statements  does not decompose or non-biodegradable shortage of landfill sites or o space visual pollution poisonous/toxic/harmful gases when <u>burnt</u> NOT carbon monoxide, sulphur dioxide. If gas named has to be a correct one eg HC <i>I</i> , HCN dangerous to animals Any TWO  conserve petroleum or save energy  NOT cheaper  TOT  Zn(OH) <sub>2</sub> = ZnO + H <sub>2</sub> O reactant [1] products [1]  it would melt or it does not decompose or it does not react NOT no change  blue (solid) to black (solid) brown gas  Mark consequentially to any error but not involving simple integers There has to be some evidence that the candidate has attempted to work through the calculation and not merely inserted whole numbers. For example 2, 1, 160 or 1, 0.5, 80 number of moles of Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> = 1/40 or 0.025 number of moles of Fe <sub>2</sub> (SO <sub>4</sub> ) <sub>3</sub> = 1/40 or 0.025 number of moles of Fe <sub>2</sub> (SO <sub>3</sub> ) formed = 1/40 or 0.025 number of moles of Fe <sub>2</sub> (SO <sub>3</sub> ) formed = 1/40 or 0.075 volume of sulphur trioxide at r.t.p. = 0.075 x 25

**TOTAL = 11** 

	raye 4	Walk Scheme	Syllabus	rapei
		IGCSE – November 2004	0620	3
8	(a) (i)	C <sub>6</sub> H <sub>12</sub> between 60 to 65°C		[1] [1]
	(ii)	C <sub>12</sub> H <sub>24</sub> COND giving some indication of the method		[1] [1]
	(b)	add bromine water <b>or</b> potassium manganate(VII) butene it goes from brown/orange/yellow to colourless		[1]
		or manganate (VII) from pink to colourless  NOT clear		[1]
		Cyclobutane it remains brown/orange/yellow or mangana or no colour change Accept does not react Provided colour of reagent somewhere in the answer [3]	, , ,	pink [1]
	(c) (i)	alcohol		[1]
	(ii)	CH <sub>3</sub> -CH <sub>2</sub> -CHC <i>l</i> -CH <sub>3</sub>		[1]
	(iii)	-CH(CH <sub>3</sub> )-CH(CH <sub>3</sub> )- or any equivalent diagram [1] for repeat unit and [1] for continuation		[2]
			-	TOTAL = 11

Mark Scheme

Page 4

Syllabus

Paper