#### **Location Entry Codes**

As part of CIE's continual commitment to maintaining best practice in assessment, CIE uses different variants of some question papers for our most popular assessments with large and widespread candidature. The question papers are closely related and the relationships between them have been thoroughly established using our assessment expertise. All versions of the paper give assessment of equal standard.

The content assessed by the examination papers and the type of questions is unchanged.

This change means that for this component there are now two variant Question Papers, Mark Schemes and Principal Examiner's Reports where previously there was only one. For any individual country, it is intended that only one variant is used. This document contains both variants which will give all Centres access to even more past examination material than is usually the case.

The diagram shows the relationship between the Question Papers, Mark Schemes and Principal Examiners' Reports that are available.

Question Paper	Mark Scheme	Principal Examiner's Report
Introduction	Introduction	Introduction
First variant Question Paper	First variant Mark Scheme	First variant Principal Examiner's Report
Second variant Question Paper	Second variant Mark Scheme	Second variant Principal Examiner's Report

Who can I contact for further information on these changes?

Please direct any questions about this to CIE's Customer Services team at: <a href="mailto:international@cie.org.uk">international@cie.org.uk</a>

The titles for the variant items should correspond with the table above, so that at the top of the first page of the relevant part of the document and on the header, it has the words:

• First variant Question Paper / Mark Scheme / Principal Examiner's Report

or

Second variant Question Paper / Mark Scheme / Principal Examiner's Report

as appropriate.





# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/31
Paper 3 (Extended)		October/Nover	nber 2008
		1 hour 1	5 minutes
Candidates and	swer on the Question Paper.		
No Additional N	Materials are required		

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES

Answer all questions.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part questions.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of 12 printed pages.



1 Complete the following table.

For Examiner's Use

gas	test for gas
ammonia	
	bleaches damp litmus paper
hydrogen	
	relights a glowing splint
	turns limewater milky

[Total: 5]

There are three types of giant structure – ionic, metallic and macromolecular. (a) Sodium nitride is an ionic compound. Draw a diagram that shows the formula of the compound, the charges on the ions and the arrangement of the valency electrons around the negative ion. Use x to represent an electron from a sodium atom. Use o to represent an electron from a nitrogen atom. [3] (b) (i) Describe metallic bonding. [3] (ii) Use the above ideas to explain why metals are good conductors of electricity, [1] metals are malleable. [2] (c) Silicon(IV) oxide has a macromolecular structure. (i) **Describe** the structure of silicon(IV) oxide (a diagram is not acceptable). [3] (ii) Diamond has a similar structure and consequently similar properties. Give **two** physical properties common to both diamond and silicon(IV) oxide.

For Examiner's Use

[Total: 14]

2

Steel is an alloy made from impure iron. (a) Both iron and steel rust. The formula for rust is Fe<sub>2</sub>O<sub>3</sub>.2H<sub>2</sub>O. It is hydrated iron(III) oxide. (i) Name the **two** substances that must be present for rusting to occur. (ii) Painting and coating with grease are two methods of preventing iron or steel from rusting. Give two other methods. (b) (i) Name a reagent that can reduce iron(III) oxide to iron. [1] (ii) Write a symbol equation for the reduction of iron(III) oxide, Fe<sub>2</sub>O<sub>3</sub>, to iron. [2] (c) (i) Calculate the mass of one mole of Fe<sub>2</sub>O<sub>3</sub>.2H<sub>2</sub>O. [1] ..... (ii) Use your answer to (i) to calculate the percentage of iron in rust. [2] ..... (d) Iron from the blast furnace is impure. Two of the impurities are carbon and silicon. These are removed by blowing oxygen through the molten iron and adding calcium oxide. (i) Explain how the addition of oxygen removes carbon. (ii) Explain how the addition of oxygen and calcium oxide removes silicon.


For Examiner's Use

3

[Total: 13]

4 Across the world, food safety agencies are investigating the presence of minute traces of the toxic hydrocarbon, benzene, in soft drinks. It is formed by the reduction of sodium benzoate by vitamin C.

For Examiner's Use



(a) Sodium benzoate is a salt, it has the formula  $C_6H_5COONa$ . It can be made by the neutralisation of benzoic acid by sodium hydroxide.

(i)	Deduce the formula of benzoic acid.	
		[1]
(ii)	Write a word equation for the reaction between benzoic acid and sodium hydro	xide
		[1]
(iii)	Name <b>two</b> other compounds that would react with benzoic acid to form sodium benzoate.	
		[2]
(b) Ber	nzene contains 92.3% of carbon and its relative molecular mass is 78.	
(i)	What is the percentage of hydrogen in benzene?	
		[1]
(ii)	Calculate the ratio of moles of C atoms: moles of H atoms in benzene.	
		••
		[2]
(iii)	Calculate its empirical formula and <b>then</b> its molecular formula.	
	The empirical formula of benzene is	

The molecular formula of benzene is

[2]

**(c)** The structural formula of Vitamin C is drawn below.

For Examiner's Use

[Total: 12]

(i)	What is its molecular formula?	
		[1]
(ii)	Name the two functional groups which are circled.	
		[2]


5

The electrolysis of concentrated aqueous sodium chloride produces three commercially important chemicals hydrogen, chlorine and sodium hydroxide.			
(a) The ions	s present are Na $^{+}$ (aq), H $^{+}$ (aq) ,C $l^{-}$ (aq) and OH $^{-}$ (aq).		
(i) Con	nplete the ionic equation for the reaction at the negative electrode (cathode)	١.	
	+ H <sub>2</sub>	[1]	
(ii) Con	nplete the ionic equation for the reaction at the positive electrode (anode).		
	C1₂	[1]	
(iii) Exp	lain why the solution changes from sodium chloride to sodium hydroxide.		
		[1]	
<b>(b) (i)</b> Why	y does the water supply industry use chlorine?		
		[1]	
(ii) Nan	ne an important chemical that is made from hydrogen.		
		[1]	
(iii) How	v is sodium hydroxide used to make soap?		
		[2]	
	[Tota	l: 7]	

6 The reactivity series lists metals in order of reactivity.

For Examiner's Use

(a) To find out which is the more reactive metal, zinc or tin, the following experiment could be carried out.



This experiment could be carried out with other metals and the results recorded in a table. Then the order of reactivity can be deduced.

(i) The order was found to be:

manganese most reactive

zinc tin

silver least reactive

Complete the table of results from which this order was determined.

aqueous	tin	manganese	silver	zinc
solution	Sn	Mn	Ag	Zn
tin(II) nitrate		R	NR	R
manganese(II) nitrate				
silver(I) nitrate				
zinc nitrate				

•
. 3

(ii) Write the ionic equation for the reaction between tin atoms and silver(I	) ions
---	--------

[2]

	(iii)	The following is a redox reaction.	
		$Mn + Sn^{2+} \longrightarrow Mn^{2+} + Sn$	
		Indicate on the equation the change which is oxidation. Give a reason for your choice.	
			[2]
	(iv)	Explain why experiments of this type cannot be used to find the position aluminium in the reactivity series.	of
			[2]
(b)	ions	assium and calcium are very reactive metals at the top of the series. Because to have different charges, $K^+$ and $Ca^{2^+}$ , their compounds behave differently wated.	
	(i)	Explain why the ions have different charges.	
			[2]
	(ii)	Their hydroxides are heated. If the compound decomposes, complete the word equation. If it does not decompose, write "no reaction".	
		Potassium hydroxide —	
		Calcium hydroxide	[2]
	(iii)	Complete the equations for the decomposition of their nitrates.	
		2KNO₃ → +	
		2Ca(NO <sub>3</sub> ) <sub>2</sub> + + + + + + + + + + + + + + + + + + +	[4]
		[Total:	17]

The crac		anes are generally unreactive. Their reactions include combustion, substitution g.	and	
(a)	The	e complete combustion of an alkane gives carbon dioxide and water.		
	(i)	10 cm <sup>3</sup> of butane is mixed with 100 cm <sup>3</sup> of oxygen, which is an excess. The mix is ignited. What is the volume of unreacted oxygen left and what is the volunt carbon dioxide formed?		
		$C_4H_{10}(g) + 6\frac{1}{2}O_2(g) \longrightarrow 4CO_2(g) + 5H_2O(I)$		
		Volume of oxygen left = cm <sup>3</sup>		
		Volume of carbon dioxide formed = cm <sup>3</sup>	[2]	
	(ii)	Why is the incomplete combustion of any alkane dangerous, particularly is enclosed space?	n an	
			[2]	
(b)	The	e equation for a substitution reaction of butane is given below.		
		$CH_3-CH_2-CH_2-CH_3 + Cl_2 \longrightarrow CH_3-CH_2-CH_2-CH_2-Cl + HCl$		
	(i)	Name the organic product.		
			[1]	
	(ii)	This reaction does not need increased temperature or pressure. What is the essential reaction condition?		
			[1]	
(	iii)	Write a different equation for a substitution reaction between butane and chloring	ne.	
			[1]	

7

(c)	Alkenes are more reactive and industrially more useful than alkanes. They are made by cracking alkanes.				
		C <sub>7</sub> H <sub>16</sub> − heptane		+ CH <sub>3</sub> -CH <sub>2</sub> -CH=CH <sub>2</sub> + H <sub>2</sub> but-1-ene	
	(i)	Draw the structura	l formula of the polyr	/mer poly(propene).	
					[2]
	(ii)	Give the structural with steam.	formula and name	of the alcohol formed when but-1-ene rea	acts
		name			[1]
		structural formula			
					[1]
	(iii)	Deduce the struct hydrogen chloride.		e product formed when propene reacts v	with
					[1]
				[Total:	12]

DATA SHEET
The Periodic Table of the Elements

	0	4 <b>He</b> Helium	20 Neon 10 At Argon 18	84 <b>Kr</b> Krypton 36	131 <b>Xe</b> Xenon 54	Rn Radon 86		175 <b>Lu</b> Lutetium 71	<b>Lr</b> Lawrencium 103
	II/		19 Fluorine 9 35.5 <b>C l</b>	80 <b>Br</b> Bromine 35	127 <b>I</b> lodine 53	At Astatine 85		<b>Yb</b> Ytterbium 70	Nobelium
	IN		16 Oxygen 8 32 \$ \$ Sulphur	Se Selenium 34	128 <b>Te</b> Tellurium	Po Polonium 84		169 <b>Tm</b> Thulium	Md Mendelevium 101
	^		Nitrogen 7 31 9 Phosphorus 15	75 <b>AS</b> Arsenic	Sb Antimony 51	209 <b>Bi</b> Bismuth		167 <b>Er</b> Erbium 68	Fm Fermium
	ΛΙ		Carbon 6 Carbon 8 Silicon 14	73 <b>Ge</b> Germanium 32	119 <b>S</b> Tin	207 <b>Pb</b> Lead		165 <b>Ho</b> Holmium 67	Es Einsteinium 99
	=		11  B Boron 5 27 At Aluminium 13	70 <b>Ga</b> Gallium 31	115 <b>In</b> Indium 49	204 <b>T t</b> Thallium 81		162 <b>Dy</b> Dysprosium 66	Cf Californium 98
				65 <b>Zn</b> Zinc 30	112 <b>Cd</b> Cadmium 48	201 <b>Hg</b> Mercury 80		159 <b>Tb</b> Terbium 65	<b>BK</b> Berkelium 97
				64 <b>Cu</b> Copper	108 <b>Ag</b> Silver 47	197 <b>Au</b> Gold		157 <b>Gd</b> Gadolinium 64	Cm Curium
Group				59 Nickel	106 <b>Pd</b> Palladium 46	195 <b>Pt</b> Platinum 78		152 <b>Eu</b> Europium 63	Am Ameridum 95
Ď				59 <b>Co</b> Cobalt	103 <b>Rh</b> Rhodium 45	192 <b>Ir</b> Iridium 77		Sm Samarium 62	Pu Plutonium 94
		T Hydrogen		56 <b>Fe</b> Iron	Ru Ruthenium	190 <b>Os</b> Osmium 76		Pm Promethium 61	Neptunium
				Manganese	Tc Technetium	186 <b>Re</b> Rhenium 75		144 <b>Nd</b> Neodymium 60	238 <b>U</b> Uranium 92
				52 <b>Cr</b> Chromium 24	96 <b>Mo</b> Molybdenum 42	184 <b>W</b> Tungsten 74		Pr Praseodymium 59	Pa Protactinium 91
				51 V Vanadium 23	Niobium N1	181 <b>Ta</b> Tantalum 73		140 <b>Ce</b> Cerium	232 <b>Th</b> Thorium
				48 Tritanium 22	2r Zirconium 40	178 <b>Hf</b> Hafnium * 72		1	nic mass bol nic) number
				Scandium 21	89 Yttrium 39	139 <b>La</b> Lanthanum 57 *	227 <b>Ac</b> Actinium	d series series	a = relative atomic mass  X = atomic symbol  b = proton (atomic) number
	=		Be Beryllum 4 24 Magnesium 12	40 Calcium	Strontium	137 <b>Ba</b> Barium 56	226 <b>Ra</b> Radium	*58-71 Lanthanoid series 190-103 Actinoid series	« <b>×</b> ∞
	_		7 Lithium 3 23 Rodum 11	39 <b>K</b> Potassium 19	Rubidium	133 <b>CS</b> Caesium 55	<b>Fr</b> Francium 87	*58-71 L	Key

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).





# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/32
Paper 3 (Exten	ded)	October/Novem	ber 2008
		1 hour 15	minutes
Candidates ans	swer on the Question Paper.		

### **READ THESE INSTRUCTIONS FIRST**

No Additional Materials are required.

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO **NOT** WRITE IN ANY BARCODES

Answer all questions.

A copy of the Periodic Table is printed on page 12.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part questions.

For Examiner's Use			
1			
2			
3			
4			
5			
6			
7			
Total			

This document consists of 12 printed pages.



1 Complete the following table.

For Examiner's Use

gas	test for gas
	turns damp red litmus paper blue
	bleaches damp litmus paper
hydrogen	
oxygen	
carbon dioxide	

[Total: 5]

There are three types of giant structure – ionic, metallic and macromolecular. (a) Sodium sulphide is an ionic compound. Draw a diagram that shows the formula of the compound, the charges on the ions and the arrangement of the valency electrons around the negative ion. Use x to represent an electron from a sodium atom. Use o to represent an electron from a sulphur atom. [3] (b) (i) Describe metallic bonding. (ii) Use the above ideas to explain why metals are good conductors of electricity, [1] metals are malleable. [2] (c) Silicon(IV) oxide has a macromolecular structure. (i) **Describe** the structure of silicon(IV) oxide (a diagram is not acceptable). [3] (ii) Diamond has a similar structure and consequently similar properties. Give **two** physical properties common to both diamond and silicon(IV) oxide. [Total: 14]

Examiner's Use

For

2


Ste	el is	an alloy made from impure iron.					
(a)		oth iron and steel rust. The formula for rust is $Fe_2O_3.2H_2O$ . s hydrated iron(III) oxide.					
	(i)	Name the <b>two</b> substances that must be present for rusting to occur.					
		and	[2]				
	(ii)	Painting and coating with grease are two methods of preventing iron or steel rusting. Give <b>two</b> other methods.	from				
			[2]				
(b)	(i)	Name a reagent that can reduce iron(III) oxide to iron.					
			[1]				
	(ii)	Write a symbol equation for the reduction of iron(III) oxide, Fe <sub>2</sub> O <sub>3</sub> , to iron.					
			[2]				
(c)	(i)	Calculate the mass of one mole of Fe <sub>2</sub> O <sub>3</sub> .2H <sub>2</sub> O.					
			[1]				
	(ii)	Use your answer to (i) to calculate the percentage of water in rust.					
			[2]				
(d)		n from the blast furnace is impure. Two of the impurities are carbon and silicese are removed by blowing oxygen through the molten iron and adding calced.					
	(i)	Explain how the addition of oxygen removes carbon.					
			[1]				
	(ii)	Explain how the addition of oxygen and calcium oxide removes silicon.					
			[2]				

3

[Total: 13]

4 Across the world, food safety agencies are investigating the presence of minute traces of the toxic hydrocarbon, benzene, in soft drinks. It is formed by the reduction of sodium benzoate by vitamin C.

For Examiner's Use



- (a) Sodium benzoate is a salt, it has the formula C<sub>6</sub>H<sub>5</sub>COONa. It can be made by the neutralisation of benzoic acid by sodium hydroxide.
  - (i) Deduce the formula of benzoic acid.

ſ	1	•
 L	-	•

(ii) Write a word equation for the reaction between benzoic acid and sodium hydroxide.

[ A ]
- 111
F.1

(iii) Name **two** other compounds that would react with benzoic acid to form sodium benzoate.

[2	1
L-	٦,


(b)	Ber	enzene contains 92.3% of carbon and its relative molecular mass is 78.				
	(i)	What is the percentage of hydrogen in benzene?				
	(ii)	Calculate the ratio of moles of C atoms: moles of H atoms in benzene.				
			[2]			
			[~]			
	(iii)	Calculate its empirical formula and <b>then</b> its molecular formula.				
		The empirical formula of benzene is				
		The molecular formula of benzene is	[2]			
(c)	The	e structural formula of Vitamin C is drawn below.				
		$O = C \qquad C - C - C - OH$ $O = C \qquad OH \qquad H$ $O = C \qquad OH \qquad H$				
	(i)	What is its molecular formula?				
			[1]			
	(ii)	Name the two functional groups which are circled.				
			[2]			
		[Total	: 12]			

	ectrolysis of concentrated aqueous sodium chloride produces three commercent chemicals; hydrogen, chlorine and sodium hydroxide.	cially				
(a) The	(a) The ions present are Na $^+$ (aq), H $^+$ (aq) ,C $l^-$ (aq) and OH $^-$ (aq).					
(i)	Complete the ionic equation for the reaction at the negative electrode (cathode	).				
	+ <u>H</u> 2	[1]				
(ii)	Complete the ionic equation for the reaction at the positive electrode (anode).					
	2C <i>l</i> ⁻	[1]				
(iii)	Explain why the solution changes from sodium chloride to sodium hydroxide.					
		[1]				
(b) (i)	Why does the water supply industry use chlorine?					
(::)	Name on important the original that is used a from budge one	[1]				
(ii)	Name an important chemical that is made from hydrogen.					
		[1]				
(iii)	Sodium hydroxide reacts with fats to make soap and glycerine What type of compound are fats?					
		[1]				
	What type of the reaction is this?					
		[1]				
	[Tota	ı./]				

5

6 The reactivity series lists metals in order of reactivity.

- For Examiner's Use
- (a) To find out which is the more reactive metal, zinc or tin, the following experiment could be carried out.



This experiment could be carried out with other metals and the results recorded in a table. Then the order of reactivity can be deduced.

(i) The order was found to be:

manganese most reactive

zinc tin

silver least reactive

Complete the table of results from which this order was determined.

aqueous	tin	manganese	silver	zinc
solution	Sn	Mn	Ag	Zn
tin(II) nitrate		R	NR	R
manganese(II) nitrate				
silver(I) nitrate				
zinc nitrate				

	_	
- 1	-4	1
L	٠.	"」

(ii) Write the equation for the reaction between zinc and silver(I) nitrate.

[2]

	(iii)	The following is a redox reaction.				
		$Mn + Sn^{2+} \longrightarrow Mn^{2+} + Sn$				
		Indicate on the equation which reagent is the oxidant or oxidizing agent. Give a reason for your choice.				
		[2]				
	(iv)	Explain why experiments of this type cannot be used to find the position of aluminium in the reactivity series.				
		[2]				
<b>(b)</b> Potassium and calcium are very reactive metals at the top of the series. Bec ions have different charges, K <sup>+</sup> and Ca <sup>2+</sup> , their compounds behave differe heated.						
	(i)	Explain why the ions have different charges.				
		[2]				
	(ii)	Their hydroxides are heated. If the compound decomposes, complete the word equation. If it does not decompose, write "no reaction".				
		Potassium hydroxide →				
		Calcium hydroxide — [2]				
	(iii)	Complete the equations for the decomposition of their nitrates.				
		2KNO₃ → +				
		$2Ca(NO_3)_2 \longrightarrow                                   $				
		[Total: 17]				

7 The alkanes are generally unreactive. Their reactions include combustion, substitution and cracking. (a) The complete combustion of an alkane gives carbon dioxide and water. (i) 20 cm<sup>3</sup> of butane is mixed with 150 cm<sup>3</sup> of oxygen, which is an excess. The mixture is ignited. What is the volume of unreacted oxygen left and what is the volume of carbon dioxide formed?  $C_4H_{10}(g) + 6\frac{1}{2}O_2(g) \longrightarrow 4CO_2(g) + 5H_2O(I)$ ..... cm<sup>3</sup> Volume of oxygen left = ..... cm<sup>3</sup> Volume of carbon dioxide formed = [2] (ii) Why is the incomplete combustion of any alkane dangerous, particularly in an enclosed space? **(b)** The equation for a substitution reaction of butane is given below.  $CH_3-CH_2-CH_2-CH_3 + Cl_2 \longrightarrow CH_3-CH_2-CH_2-CH_2-Cl_1 + HCl_2$ (i) Name the organic product. [1] (ii) This reaction does not need increased temperature or pressure. What is the essential reaction condition? [1] (iii) Write a different equation for a substitution reaction between butane and chlorine.

For Examiner's Use

[1]

		11						
(c)	Alkenes are more reactive and industrially more useful than alkanes.  They are made by cracking alkanes.							
		$C_7H_{16} \longrightarrow CH_3-CH=CH_2 + CH_3-CH_2-CH=CH_2 + H_2$ heptane propene but-1-ene						
	(i)	Draw the structural formula of the polymer poly(propene).						
		[2]						
	(ii)	Give the structural formula and name of the alcohol formed when propene reacts with steam.						
		name[1]						
		structural formula						
(	(iii)	[1] Deduce the structural formula of the product formed when but-1-ene reacts with hydrogen chloride.						

[1]

For Examiner's Use

[Total: 12]

DATA SHEET
The Periodic Table of the Elements

	0	4 He Helium	20 <b>Ne</b> Neon	40 <b>Ar</b> Argon	84 Krypton	36	<u></u> 8	Xenon 54	ć	Radon 86		175 <b>Lu</b> Lutetium 71		۲	Lawrencium 103
	II/		19 <b>F</b> Fluorine 9	35.5 <b>C 1</b> Chlorine	80 <b>Br</b> Bromine	35	<b>I</b>	lodine 53	Š	At Astatine 85		173 <b>Yb</b> Ytterbium 70		8	Nobelium 102
	>		16 <b>O</b> Oxygen 8	32 <b>S</b> Sulphur 16	79 <b>Se</b> Selenium	34	٦ <b>و</b>	Tellurium 52	ć	Polonium 84		169 <b>Tm</b> Thulium 69		Md	Mendelevium 101
			14 <b>N</b> Nitrogen 7	31 Phosphorus 15	75 <b>AS</b> Arsenic	33	Sb	Antimony 51	209	Bismuth 83		167 <b>Er</b> Erbium 68		Fm	Fermium 100
	2	:	12 <b>C</b> Carbon 6	28 <b>Si</b> Silicon	73 <b>Ge</b> Germanium	32	Sn		207	Lead 82		165 <b>Ho</b> Holmium 67		Es	Einsteinium 99
Group	≡		11 Boron 5	27 <b>A1</b> Aluminium	70 <b>Ga</b>	31	In	Indium 49	204	Thallium 81		162 <b>Dy</b> Dysprosium 66		రే	Californium 98
					65 <b>Zn</b> Zinc	30	S	Cadmium 48	201	Mercury 80		159 <b>Tb</b> Terbium 65		Æ	Berkelium 97
					64 Copper	29	<sup>™</sup> Ag		197	Au Gold 79		157 <b>Gd</b> Gadolinium 64		Cm	Curium 96
					S9 Nickel	28	Pq 1	Palladium 46	195	Platinum 78		152 <b>Eu</b> Europium 63		Am	Americium 95
Ğ					S9 Cobalt	27	<b>8</b>	Rhodium 45	192	II Iridium 77		Sm Samarium 62		Pn	Plutonium 94
		1 Hydrogen			56 Iron	26	R <sub>u</sub>	Ruthenium 44	190	Osmium 76		Pm Promethium 61	_	Ν d	Neptunium 93
					Manganese	25	٦ <sub></sub>	Technetium 43	186	Rhenium 75		Neodymium 60	238	<b>-</b>	Uranium 92
					52 Chromium	24	» <b>e</b>	Molybdenum 42	184	Tungsten 74		141 <b>Pr</b> Praseodymium 59		Ра	Protactinium 91
					51 <b>V</b> Vanadium	23	S Q	Niobium 41	181	Tantalum 73		140 <b>Ce</b> Cerium 58	232	두	Thorium 90
					48 Titanium	22	Ž	Zirœnium 40	178	Hafnium 72		1	mic mass	loqu	nic) number
				I	45 <b>Sc</b> Scandium	21	ଛ ≻	Yttrium 39	139	Lanthanum 57	227 <b>AC</b> Actinium	d series series	a = relative atomic mass	X = atomic symbol	b = proton (atomic) number
	=		9 <b>Be</b> Beryllium	24 <b>Mg</b> Magnesium		20	დ	Strontium 38	137	Barium 56	226 <b>Ra</b> Radium	*58-71 Lanthanoid series	a	× ×	٩ 
	-		7 <b>Li</b> Lithium 3	23 Na Sodium	39 <b>X</b> Potassium	9	8 <b>8</b>	Rubidium 37	133	Caesium 55	<b>Fr</b> Francium 87	*58-71 L		Key	Ω

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

The volume of one mole of any gas is  $24\,\mathrm{dm}^3$  at room temperature and pressure (r.t.p.).