

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CHEMISTRY		0620/33
CENTRE NUMBER	CANDIDATE NUMBER	
CANDIDATE NAME		

Paper 3 (Extended)

October/November 2010

1 hour 15 minutes

Candidates answer on the Question Paper.

No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

A copy of the Periodic Table is printed on page 16.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [] at the end of each question or part question.

For Examiner's Use	
1	
2	
3	
4	
5	
6	
7	
Total	

This document consists of 14 printed pages and 2 blank pages.



1 The diagrams below show the electron arrangement in two compounds.

00 1	$\times \times$
$\overset{\circ}{K}\overset{\bullet}{\overset{\bullet}{K}}$	${}^{ imes}_{\circ}Cl_{ imes}^{ imes}$
\circ \cap	$\bigcirc \cup \iota_{\times}$
\cap	$\times \times$



(a)	In a water molecule, each hydrogen atom is bonded to the oxygen atom by sharing a pair of electrons. Why does an oxygen atom share two pairs of electrons rather than just one pair?
	[1]
(b)	Describe how a potassium atom becomes a potassium ion.
(2)	
(C)	Why is there a bond between the ions in potassium chloride?
	[1]
(d)	Solid potassium chloride is a poor conductor of electricity. When dissolved in water it is a good conductor. Explain.
	[2]
	[Total: 5]

- 2 Vanadium is a transition element.
 - (a) An atom of the most common isotope of vanadium can be represented as $^{51}_{23}\mathrm{V}$.

Complete the following table to show the number of protons, electrons and neutrons in each particle.

particle	number of protons	number of electrons	number of neutrons
⁵¹ ₂₃ V			
⁵¹ ₂₃ V ³⁺			
⁵⁰ ₂₃ V			

[3]


(b)	The	e major use of vanadium is to make vanadium steel alloys.
	(i)	Explain the phrase steel alloys.
		[2]
	(ii)	State the name and use of another steel alloy.
		name
		use[2]
(c)	Two	o of the oxidation states of vanadium are +3 and +4.
	(i)	Write the formula of vanadium(III) oxide and of vanadium(IV) oxide.
		vanadium(III) oxide
		vanadium(IV) oxide[2]
	(ii)	lem:lem:lem:lem:lem:lem:lem:lem:lem:lem:
		[3]
		[Total: 12]

- 3 The reactions of a metal and the thermal stability of some of its compounds are determined by the position of the metal in the reactivity series.
 - (a) To find the order of reactivity of the metals, cobalt, magnesium, silver and tin, the following experiments were carried out.

experiment	result
tin plus silver(I) nitrate solution	silvery layer on tin
magnesium plus tin(II) nitrate solution	grey deposit on magnesium
tin plus cobalt nitrate solution	no reaction

		tin plus cobalt nitrate solution	no reaction	
	(i)	Give as far as possible the order of reac Write the least reactive first.	tivity of these metals.	
				[2]
	(ii)	What additional experiment needs to be reactivity?	oe done to put all four metal:	s in order of
				[1]
	(iii)	Write an ionic equation for the reaction b on the equation the change which is oxide	The state of the s	ons. Indicate
				[3]
(b)		lium is a more reactive metal than magner magnesium compounds.	esium. Sodium compounds are	more stable
		n experiment, their hydroxides were heate reaction' otherwise complete the equatio	•	ompose write
	NaC	DH →		
	Mg(OH) ₂ →		[2]

(c) A cell consists of two different metal electrodes in an electrolyte. Three possible cells are shown below.



(i)	Why is the more reactive metal the negative electrode?	
(ii)	How can you deduce that zinc is more reactive than tin?	[∠]
(iii)	How could you change the zinc/copper cell to have a voltage greater than 1.1 V	
		[1]
(iv)	Complete the labelling of the zinc/tin cell.	[2]
	[Tota	l: 14]

4 The electrolysis of concentrated aqueous sodium chloride, between inert electrodes, is used to make four important chemicals.

hydrogen chlorine sodium hydroxide sodium chlorate(I)

- (a) The ions present in the electrolyte are Na $^+$, H $^+$, C l^- and OH $^-$.
 - (i) Hydrogen ions are discharged at the negative electrode (cathode). Write an equation for this reaction.

[2]

(ii) The hydrogen ions are from the water.

 $H_2O \rightleftharpoons H^+ + OH^-$

Suggest an explanation why the concentration of hydroxide ions increases.

[2]

(iii) When a dilute solution of sodium chloride is used, chlorine is not formed at the positive electrode (anode), a different gas is produced. Name this gas.

......[1]

(iv) State an example of an inert electrode.

.....[1]

(b) (i) State a use of hydrogen.

......[1]


(ii) Why is chlorine used to treat the water supply?

.....[1]

(c) Sodium chlorate(I) is made by the reaction between chlorine and sodium hydroxide. It is used as bleach but over time it decomposes.

$$2NaClO(aq) \rightarrow 2NaCl(aq) + O_2(g)$$

The rate of decomposition can be studied using the apparatus shown below.



(i)	How could you measure the rate of decomposition of sodium chlorate(I)?
	[1]
(ii)	Describe how you could show that the rate of decomposition of sodium chlorate(I) is a photochemical reaction.
	[2]

[Total: 11]

5 Carboxylic acids contain the group

- (a) Ethanoic acid is a typical carboxylic acid. It forms ethanoates.
 - (i) Complete the following equations.

$Mg \; + \; CH_{3COOH} \; \rightarrow \; \; + \;$	
	[2
sodium + ethanoic → + +	

(ii) Ethanoic acid reacts with ethanol to form an ester. Give the name of the ester and draw its structural formula. Show all of the bonds.

name	

structural formula

[2]

[1]

- **(b)** Maleic acid is an unsaturated acid. 5.8 g of this acid contained 2.4 g of carbon, 0.2 g of hydrogen and 3.2 g of oxygen.
 - (i) How do you know that the acid contained only carbon, hydrogen and oxygen?

.....[1]

(ii) Calculate the empirical formula of maleic acid.

Number of moles of carbon atoms =

Number of moles of hydrogen atoms =

Number of moles of oxygen atoms =

The empirical formula is[3]

	· ·	1
(iii)	The mass of one mole of maleic acid is 116 g. What is its molecular formula?	Ex
	[2]	
(iv)	Maleic acid is dibasic. One mole of acid produces two moles of H ⁺ . Deduce its structural formula.	
	ro.	
	121	I

For xaminer's Use

[Total: 13]

For
Examiner's
Use

[2]

6 The Kinetic Theory explains the properties of matter in terms of the arrangement and movement of particles.


- (a) Nitrogen is a gas at room temperature. Nitrogen molecules, N₂, which are spread far apart move in a random manner at high speed.
 - (i) Draw a diagram showing the arrangement of the valency electrons in a nitrogen molecule.

Use \times to represent an electron from a nitrogen atom.

	(ii)	How does the movement and arrangement of the molecules in a crystal of nitrogen differ from those in gaseous nitrogen?
		[3]
(b)	Use	the ideas of the Kinetic Theory to explain the following.
	(i)	A sealed container contains nitrogen gas. The pressure of a gas is due to the molecules of the gas hitting the walls of the container. Explain why the pressure inside the container increases when the temperature is increased.

(ii) The following apparatus can be used to measure the rate of diffusion of a gas.

For Examiner's Use



The following results were obtained.

gas	temperature /°C	rate of diffusion in cm³/min
nitrogen	25	1.00
chlorine	25	0.63
nitrogen	50	1.05

Explain why nitrogen diffuses faster than chlorine.

[2]
Explain why the nitrogen diffuses faster at the higher temperature.

[1]

[Total: 10]

- 7 Synthetic polymers are widely used in the modern world.
 - (a) Their use has brought considerable advantages to modern life as well as some disadvantages.
 - (i) Suggest **two** advantages of a plastic bucket compared to a steel bucket.

.....[2]

(ii) Name two uses of man-made fibres, such as nylon and Terylene.

[2]

(iii) Describe the pollution caused by synthetic polymers.

.....[3]

- **(b)** One type of polymer is formed by addition polymerisation.
 - (i) The structural formula of an addition polymer is given below.

Give the name and structural formula of the monomer.


name of monomer[1]

structural formula of monomer

[1]

(ii) Draw the structural formula of the addition polymer formed by the polymerisation of phenylethene. The structural formula of phenylethene is given below.

For Examiner's Use



[2]

(c) Nylon is made by condensation polymerisation. It has the structural formula shown below.

(i)	Name the	linkage	in	this	poly	mer
-----	----------	---------	----	------	------	-----

E 4.7	
111	ı.
 1.1	1

(ii) Name the natural macromolecules which have the same linkage.

 [1]

(iii) Deduce the formulae of the two monomers which reacted to form the nylon and water.

monomer	
IIIOHOHICH	

monomer

[2]

[Total: 15]

BLANK PAGE

BLANK PAGE

DATA SHEET
The Periodic Table of the Elements

	0	4 He lium	2	20	Ne	Neon 10	40	Ā	Argon 18	84	첫	Krypton 36	131	Xe	Xenon 54		Ru	Radon 86				175	Γn	Lutetium 71		בֿ	Lawrendum 103
	ΙΙΛ			19	ш	Fluorine 9	35.5	CI	Chlorine 17	80		Bromine 35		Ι	lodine 53		Αt	Astatine 85					Υp	Ytterbium 70		8 N	Nobelium 102
				16	0	Oxygen 8	32	S	Sulfur 16	62		_	128	<u>a</u>	Tellurium 52		Ъо	Polonium 84				169	H	Thulium 69		Md	Mendelevium 101
	Λ			14	z	Nitrogen 7			Phosphorus 15						Antimony 51			Bismuth 83					Ē			Fm	Fermium 100
	<u>N</u>			12	ပ				Silicon 14		Ge	Germanium 32	119	Su	Tin 50		Pp	Lead 82				165	운	Holmium 67		Es	Einsteinium 99
	=			7	ш	Boron 5	27	ΝI	Aluminium 13	70	Ga	Gallium 31	115	In	Indium 49	204	11	Thallium 81				162	۵	Dysprosium 66		ర	Californium 98
											Zn	Zinc 30	112	පි	Cadmium 48	201	Нg	Mercury 80				159	<u>Q</u>	Terbium 65		æ	Berkelium 97
										64	చె	Copper 29	108	Ag	Silver 47	197	Αn	Gold 79				157		Gadolinium 64			Curium 96
Group										29	Z	Nickel 28	106	Pd	Palladium 46	195	Ŧ	Platinum 78				152	Ш	Europium 63		Am	Americium 95
Ğ										29	ပိ	Cobalt 27	103	R	Rhodium 45	192	ľ	Iridium 77				150	Sm	Samarium 62			Plutonium 94
		1 Hydrogen	-							26	æ	Iron 26			Ruthenium 44	190	Os	Osmium 76						Promethium 61		ď	Neptunium 93
										55	M	Manganese 25		ပ	n Technetium 43	186	Re	Rhenium 75					Š	Neodymium 60	238	>	Uranium 92
										52	ပ်	Chromium 24	96	Mo	Molybdenum 42	184	>	Tungsten 74				141	Ā	Praseodymium 59		Ра	Protactinium 91
										51	>	Vanadium 23	93	Q Q	Niobium 41	181	Та	Tantalum 73				140	Ce	Cerium 58	232	Ļ	Thorium 90
										48	j	Titanium 22	91	Zr	Zirconium 40	178	Ξ	* Hafnium							mic mass	loqu	mic) number
			ſ							45	လွ	Scandium 21	88	>	Yttrium 39	139	Ľ	Lanthanum 57	227	Actinium	. 68	d corion	r series	2	a = relative atomic mass	X = atomic symbol	b = proton (atomic) number
	=			6	Be	Beryllium 4	24	Mg	Magnesium 12	40	Sa	Calcium 20	88	જ	Strontium 38	137	Ba	Barium 56	226	226 Rad ium Radium			30-7 1 Eartinaidus seire 190-103 Artinoid series		а	× ×	
	_			7	=	Uthium 3	23	Na	Sodium 11	39	¥	Potassium 19	82	Rb	Rubidium 37	133	S	Caesium 55	ı	Francium	87	*58-71 Lanthanoid series	190-103	2		Key	Q

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.