

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

CANDIDATE NAME						
CENTRE NUMBER			CANDIDA NUMBER			

CHEMISTRY 0620/51

Paper 5 Practical Test

October/November 2010

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

For Examiner's Use	
1	
2	
Total	

This document consists of 8 printed pages.



1 You are going to investigate what happens when two different solids, **A** and **B**, dissolve in water.

Read all instructions below carefully before starting the experiments.

#### Instructions

You are going to carry out two sets of experiments.

## (a) Experiment 1

Using a measuring cylinder, pour 20 cm<sup>3</sup> of distilled water into the polystyrene cup provided. Put the cup into a 250 cm<sup>3</sup> beaker for support. Measure the temperature of the water and record it in the table below.

Add 2 g of solid **A** provided to the cup and stir the mixture with a thermometer. Measure and record the temperature of the solution after one minute. Pour the solution away and rinse the polystyrene cup.

Repeat the experiment using 3g of the solid **A** provided. Record your results in the table. Repeat the experiment using 4g of the solid **A** provided. Record your results in the table. Repeat the experiment using 6g of the solid **A** provided. Record your results in the table.

mass of solid A/g	initial temperature/°C	final temperature/°C
2		
3		
4		
6		

[3]

### (b) Experiment 2

Repeat experiment 1 using 2 g, 3 g and 4 g of solid **B** respectively. Record your results in the table below.


mass of solid <b>B</b> /g	initial temperature/°C	final temperature/°C
2		
3		
4		

[2]

For Examiner's Use

(c) Plot the results of the experiments on the grid below. Draw two best-fit straight line graphs.

Clearly label your graphs.



[6]

(d)	(i)	<b>Use your graph</b> to estimate the temperature of the reaction mixture if 6 g of solid <b>B</b> was added to 20 cm <sup>3</sup> of water. Show clearly <b>on the grid</b> how you worked out your answer.
		[2]
	(ii)	<b>From your graph</b> , work out the temperature of the reaction mixture if 5 g of solid <b>A</b> was added to 20 cm³ of water.  Show clearly <b>on the graph</b> how you worked out your answer.
		[2]
(e)	Wh	at type of chemical reaction occurs when solid <b>A</b> dissolves in water?
		[1]
(f)		plain how the temperature changes would differ in the experiments if 40 cm <sup>3</sup> of water s used.
		[2]
(g)	Pre	edict the effect of using lumps of solid <b>B</b> in Experiment 2. Explain your answer.
		[2]
(h)	-	ggest <b>one</b> change you could make to the <b>apparatus</b> used in the experiments to obtain re accurate results.
		[1]
		[Total: 21]

2 You are provided with a mixture of two solids, **C** and **D**. Solid **C** is water-soluble and **D** is insoluble. Carry out the following tests on **C** and **D**, recording all of your observations in the table.

Conclusions must **not** be written in the table.

tests		observations
the tube	d 15 cm <sup>3</sup> of distilled water to the mixture in boiling tube. Stopper and shake the boiling e for two minutes. Filter the contents of the e, keeping the filtrate and the residue for following tests.	
test	on the filtrate	
(a)	To about 1 cm³ of the solution, add a few drops of dilute nitric acid and about 1 cm³ of aqueous potassium iodide.	[2]
(b)	To about 1 cm <sup>3</sup> of the solution add about 1 cm <sup>3</sup> of dilute hydrochloric acid.	[1]
(c)	To about 1 cm <sup>3</sup> of the solution add an equal volume of aqueous sodium hydroxide. Now add a small spatula measure of aluminium powder and warm the mixture <b>carefully</b> . Test any gases given off.	[2]

		tests	observations
tests on the residue			
Wash the residue in the filter paper with a little distilled water.		• • •	
Using a spatula, transfer some of the solid residue from the filter paper into two test-tubes.		sidue from the filter paper into two	
(d) Heat the solid in the first test-tube gently and then strongly. Leave the test-tube to cool.		then strongly. Leave the test-tube to	[2]
(e) Add about 2 cm³ of dilute hydrochloric acid to the second test-tube. Test the gas given off with limewater.		e second test-tube. Test the gas given	[2]
(f) After 2 minutes, add an equal volume of distilled water and shake the test-tube. Decant off the liquid and divide into two approximately equal portions.		lled water and shake the test-tube. ant off the liquid and divide into two	
	(i)	To the first portion add aqueous sodium hydroxide a little at a time until in excess.	[2]
	(ii)	To the second portion add aqueous ammonia a little at a time until in excess.	[2]

(g)	Identify the gas given off in test (c).
	[1]
(h)	Identify solid C.
	[2]
(i)	What conclusions can you draw about solid <b>D</b> ?
	[3]
	[Total: 19]

## NOTES FOR USE IN QUALITATIVE ANALYSIS

#### **Test for anions**

anion	test	test result
carbonate (CO <sub>3</sub> <sup>2-</sup> )	add dilute acid	effervescence, carbon dioxide produced
chloride (Cl <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	white ppt.
iodide (I <sup>-</sup> ) [in solution]	acidify with dilute nitric acid, then add aqueous silver nitrate	yellow ppt.
nitrate (NO <sub>3</sub> <sup>-</sup> ) [in solution]	add aqueous sodium hydroxide then aluminium foil; warm carefully	ammonia produced
sulfate (SO <sub>4</sub> <sup>2-)</sup> [in solution]	acidify with dilute nitric acid, then aqueous barium nitrate	white ppt.

# Test for aqueous cations

cation	effect of aqueous sodium hydroxide	effect of aqueous ammonia
aluminium (Al³+)	white ppt., soluble in excess giving a colourless solution	white ppt., insoluble in excess
ammonium (NH <sub>4</sub> +)	ammonia produced on warming	_
calcium (Ca <sup>2+</sup> )	white ppt., insoluble in excess	no ppt., or very slight white ppt.
copper (Cu <sup>2+</sup> )	light blue ppt., insoluble in excess	light blue ppt., soluble in excess giving a dark blue solution
iron(II) (Fe <sup>2+</sup> )	green ppt., insoluble in excess	green ppt., insoluble in excess
iron(III) (Fe <sup>3+</sup> )	red-brown ppt., insoluble in excess	red-brown ppt., insoluble in excess
zinc (Zn <sup>2+</sup> )	white ppt., soluble in excess giving a colourless solution	white ppt., soluble in excess giving a colourless solution

# **Test for gases**

gas	test and test results	
ammonia (NH <sub>3</sub> )	turns damp red litmus paper blue	
carbon dioxide (CO <sub>2</sub> )	turns limewater milky	
chlorine (Cl <sub>2</sub> )	bleaches damp litmus paper	
hydrogen (H <sub>2</sub> )	'pops' with a lighted splint	
oxygen (O <sub>2</sub> )	relights a glowing splint	

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.