

# UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

| CANDIDATE<br>NAME |  |                     |  |  |
|-------------------|--|---------------------|--|--|
| CENTRE<br>NUMBER  |  | CANDIDATE<br>NUMBER |  |  |

CHEMISTRY 0620/53

Paper 5 Practical Test

October/November 2011

1 hour 15 minutes

Candidates answer on the Question Paper.

Additional Materials: As listed in the Confidential Instructions

#### **READ THESE INSTRUCTIONS FIRST**

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

You may use a pencil for any diagrams, graphs or rough working.

Do not use staples, paper clips, highlighters, glue or correction fluid.

DO NOT WRITE IN ANY BARCODES.

Answer all questions.

Practical notes are provided on page 8.

At the end of the examination, fasten all your work securely together.

The number of marks is given in brackets [ ] at the end of each question or part question.

| For Examiner's Use |  |
|--------------------|--|
| 1                  |  |
| 2                  |  |
| Total              |  |

This document consists of 6 printed pages and 2 blank pages.



1 You are going to investigate the reaction between dilute sulfuric acid and three aqueous solutions of sodium hydroxide of different concentrations, labelled **A**, **B** and **C**.

Read all the instructions below carefully before starting the experiments.

#### Instructions

You are going to carry out three experiments.

## (a) Experiment 1

Fill the burette with the dilute sulfuric acid provided to the 0.0 cm<sup>3</sup> mark.

Use a measuring cylinder to pour 20 cm³ of solution **A** into a conical flask. Add a few drops of phenolphthalein indicator to the flask.

Add the sulfuric acid from the burette 1 cm<sup>3</sup> at a time, while shaking the flask, until the colour of the phenolphthalein changes. Record the burette readings in the table.

## (b) Experiment 2

Fill the burette with dilute sulfuric acid to the 0.0 cm<sup>3</sup> mark.

Empty the conical flask and rinse it with water. Use a measuring cylinder to pour 20 cm<sup>3</sup> of solution **B** into the conical flask. Add a few drops of phenolphthalein to the flask. Add the sulfuric acid from the burette 1 cm<sup>3</sup> at a time, while shaking the flask, until the colour of the phenolphthalein changes. Record the burette readings in the table.

## (c) Experiment 3

Repeat Experiment 2, using solution **C** instead of solution **B**. Record your burette readings in the table and complete the table.

|                                 | experiment 1 | experiment 2 | experiment 3 |
|---------------------------------|--------------|--------------|--------------|
| final reading/cm <sup>3</sup>   |              |              |              |
| initial reading/cm <sup>3</sup> |              |              |              |
| difference/cm <sup>3</sup>      |              |              |              |

| г | $\sim$ 1 |
|---|----------|
| ı | nı       |
| ı | vi       |

| (d) | What colour change was observed after the sulfuric acid was added to the flask?        |     |
|-----|--|-----|
|     | from to  | [2] |
| (e) | What type of chemical reaction occurs when sulfuric acid reacts with sodium hydroxide? |     |
|     |  | [1] |

| (†) | (1)  | Complete the sentences below.   |  |  |
|-----|------|---|--|--|
|     |      | Aqueous sodium hydroxide labelled needed the smallest volume of sulfuric acid to change the colour of the phenolphthalein.                          |  |  |
|     |      | Aqueous sodium hydroxide labelled needed the largest volume of sulfuric acid to change the colour of the phenolphthalein. [1]                       |  |  |
|     | (ii) | The order of concentration of the solutions of sodium hydroxide is  |  |  |
|     |      | least concentrated  |  |  |
|     |      | <b>\</b>  |  |  |
|     |      | most concentrated   |  |  |
| (g) | Cor  | mpare the volumes of sulfuric acid used in Experiments 1 and 2.   |  |  |
|     |      | [1]   |  |  |
| (h) |      | experiment 3 was repeated using 40 cm <sup>3</sup> of solution <b>C</b> , what volume of sulfuric acid ald be used?                                 |  |  |
|     |      | [2]   |  |  |
| (i) |      | What would be a more accurate method of measuring the volume of the aqueous sodium hydroxide?   |  |  |
|     |      | [1]   |  |  |
| (j) |      | at would be the effect on the results if the solutions of sodium hydroxide were warmed ore adding the sulfuric acid? Give a reason for your answer. |  |  |
|     | effe | ect on results  |  |  |
|     | rea  | son[2]  |  |  |
| (k) |      | ggest a different method of finding the order of concentrations of the solutions of lium hydroxide.   |  |  |
|     |      |   |  |  |
|     |      |   |  |  |
|     |      |   |  |  |
|     |      |   |  |  |
|     |      | [3]   |  |  |
|     |      | [Total: 21]   |  |  |

You are provided with two different salts, D and E.
 D is an aqueous solution of the salt and E is a solid.
 Carry out the following tests on each salt, recording all of your observations in the table.
 Conclusions must not be written in the table.

|                         | tests   | observations |
|-------------------------|---|--------------|
| (a) De                  | escribe the appearance of   |              |
| (i)                     | solution <b>D</b> ,   | [1]          |
| (ii)                    | solid <b>E</b> .  | [1]          |
| tests o                 | n solution <b>D</b>   |              |
| in                      | vide the solution into five equal portions test-tubes, and carry out the following sts.                                 |              |
| (i)                     | Add about 1 cm <sup>3</sup> of dilute nitric acid to the first portion of the solution and then aqueous barium nitrate. | [1]          |
| (ii)                    | To the second portion of the solution, add about 1 cm <sup>3</sup> of dilute nitric acid and aqueous silver nitrate.    | [2]          |
| (iii)                   | To the third portion of the solution, add an excess of aqueous sodium hydroxide.  | [2]          |
| (iv)                    | Add an excess of aqueous ammonia solution to the fourth portion.  | [1]          |
|                         | Keep the remaining portion of the solution for use in test (c)(ii).   |              |
| tests on solid <b>E</b> |   |              |
| (c) (i)                 | Place about half of solid <b>E</b> in a test-tube. Heat the test-tube gently, then strongly. Test any gas given off.    | [3]          |
|                         | Leave the test-tube to cool for five minutes. Then add about 1 cm³ of dilute nitric acid to the test-tube.              | [1]          |
| (ii)                    | Add the rest of solid <b>E</b> to the remaining portion of solution <b>D</b> in a test-tube.                            | [1]          |

| (d) | Identify salt <b>D</b> .                             |
|-----|--|
|     | [3]  |
| (e) | Identify the gas given off in test (c)(i).           |
|     | [1]  |
| (f) | What conclusions can you draw about solid <b>E</b> ? |
|     |  |
|     | [2]  |
|     | [Total: 19]  |

# **BLANK PAGE**

# **BLANK PAGE**

## NOTES FOR USE IN QUALITATIVE ANALYSIS

### **Test for anions**

| anion  | test   | test result                            |
|--|--|--|
| carbonate (CO <sub>3</sub> <sup>2-</sup> )               | add dilute acid  | effervescence, carbon dioxide produced |
| chloride (Cl <sup>-</sup> ) [in solution]                | acidify with dilute nitric acid, then add aqueous silver nitrate | white ppt.                             |
| iodide (I <sup>-</sup> )<br>[in solution]                | acidify with dilute nitric acid, then add aqueous silver nitrate | yellow ppt.                            |
| nitrate (NO <sub>3</sub> <sup>-</sup> )<br>[in solution] | add aqueous sodium hydroxide then aluminium foil; warm carefully | ammonia produced                       |
| sulfate (SO <sub>4</sub> <sup>2-)</sup><br>[in solution] | acidify with dilute nitric acid, then aqueous barium nitrate     | white ppt.                             |

## Test for aqueous cations

| cation                        | effect of aqueous sodium hydroxide                         | effect of aqueous ammonia                                      |
|-------------------------------|--|--|
| aluminium (Al³+)              | white ppt., soluble in excess giving a colourless solution | white ppt., insoluble in excess                                |
| ammonium (NH <sub>4</sub> +)  | ammonia produced on warming                                | _  |
| calcium (Ca <sup>2+</sup> )   | white ppt., insoluble in excess                            | no ppt., or very slight white ppt.                             |
| copper (Cu <sup>2+</sup> )    | light blue ppt., insoluble in excess                       | light blue ppt., soluble in excess giving a dark blue solution |
| iron(II) (Fe <sup>2+</sup> )  | green ppt., insoluble in excess                            | green ppt., insoluble in excess                                |
| iron(III) (Fe <sup>3+</sup> ) | red-brown ppt., insoluble in excess                        | red-brown ppt., insoluble in excess                            |
| zinc (Zn <sup>2+</sup> )      | white ppt., soluble in excess giving a colourless solution | white ppt., soluble in excess giving a colourless solution     |

## **Test for gases**

| gas                               | test and test results            |  |
|-----------------------------------|----------------------------------|--|
| ammonia (NH <sub>3</sub> )        | turns damp red litmus paper blue |  |
| carbon dioxide (CO <sub>2</sub> ) | turns limewater milky            |  |
| chlorine (Cl <sub>2</sub> )       | bleaches damp litmus paper       |  |
| hydrogen (H <sub>2</sub> )        | 'pops' with a lighted splint     |  |
| oxygen (O <sub>2</sub> )          | relights a glowing splint        |  |

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

www.theallpapers.com