

UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS International General Certificate of Secondary Education

	CANDIDATE NAME					
	CENTRE NUMBER	CANDIDATE NUMBER				
*						
	CAMBRIDGE IN	0607/41				
	Paper 4 (Extend	May/June 2013				
			2 hours 15 minutes			
4 2	Candidates ans					
3 3 2 *	Additional Mater	rials: Geometrical Instruments Graphics Calculator				

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.

Write in dark blue or black pen.

Do not use staples, paper clips, highlighters, glue or correction fluid.

You may use a pencil for any diagrams or graphs.

DO NOT WRITE IN ANY BARCODES.

Answer all the questions.

Unless instructed otherwise, give your answers exactly or correct to three significant figures as appropriate. Answers in degrees should be given to one decimal place.

For π , use your calculator value.

You must show all the relevant working to gain full marks and you will be given marks for correct methods, including sketches, even if your answer is incorrect.

The number of marks is given in brackets [] at the end of each question or part question. The total number of marks for this paper is 120.

For Examiner's Use				


This document consists of **16** printed pages.




UNIVERSITY of CAMBRIDGE International Examinations

Formula List

For the equation	$ax^2 + bx + c = 0$	$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$
Curved surface area, A, of cylin	nder of radius r, height h.	$A = 2\pi rh$
Curved surface area, A, of cone	e of radius <i>r</i> , sloping edge <i>l</i> .	$A = \pi r l$
Curved surface area, A, of sphe	ere of radius <i>r</i> .	$A = 4\pi r^2$
Volume, V , of pyramid, base as	rea A, height h.	$V = \frac{1}{3}Ah$
Volume, V , of cylinder of radiu	is r , height h .	$V = \pi r^2 h$
Volume, V , of cone of radius r ,	height <i>h</i> .	$V = \frac{1}{3}\pi r^2 h$
Volume, V , of sphere of radius	r.	$V = \frac{4}{3}\pi r^3$
\bigwedge^{A}		$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$
		$a^2 = b^2 + c^2 - 2bc \cos A$
		Area = $\frac{1}{2}bc\sin A$
в <u></u> а	\longrightarrow_{C}	




www.theallpapers.com




© UCLES 2013








The diagram shows a vertical radio mast *PQR* supported by 6 straight wires. *A*, *B*, *C* and *P* are on level horizontal ground. RA = RB = RC and QA = QB = QC. PQ = 30 m, QR = 20 m and angle AQP = angle BQP = angle $CQP = 65^{\circ}$.




(a) Show that QC = 70.99 m, correct to 2 decimal places.








0607/41/M/J/13



© UCLES 2013

www.theallpapers.com



(b) The speeds of another 140 cars were measured on road B. The results are shown in this table.

For Examiner's Use

[Turn over www.theallpapers.com

Right-handed

Total

Left-handed


			Lett-nanueu	Right-handed	Total	
		Girls	4	14	18	
		Boys	3	11	14	
		Total	7	25	32	
(a)		e chosen at rando ility that they are		class.		
(b)		are chosen at ran ility that exactly o	dom.			[2
(c)	Two of the right	-handed students				[
	Find the probabi	ility that at least o	one is a girl.			

8 The table shows the number of left-handed and right-handed girls and boys in a class.

For Examiner's Use

The resistance, R ohms, of a standard length of wire varies inversely as the square of its diameter, d mm.				
(a)	The	e resistance of a standard length of wire of diameter 0.5mm is 0.8 ohms.		
	(i)	Find a formula for <i>R</i> in terms of <i>d</i> .		
		Answer(a)(i) $R =$	[3]	
	(ii)	Find the resistance of a standard length of the same type of wire with diameter 2 mm.		
	(iii)	<i>Answer(a)</i> (ii) ohms The resistance of a standard length of the same type of wire is 4 ohms. Find the diameter of this wire.	5 [1]	
(b)		<i>Answer(a)</i> (iii) mm a different type of wire the resistance of a standard length is 2 ohms. d the resistance of a standard length of this wire when the diameter is doubled.	[2]	
		Answer(b) ohms	5 [2]	





(a) On the diagram, sketch the graph of y = f(x), where

$$f(x) = \frac{(x-1)}{(x+3)}$$
 between $x = -6$ and $x = 3$. [3]

(b) Find the co-ordinates of the point where the graph crosses the *x*-axis.

$$Answer(b) \quad (\dots, , \dots,) \quad [1]$$
(c) Find the equations of the asymptotes of $y = f(x)$.

$$Answer(c) \quad \dots \quad and \quad \dots \quad [2]$$
(d) Find the range of $f(x)$ for $x \ge 0$.

$$Answer(d) \quad \dots \quad [2]$$
(e) Find the solutions to the equation $\frac{(x-1)}{(x+3)} = -5 - 2x$.

$$Answer(e) \ x = \dots \quad \text{or } x = \dots \quad [3]$$
(f) On the diagram, sketch the graph of $y = f(x-3)$.


10

www.theallpapers.com

ForExaminer's

Use

y



Question 12 is printed on the next page.

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.