## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

**International General Certificate of Secondary Education** 

## MARK SCHEME for the May/June 2011 question paper for the guidance of teachers

## 0581 MATHEMATICS

0581/21

Paper 2 (Extended), maximum raw mark 70

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2011	0581	21

## **Abbreviations**

cao correct answer only cso correct solution only

dep dependent

ft follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

www without wrong working

Qu.	Answers	Mark	Part Mark
1	847	1	
2	correct regions shaded	1, 1	
3	48	2	B1 for 3 and 16 seen
4	<b>(a)</b> 10	1	
	<b>(b)</b> 5.5 oe	1	
5	(a) 86400	1	
	<b>(b)</b> $8.64 \times 10^4$	1ft	
6	108	2	M1 for $3^3$ or $27$ or $\left(\frac{1}{3}\right)^3$ or $\frac{1}{27}$ seen
7	13	3	<b>B1</b> for 12, 5 seen <b>M1</b> for (their 12) <sup>2</sup> + (their 5) <sup>2</sup> or <b>M2</b> $\sqrt{[(-8-4)^2+(1-6)^2]}$ oe or <b>M1</b> if $\sqrt{\text{missing}}$
8	6.70	3	M1 for $(r^3 = )$ 1260 $\times \frac{3}{4\pi}$ oe seen M1 for $\sqrt[3]{}$ of their $r^3$ seen or implied
9	22.5 oe	3	<b>B2</b> $180 = 5x + 2x + x$ oe or better <b>B1</b> for $2x$ or $6x$ marked in the correct place on the diagram.
10	x = 13 $y = -9$	3	M1 for consistent multiplication and addition/subtraction A1 for $x = 13$ or A1 for $y = -9$
11	(a) 85.8	2	M1 for 23.25 and 19.65 seen
	<b>(b)</b> 456.8625 cao	1	
12	(a) (0)8(.)01 (am)	1	Not 8.01pm
	<b>(b)</b> 78.4 or 78.38 to 78.39	3	<b>M2</b> for 827 ÷ 10.55 or <b>M1</b> for figs 827 ÷ their time
13	(a) 0.54	2	<b>M1</b> for $\frac{2.7 \times 20000}{100000}$ oe
			or SC1 for figs 54 in answer
	<b>(b)</b> 1.61	2	<b>SC1</b> for figs 161 or <b>M1</b> 200 <sup>2</sup> or 20 000 <sup>2</sup> seen

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	IGCSE – May/June 2011	0581	21

	T	I	
14	-2.64, 1.14 cao with working	4	<b>B1</b> for $\sqrt{3^2 - 4(2)(-6)}$ or better seen anywhere <b>B1</b> for $p = -3$ and $r = 2 \times 2$ or better as long as in the form $\frac{p + \sqrt{q}}{r}$ or $\frac{p - \sqrt{q}}{r}$
			After <b>B0B0</b> , <b>SC1</b> for –2.6 or –2.637(45) and 1.1 or 1.137(45)
15	(a) 4	1	
	<b>(b) (i)</b> $\frac{12}{36}$ oe 0.333	1	
	(ii) $\frac{11}{36}$ , 0.306 or 0.3055 to	1	
	0.3056		
		1	
	(c) $\frac{8}{15}$ oe 0.533(3)		
16	(a) Answer given	2	$\mathbf{M1} \ (A =)k^2 - \pi \left(\frac{k}{2}\right)^2$
			$\mathbf{E1} \ A = k^2 - \frac{\pi k^2}{4}$
			correctly completed to $4A = 4k^2 - \pi k^2$
	<b>(b)</b> $k = (\pm) \sqrt{\frac{4A}{(4-\pi)}}$ or $2\sqrt{\frac{A}{(4-\pi)}}$	3	<b>M1</b> factorising (must contain a $\pi$ )
	$\bigvee (4-\pi) \qquad \bigvee (4-\pi)$		M1 division (by coefficient of $k^2$ )
			M1 square root
17	(a) 66°	2	<b>M1</b> for 90° clearly identified as $A$
	<b>(b)</b> 33°	1	
	(c) 123°	2	<b>B1</b> for $OBA$ or $OAB = 57^{\circ}$
18	(a) (i) $-r + q$ or $q - r$	1	
	(ii) $\frac{1}{2}(3\mathbf{q} - \mathbf{r})$ oe	1	Must be simplified
	(b) correct working	3	M1 for $MX = \frac{1}{2} r + \frac{3}{4}$ their $(-r + q)$
	(1)		M1 using a different route for $XS$ or $\frac{1}{2}MS$
			E1 dep correct simplification and conclusion
19	(a) 480	1	
	<b>(b)</b> 9900	3	M1 for attempt at area under graph M1 for $0.5 \times 15 \times$ (their (a) + $14 \times 60$ ) oe
			or $0.5 \times 15 \times (8 + 14)$ oe
	(c) $0.125 \text{ or } \frac{1}{8}$	2	M1 for numerical vertical/horizontal or numerical use of $v = u + at$ but $t \le 120$ or $t \le 2$
20	(a) (i) 9 (ii) $8x^3$ cao	1 1	
	(b) 4 www	3	<b>M1</b> for $(2x-3)^3 = 125$ <b>M1</b> $2x-3=5$
	(c) $\frac{x+3}{2}$	2	<b>M1</b> for $x \pm 3 = 2y$ or $x = \frac{y \pm 3}{2}$
	İ	1	I