## UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS

**International General Certificate of Secondary Education** 

## MARK SCHEME for the October/November 2008 question paper

## 0580 and 0581 MATHEMATICS

**0580/04 and 0581/04** Paper 04 (Extended), maximum raw mark 130

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2008 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

## **Abbreviations**

cao correct answer only cso correct solution only

dep dependent

ft follow through after error isw ignore subsequent working

oe or equivalent SC Special Case

www without wrong working

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

1 (a) (i)	(\$) 6 000 cao	B2	<b>M1</b> for $0.1 \times 10\ 000 + 0.25 \times 20\ 000$ oe
(ii)	15 (%) cao	B2	M1 for $\frac{their(a)(i)}{40000} \times 100$
(b)	(\$) 11 200 ft	B1 ft	ft 17200 – their (a)(i)
(c) (i)	(\$) 7500 cao	B2	M1 for $\frac{12000}{5+3} \times 5$ oe After M0, SC1 for 4500
(ii)	9/80 cao	B1	Ignore decimals or %'s seen Mark final fraction
(d)	(\$) 8640 cao	B2	<b>M1</b> for 10 800 ÷ 1.25 oe [10]

		,	
2 (a) (i)	x(x+4) / 2 = 48 oe $x^2 + 4x - 96 = 0$	M1	Eqn must include 48
	$x^2 + 4x - 96 = 0$	E1	Dep on M1 + shows one intermediate algebraic step with no errors seen
(ii)	- 12 or 8	B1B1	Allow deletion of negative root
(iii)	12 (cm) correct or ft	B1ft	Accept 12 or ft their positive root in part (ii) (if only one) + 4.
(b)	$\frac{4}{5}$ oe	B2	M1 for $\frac{x}{x+4} = \frac{1}{6}$ oe
(c) (i)	$(x + 4)^2 + x^2 = 9^2$ oe or $x^2 + 8x + 16 + x^2 = 81$ $2x^2 + 8x - 65 = 0$	M1 E1	Accept $2^{nd}$ line for M1 or $2x^2 + 8x + 16 = 81$ Dep on M1 with no errors, expanded brackets step needed
(ii)	$\frac{p+(-)\sqrt{q}}{r}$ where $p = -8$ and $r = 2 \times 2$ and $q = 8^2 - 4(2)(-65)$ oe (584)	M1 M1	Allow second mark if in form $p \pm \frac{\sqrt{q}}{r}$
	- 8.04, 4.04 cao www	A1A1	SC2 if correct solutions but no working shown or SC1 for -8.041522987 and 4.041522987 rounded or truncated
(iii)	21.08 or 21.1 (cm) strict ft	B1ft dep	ft 4.04 in part (ii) or 2 × a positive root + 13  [14]

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

3	(a)	5.(04), 0(.0), 8.7 or 8.66(6) or better	В3	1 each
		seen		
	<b>(b)</b>	Correct axes for domain and range	<b>S1</b>	
		10 correct points, on correct grid line or	P3ft	<b>P2ft</b> for 8 or 9 correct
		within correct 2mm square vertically		P1ft for 6 or 7 correct
		Reasonable curve through 10 points	C1ft	Correct shape, not ruled, within 1 mm of points
		condone curvature around $x = -0.2$ and 0.2		(curves could be joined)
		Two separate branches	B1ft	Independent but needs two 'curves' on either
		The state of the s		side of y-axis
	(c) (i)	y = -3x ruled correctly	L1	Check at $(-1, 3)$ to $(1, -3)$ within 1 mm (can be
	(•) (-)	y en raisea correcting		shorter)
		-2.95 to $-2.6$ , $-0.75$ to $-0.6$ , $0.5$ to $0.6$	B2	<b>B1</b> for 2 correct.
		2.55 to 2.0, 0.75 to 0.0, 0.5 to 0.0	22	isw $y$ – values
				No penalty for each extra value if curve is cut
				more than 3 times
				more than 5 times
	(ii)	(r-)2 $(k-)1$	D1D1	After 0,0 <b>SC1</b> for $x^3 + 3x^2 - 1 = 0$
	(11)	(a =) 3   (b =) -1	B1B1	After 0,0 SC1 for $x + 5x - 1 = 0$
	(A)	T	TP1	Must be a generally top sout allow allow
	<b>(d)</b>	Tangent to their curve ruled at $x = -2$	<b>T1</b>	Must be a reasonable tangent allow slight
		. ,	3.54	daylight <1mm
		rise/run using correct scales	M1	Dep on T1 (implied by answer 3 to 4.5)
		4.5.		Must show working if answer out of range
		−4.5 to −3	<b>A1</b>	
				[17]

4 (a)	72	B1	
(b) (i)	$0.5 \times 15 \times 15 \sin (their 72)$ oe 106.9 to 107 (cm <sup>2</sup> ) cso	M1 A1	not for 90° www2
(ii)	534.5 to 535 (cm <sup>2</sup> ) ft	B1 ft	ft their (i) × 5
(iii)	$\pi \times 15^2 \times 50$ their (ii) × 50 Vol of cylinder – prism 8590 - 8625 (cm <sup>3</sup> ) cao	M1 M1 M1 A1	(707 or 35350) or $\pi \times 15^2$ (26750) or $\pi \times 15^2$ - their (b) (ii) Dep on M2 then $\times 50$ www4
(c)	$(AB =) 15\sin(their36) \times 2$ oe (17.63) (not 30° or 45°)	M1	or $\sqrt{15^2 + 15^2 - 2 \times 15 \times 15 \times \cos(their72)}$ Not for 90° or 60° or sine rule
	Area of one rectangle = their $AB \times 50$ 5 (50 × a length) + 2 × their (b)(ii) 5470 - 5480 (cm <sup>2</sup> ) cao	M1 M1 A1	dep on $1^{\text{st}}\mathbf{M}$ (881.5) not $15 \times 50$ Indep (4407.5 + 1070) www4

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

5	(a)	(60 + 40)/35	M1	(2.857) could be in parts
		Correct method to convert a decimal time to minutes	M1	ft a decimal
		to minutes		either full answer or decimal part × 60 (e.g. 51.(428), 171.(4 )or 2hrs 51 or 51 m)
		14 46 or 2 46 pm cao	A1	(e.g. 31.(428), 171.(4)01 21118 31 01 31 111)
		of the second second		
	(b) (i)	260	B1	
	(ii)	145	B1ft	ft their <b>(b) (i)</b> – 115
	(c)	$(AC^2 = )40^2 + 60^2 - 2 \times 40 \times 60 \times \cos 115$	M2	M1 for correct implicit version
		(AC=) $$ of a correct combination	M1	dependent (7229)
		85(.0 km) cao	<b>A1</b>	www4
	(d)	$\frac{\sin A}{\sin A} = \frac{\sin 115}{\sin A}$ oe	M1	Implicit equation
		$\frac{-}{60} = \frac{1}{their(c)}$ de		Could use cosine rule M1 for implicit
		, ,		and M1 for explicit form
		$\sin 1.0$ $\sin 1.0$	M1	Dep on M1 Explicit equation
		$(\sin A =) \frac{\sin 115}{their(c)} \times 60$		- or one and
		39.76 to 39.8 cao	<b>A1</b>	www3
	(e)	40sin80 + 60sin35 oe	M2	their (c) $\times \sin(100 - their$ (d))
	(C)	(39.4) (34.4)	1412	or their (c) $\times$ cos (their (d) – 10)
		()		M1 for either 40sin80 or 60sin35
				or implicit trig version using their (c)
		73.76 – 73.81 (km) cao	<b>A1</b>	www3
				[15]

6	(a) (i)	30	B1	
	(ii)	30, 30.5, 31	B1 B1	Penalty 1 for each extra value
	( )	,	<b>B</b> 1	Ignore repeated values
	(iii)	$\frac{10 \times 30 + 7 \times 31 + x \times 32}{10 \times 30 \times 30 \times 30} = 30.65$	3.71	
		$\frac{10+7+x}{10+3}$	M1	
		correct clearance of fraction	М1	Den en M1
		correct creatance of maction	M1	Dep on M1
		3 cao	A 1	e.g. $517 + 32x = 521.05 + 30.65x$ oe www3
		5 640	A1	www3
	(b) (i)	$35 \times 15 + 115 \times 21 + 26 \times 23 + 24 \times 27$	<b>M3</b>	(4186/200) <b>M1</b> for use of 15, 21, 23, 27 (allow
		200		one error)
		_00		and <b>M1</b> for use of $\sum fx$ with value of x in
				correct range used (allow one further error)
				and M1 dep on $2^{nd}$ M for dividing by $\sum f$ or
				_
			۸.1	200
		20.93 or 20.9 cao	A1	www4 Accept 21 after M3 earned
	(ii)	2.6 cao	B1	
	( )			
		0.7 and 0.8	<b>B4</b>	B3 for one correct
				or <b>B2</b> for 3.5 and 4 seen
				or <b>B1</b> for 4 seen
				[16]

Page 6	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

7 (a) (i)	Translation only  (0) oe	B1 B1	Throughout parts (i) to (v) if more than one transformation is given then no marks at all for that part Accept T
(ii)	Reflection only $x = 1$ oe only	B1 B1	Accept M
(iii)	Reflection only $y = -x$ oe only	B1 B1	Accept M
(iv)	Enlargement only (centre)(2, 0), only (scale factor) 0.5 oe only	B1 B1 B1	Accept E
(v)	Stretch only (factor) 2, only x-axis oe invariant cao only	B1 B1 B1	Accept S Ignore parallel to <i>y</i> -axis
(b) (i)	$\begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}$	B2	B1 each column
(ii)	$\begin{pmatrix} 1 & 0 \\ 0 & 2 \end{pmatrix}$	B2	B1 for right hand column [16]

8 (a)	<i>x</i> = 78	<b>B</b> 1	
	alternate angles	R1	Dep on <b>B1</b> Accept Z <u>angle</u> , extras can spoil Accept longer reasons using correct language and clarity with angles used. e.g. allied angles gives 102° and angles on a straight line = 180°
	either $y = 144$ or $z = 102$ (opposite angles of) cyclic quad (= 180)	B1 R1	Dep on <b>B1</b> , extras can spoil
	and $z = 102$ or $y = 144$ Angles (in (a)) quadrilateral (= 360) or (opp angles of) cyclic quad (= 180)	B1 R1	Dep on <b>B1</b> extras can spoil
(b)	Their $z + 36 \neq 180$ oe	R1	Could also use their angles $x$ and $y$ provided $x + y \neq 180$ . Could be a longer reason involving angles must be clearly explained.
(c)	72 or 288	B1	
(d)	51 cao	B1	[9]

Page 7	Mark Scheme	Syllabus	Paper
	IGCSE – October/November 2008	0580/0581	04

9 (a)	(p=) 5 cao,	B1	Accept in correct order if no labels
	(q =) 12  cao (r =) 1  ft	B1 B1ft	ft for $r = 18 - their p - their q$ provided $r$ not
	(/ –) 1 11	Dill	negative $p = men \ q \text{ provided } r \text{ not}$
			inegative
(b) (i)	17 cao	B1	
(ii)	12 cao	B1	
(c) (i)	26 cao	B1	
(ii)	57 ft	B1ft	ft $45 + their q$
(11)		Din	it is then q
(d) (i)	8	B1	
	$\frac{3}{100}$ oe isw		
(ii)	45 oe isw	B1	
	$\frac{1}{100}$ de isw		
(e)	Any fraction with denominator 74 seen	B1	
	$\frac{37}{3} \times \frac{36}{3}$	M1	ft their fraction i.e. one taken off each part
	$\frac{74}{74} \times \frac{73}{73}$		$\frac{k}{l} \times \frac{k-1}{l-1}$ N.B $\frac{1}{2} \times \frac{36}{73}$ gets <b>B1M1</b>
	18		
	$\frac{18}{73}$ oe isw cao	A1	$\frac{1332}{5402}$ www3 (if decimal then 0.247 or better)
	13		Do not accept ratio or in words
			[12]
40 ( ) (*)	1 - 4 - 11	1	
10 (a) (i)	$\frac{8 \times (8+1)}{3} = 36$	E1	
	)		

10 (a) (i)	$\frac{8 \times (8+1)}{2} = 36$ $1 + 2 + 3 + \dots + 8 = 36$	E1	
	$1 + 2 + 3 + \dots + 8 = 36$	<b>E</b> 1	
(ii)	80 200	B1	
(b) (i)	$2 (1 + 2 + 3 + \dots + n) = 2 \times \frac{n(n+1)}{2} = n (n+1)$	E1	both steps must be shown
(ii)	40 200	B1	
(iii)	40 000	B1ft	ft their (a)(ii) – their(b)(ii) or their (b)(ii) – 200 ft Not for zero or negative answer
(c) (i)	$\frac{2n(2n+1)}{2}$ oe final answer	B1	e.g. $2n^2 + n$
(ii)	$n^2$ cao	B2	M1 for their (c)(i) – $n(n + 1)$ or $n(n + 1) - n$ or $n/2(2+2(n-1))$