

**CAMBRIDGE**  
INTERNATIONAL EXAMINATIONS


**NOVEMBER 2002**

**INTERNATIONAL GCSE**



|   |
|---|
| <b>MARK SCHEME</b>  |
| <b>MAXIMUM MARK : 80</b>  |
| <b>SYLLABUS/COMPONENT : 0606/2</b><br><b>ADDITIONAL MATHEMATICS</b><br><b>(Paper 2)</b> |

| Page 1 | Mark Scheme                        | Syllabus | Paper |
|--------|------------------------------------|----------|-------|
|        | IGCSE Examinations – November 2002 | 0606     | 2     |

|       |  |  |
|-------|--|--|
| 1 [4] | $\text{Inverse} = \begin{pmatrix} 6 & -3 \\ -7 & 4 \end{pmatrix} \times \frac{1}{3}$ $\begin{pmatrix} x \\ y \end{pmatrix} = \frac{1}{3} \begin{pmatrix} 6 & -3 \\ -7 & 4 \end{pmatrix} \begin{pmatrix} -7 \\ -16 \end{pmatrix} = \begin{pmatrix} 2 \\ -5 \end{pmatrix}$   | B1 B1<br>M1 A1   |
| 2 [4] | $2^6 + 6 \times 2^5 \times x + \frac{6 \times 5}{1 \times 2} \times 2^4 \times x^2$ $= 64 + 192x + 240x^2$ <p>Replace <math>x</math> by <math>x - x^2 \Rightarrow</math> coefficient of <math>x^2 = -192 + 240 = 48</math></p>   | B2, 1, 0<br><i>(-1 each, incorrect or missing term)</i><br>M1 A1<br>C.S.C. |
| 3 [5] | <p>(i) Either <math>\frac{1 + \frac{1}{\sqrt{3}}}{1 - \frac{1}{\sqrt{3}}} \times \frac{1 + \frac{1}{\sqrt{3}}}{1 + \frac{1}{\sqrt{3}}}</math> or <math>\frac{\sqrt{3} + 1}{\sqrt{3} - 1} \times \frac{\sqrt{3} + 1}{\sqrt{3} + 1}</math></p> <p>Simplify <math>\Rightarrow 2 + \sqrt{3}</math></p> <p>(ii) <math>\frac{1}{p} = \frac{1}{2 + \sqrt{3}} \times \frac{2 - \sqrt{3}}{2 - \sqrt{3}} = \frac{2 - \sqrt{3}}{4 - 3}</math></p> $p - \frac{1}{p} = 2 + \sqrt{3} - (2 - \sqrt{3}) = 2\sqrt{3}$ <p>Or <math>p - \frac{1}{p} = 2 + \sqrt{3} - \frac{1}{2 + \sqrt{3}} = \frac{6 + 4\sqrt{3}}{2 + \sqrt{3}}</math></p> <p>Multiply by <math>\frac{2 - \sqrt{3}}{2 - \sqrt{3}} \Rightarrow 2\sqrt{3}</math></p> | M1<br>A1<br>M1 A1 ✓<br>A1<br>B1 ✓<br>M1 A1                                 |
| 4 [6] | <p>Solving inequalities:</p> <p>A <math>x &lt; 3.5</math></p> <p>B <math>x^2 - x - 2 = 0 \Rightarrow (x - 2)(x + 1) = 0 \Rightarrow x = -1, 2</math><br/> <math>x^2 - x - 2 &gt; 0 \Rightarrow x &lt; -1, x &gt; 2</math></p> <p>Required values <math>-5 &lt; x &lt; -1</math><br/> <math>2 &lt; x &lt; 3.5</math></p>  | B1<br>M1 A1<br>A1<br>M1 A1   |

|       |   |  |
|-------|---|--|
| 5 [6] | <p>(a) Either <math>{}_5C_3 = \frac{5 \times 4 \times 3}{1 \times 2 \times 3}</math> or <math>{}_4C_2 = \frac{4 \times 3}{1 \times 2}</math></p> <p>Product = <math>10 \times 6 = 60</math></p> <p>(b) Either, ending in 1 (or 3) <math>\Rightarrow 2 \times 5 \times 4</math> or, ending in 5 (or 7) <math>\Rightarrow 3 \times 5 \times 4</math></p> <p>Adding all 4 cases <math>\Rightarrow 40 + 40 + 60 + 60 = 200</math></p>   | B1<br>M1 A1<br>B1<br>M1 A1   |
| 6 [6] | <p>(i) <math>f(x) = -(x-1)(x-2)(x-k)</math></p> <p><math>f(3) = -2 \times 1 \times (3-k) = 8 \Rightarrow k = 7</math></p> <p>(ii) <math>f(-3) = -(-4)(-5)(-10) = 200</math></p>   | M1 A1<br>M1 A1<br>M1 A1  |
| 7 [6] | <p>(i) <math>\frac{d}{dx}(x \sin x) = \sin x + x \cos x</math></p> <p>(ii) <math>\int x \cos x dx = x \sin x - \int \sin x dx</math></p> <p><math>\int \sin x dx = -\cos x</math></p> <p><math>x \sin x + \cos x</math></p> <p><math>\frac{\pi}{2} - 1 \approx 0.571</math></p>   | M1 A1<br>M1<br>DM1<br>A1 A1 e.s.o.                                       |
| 8 [6] | <p>(i)  <math>[-\infty \text{ as } x \rightarrow 0; \text{ thro' } (1,0); \infty \text{ as } x \rightarrow \infty]</math></p> <p>(ii) Take logs <math>\ln x^2 + \ln e^{x-2} = \ln 1</math></p> <p><math>\Rightarrow 2 \ln x + x - 2 = 0</math></p> <p>Make <math>\ln x</math> the subject <math>\Rightarrow \ln x = -\frac{1}{2}(x-2) \Rightarrow</math> line is <math>y = 1 - x/2</math></p>                                      | B2,1,0<br>M1<br>A1<br>M1 A1  |
| 9 [7] | <p>(a) Correct combination of indices</p> <p>Either <math>(a^{2/3} - a^{1/3}b^{2/3} + b^{4/3}) \times a^{1/3} = a - a^{2/3}b^{2/3} + a^{1/3}b^{4/3}</math></p> <p>Or <math>(a^{2/3} - a^{1/3}b^{2/3} + b^{4/3}) \times b^{2/3} = a^{2/3}b^{2/3} - a^{1/3}b^{4/3} + b^2</math></p> <p>Sum = <math>a + b^2</math></p> <p>(b) <math>2^{2x+2} = 4 \times 2^{2x}</math> or <math>2^2 \times 4^x</math>      <math>5^{x-1} = 5^x \div 5</math>      <math>8^x = 2^{3x}</math></p> <p><math>\therefore 10^x = 4/5</math></p> | M1<br>A1<br>A1<br>B2,1,0<br>(-1 each incorrect or missing term)<br>M1 A1 |

|        |                                    |          |       |
|--------|------------------------------------|----------|-------|
| Page 3 | Mark Scheme                        | Syllabus | Paper |
|        | IGCSE Examinations – November 2002 | 0606     | 2     |

|                   |   |  |
|-------------------|---|--|
| 10 [9]            |  <p>(i) <math>AP = b/3 - a</math>                      <math>OM = a/2 + b/2</math></p> <p>(ii) <math>OQ = \lambda (a/2 + b/2)</math></p> <p>(iii) <math>OQ = OA + \mu AP = a + \mu (b/3 - a)</math></p> <p>(iv) Comparing coefficients    <math>\lambda/2 = 1 - \mu</math>    and    <math>\lambda/2 = \mu/3</math></p> <p>Solving                      <math>\lambda = 1/2</math>    <math>\mu = 2/3</math></p>   | B1    M1 A1<br>B1√<br>M1 A1√<br>M1<br>M1 A1              |
| 11 [11]           | <p>(i) <math>v = \int (\frac{3t}{2} - 6) dt = \frac{3t^2}{4} - 6t \quad (+c)</math></p> <p><math>[v]_{t=0} = 20 \Rightarrow c = 20</math>                      <math>[v]_{t=4} = 12 - 24 + 20 = 8</math></p> <p>(ii) <math>\int (\frac{3t^2}{4} - 6t + 20) dt = \frac{t^3}{4} - 3t^2 + 20t</math></p> <p><math>AB = [ ]_0^4 = 16 - 48 + 80 = 48</math></p> <p>(iii) <math>v_B = 8, \quad v_C = 20 \Rightarrow t_{BC} = (20 - 8) / 2 = 6</math></p> <p>(iv) </p> <p>curve</p> <p>straight line</p>                            | M1 A1<br>A1    A1<br>M1 A1√<br>A1<br>M1 A1√<br>B1<br>B1√ |
| 12 [10]<br>Either | <p><math>A = \pi r^2 + \pi r l \Rightarrow l = (120 - \pi r^2) / \pi r</math></p> <p><math>V = \frac{1}{2} \pi r^2 \left( \frac{\text{expression}}{\text{for } l} \right) = 60r - \frac{1}{2} \pi r^3 \quad (\text{AG})</math></p> <p><math>dV/dr = 60 - 3\pi r^2/2 = 0</math> when <math>r^2 = 40/\pi \approx 3.57</math></p> <p>Stationary value of <math>V \approx 143 \quad (142.73)</math></p> <p><math>d^2V/dr^2 = -3\pi r &lt; 0</math> for <math>r &gt; 0 \Rightarrow</math> maximum    [ or any valid method ]</p>   | B 1 M1<br>M1 A1<br>B1    M1 A1<br>A1<br>M1 A1            |
| Or                | <p>(i) <math>dy/dx = x^2 \times 1/x + 2x \ln x</math></p> <p>At Q, <math>y = 0 \Rightarrow \ln x = 0 \Rightarrow x = 1</math>                      <math>[dy/dx]_{x=1} = 1 \text{ c.s.o.}</math></p> <p>(ii) At P, <math>dy/dx = 0 \Rightarrow x(1 + 2 \ln x) = 0 \Rightarrow \ln x = -1/2</math></p> <p><math>\Rightarrow x = e^{-1/2} = 1/\sqrt{e} (\approx 0.6065) \quad (\text{AG})</math></p> <p>(iii) <math>d^2y/dx^2 = d(x + 2x \ln x) / dx = 1 + 2 \ln x + (2x \times 1/x)</math></p> <p><math>= 3 + 2 \ln x</math></p> <p><math>[d^2y/dx^2]_{x=1/\sqrt{e}} = 3 + 2(-1/2) = 2 \text{ c.s.o.}</math></p> | M1 A1<br>B1    A1<br>M1    A1<br>A1<br>M1 A1<br>A1       |