

**MARK SCHEME for the October/November 2006 question paper**

**0652 PHYSICAL SCIENCE**

**0652/03**

Paper 3 (Extended Theory), maximum raw mark 80

This mark scheme is published as an aid to teachers and students, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began.

All Examiners are instructed that alternative correct answers and unexpected approaches in candidates' scripts must be given marks that fairly reflect the relevant knowledge and skills demonstrated.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

The grade thresholds for various grades are published in the report on the examination for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses.

- CIE will not enter into discussions or correspondence in connection with these mark schemes.

CIE is publishing the mark schemes for the October/November 2006 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.



<b>Page 2</b>	<b>Mark Scheme</b>	<b>Syllabus</b>	<b>Paper</b>
	<b>IGCSE - OCT/NOV 2006</b>	<b>0652</b>	<b>3</b>

**General Rules.**

Apply unit penalty only once per question.

+ marks can only be scored if the previous mark has been scored.

In calculations, if the working/equation has not been asked for, and the answer is correct, then **all** the marks for that section must be scored.

Words in brackets preferable but not obligatory

Page 3	Mark Scheme	Syllabus	Paper
	IGCSE - OCT/NOV 2006	0652	3

1 (a)	one arrow upwards,	1	
	one arrow downwards	1	
	2.5N	1	
	2.5N or same as previous one	1	4
(b)	through the origin and linear to start with	1	
	curves upwards	1	
	limit of proportion clearly marked at beginning of curve	1	3
(c) (i)	mgh implied or seen	1	
	0.2 J	1	
	(ii) $\frac{1}{2}mv^2$ implied or seen equated to candidate's answer in (i)	1	
		1	
	1.3 m/s	1	5
			<b>Total 12</b>
2 (a) (i)	haematite	1	
	(ii) <u>carbon</u> burns/reacts with oxygen (producing carbon dioxide) carbon dioxide is reduced by / reacts with more carbon making carbon monoxide	1	
		1	
	(iii) $Fe_2O_3 + 3CO \rightarrow 2Fe + 3CO_2$ all formulae correct, equation balanced	1	
		+1	5
(b)	mass of iron(III) oxide in ore = $1 \times 80/100 = 0.8$ tonne	1	
	$Fe_2O_3 = 112 + 48 = 160$	1	
	mass of iron = $0.8 \times 112/160$	1	
	= 0.56	1	4
			<b>Total 9</b>
3 (a) (i)	reflection	1	
	(ii) $0.5 \pm 0.1$ (cm) $2.5 \text{ cm} \pm 0.5$ from candidate's figure	1	
		1	
	(iii) $v = f\lambda$ seen or implied 5.0 cm (ecf)	1	
		1	5
(b) (i)	diffraction	1	
	(ii) amount of diffraction/spreading/curvature depends on slit width larger slit less diffraction etc. (or vv)	1	
		1	3
			<b>Total 8</b>

Page 4	Mark Scheme	Syllabus	Paper
	IGCSE - OCT/NOV 2006	0652	3


4 (a)	(i)	copper		1	
	(ii)	Mg + CuSO <sub>4</sub> → MgSO <sub>4</sub> + Cu (all formulae correct, equation balanced)		1 +1	3
(b)		magnesium, aluminium, iron, copper (1 mark each for: Al after Mg; Fe after Al; Cu after iron)		3	3
(c)	(i)	Al is covered with a layer of aluminium oxide which protects the metal from contact with oxygen/water/air		1 1	
	(ii)	coat with zinc/galvanise OR mix with chromium to stop air/water reaching it/ to form stainless steel sacrificial layer		1 1	4
					<b>Total 10</b>
5 (a)	(i)	electromagnetic induction		1	
	(ii)	stronger magnets (not bigger)		1	
		faster rotation		1	
	(iii)	more turns in the coil		1	
change of flux (linkage) induces current each side cuts field upwards then downwards thus current induced in opposite directions		ANY 2		1 + 1	6
(b)	(i)	diode or l.e.d shown complete circuit with output terminals shown/load resistor included		1 +1	
	(ii)	opposite directions		1	3
					<b>Total 9</b>
6 (a)	(i)	diamond has a higher melting point diamond is harder diamond does not conduct electricity, graphite does	ANY 2	1 + 1	
	(ii)	<u>melting point &amp; hardness</u> diamond strong covalent bonds graphite weak (van der Waals) forces between layers	<u>conduction</u> diamond all electrons tightly bound graphite has mobile/free electrons	1 1	4
(b)	(i)	sea of electrons between particles/atoms/ions which move to produce the electric current		1 1	
	(ii)	particles/atoms/ions are in sheets/layers that can slide over each other		1 1	4
(c)	(i)	alloy is less malleable		1	1
	(ii)	new atoms are different size to original/layers/arrangement of atoms disrupted layers in alloy do not slide across each other as easily		1 1	2
					<b>Total 11</b>

Page 5	Mark Scheme	Syllabus	Paper
	IGCSE - OCT/NOV 2006	0652	3

7 (a)	(i)	evaporation at all temperatures – boiling at specific temperature	1	
		evaporation at surface – boiling in body of liquid	1	
		boiling the molecules have more energy than evaporation/higher energy molecules escape	1	3
(b)		liquid molecules much closer together or vv	1	
		intermolecular forces therefore much greater in liquids or vv	1	2
(c)		warms the room	1	1
(d)	(i)	P = VI seen or implied	1	
		I = 0.5 (A)	1	
	(ii)	R = V/I seen or implied	1	
		440 ( $\Omega$ )	1	
		<b>Both</b> units correct	1	5

**Total 11**

8 (a)	(i)	a family of compounds with similar properties/characteristics/reactions due to the presence of the same functional group/general molecular formula/of form $C_nH_{(2n+1)}OH$	1	
			1	2
(b)	(i)	ethene is reacted with steam	1	
		at high pressure/using a catalyst	+1	
	(ii)	$C_2H_4 + H_2O \rightarrow C_2H_5OH$	1	
	(iii)	fermentation/accept good description	1	
	(iv)	solvent/fuel	1	5



four covalent bond pairs of electrons shown on the carbon atom	1	
two covalent bond pairs of electrons shown on the oxygen atom	1	
four extra electrons shown on oxygen atom	1	3

(electrons do not need to be distinguished in any way)

**Total 10**