## UNIVERSITY OF CAMBRIDGE LOCAL EXAMINATIONS SYNDICATE

Joint Examination for the School Certification and General Certificate of Education Ordinary Level

CHEMISTRY 5070/1

PAPER 1 Multiple Choice

**OCTOBER/NOVEMBER SESSION 2002** 

1 hour

Additional materials:

Multiple Choice answer sheet
Soft clean eraser
Soft pencil (type B or HB is recommended)

TIME 1 hour

#### **INSTRUCTIONS TO CANDIDATES**

## Do not open this booklet until you are told to do so.

Write your name, Centre number and candidate number on the answer sheet in the spaces provided unless this has already been done for you.

There are **forty** questions in this paper. Answer **all** questions. For each question, there are four possible answers, **A**, **B**, **C** and **D**. Choose the **one** you consider correct and record your choice in **soft pencil** on the separate answer sheet.

Read very carefully the instructions on the answer sheet.


#### INFORMATION FOR CANDIDATES

Each correct answer will score one mark. A mark will not be deducted for a wrong answer.

Any rough working should be done in this booklet.

A copy of the Periodic Table is printed on page 16.

- 1 Which property of a gas affects the rate at which it spreads throughout a laboratory?
  - A boiling point
  - **B** molecular mass
  - **C** reactivity
  - D solubility in water
- 2 The graph gives the melting points of mixtures of lead and tin.



The graph shows that any mixture of lead and tin must have a melting point

- A above that of tin.
- **B** below that of lead.
- **C** below that of both tin and lead.
- **D** between that of tin and lead.
- **3** From which mixture can the underlined substance be obtained by adding water, stirring and filtering?
  - A <u>calcium carbonate</u> and sodium chloride
  - **B** copper(II) sulphate and sodium chloride
  - C ethanoic acid and ethanol
  - **D** iron and <u>magnesium</u>

4 An aqueous solution of a sulphate is made from a solid hydroxide, of a metal **M**, by the reaction:

$$M(OH)_2(s) + H_2SO_4(aq) \longrightarrow MSO_4(aq) + 2H_2O(l)$$

For which hydroxide would the method not work?


- A barium hydroxide
- B copper(II) hydroxide
- **C** iron(II) hydroxide
- **D** magnesium hydroxide
- 5 Which ion has the most shells that contain electrons?
  - **A**  $Al^{3+}$
  - **B** Be<sup>2+</sup>
  - **C** N<sup>3-</sup>
  - **D** S<sup>2-</sup>
- **6** The table gives data about four substances.

Which substance could be an ionic compound?

compound	melting point/°C	electrical conductivity in aqueous solution
Α	<b>-73</b>	good
В	32	poor
С	474	poor
D	805	good

7 The number of outer shell electrons for the atoms of the first 12 elements in the Periodic Table is plotted against the proton number of the element.

Which graph is obtained?



8 The table shows the electron structures of four elements.

element	electronic structure
W	2, 6
X	2, 8
Y	2, 8, 1
z	2, 8, 7

Which pair of atoms will form a covalent substance?

- Α two atoms of W
- В two atoms of X
- C an atom of W and an atom of X
- an atom of Y and an atom of Z D
- 9 Which substance contains covalent bonds, but also conducts electricity?
  - Α brass
  - В graphite
  - C iodine
  - D steel
- **10** One mole of each of the following compounds is burnt in excess oxygen.

Which compound will produce three moles of carbon dioxide and three moles of steam only?


- $A C_3H_8$

- **B**  $C_3H_7OH$  **C**  $C_3H_7CO_2H$  **D**  $CH_3CO_2CH_3$
- 11 When zinc reacts with dilute sulphuric acid a gas is released.

What happens to the zinc and what is the gas released?

	the zinc is	the gas is
Α	oxidised	hydrogen
В	oxidised	sulphur dioxide
С	reduced	hydrogen
D	reduced	sulphur dioxide

12 Which pair of metals **X** and **Y** will produce the highest voltage when used as electrodes in a simple cell?




	metal X	metal <b>Y</b>
Α	copper	silver
В	magnesium	silver
С	magnesium	zinc
D	zinc	copper

**13** Four electrolytes were electrolysed using carbon electrodes.

Which set of data is correct?


	electrolyte	produ	uct at
	electrolyte	anode	cathode
Α	CuSO <sub>4</sub> (aq)	oxygen	copper
В	NaC <i>l</i> (aq)	chlorine	sodium
С	NaH (I)	sodium	hydrogen
D	PbBr <sub>2</sub> (I)	lead	bromine

- 14 Which pair of substances are isotopes?
  - **A**  ${}^{12}_{6}$ C and  ${}^{14}_{6}$ C
  - B carbon dioxide and carbon monoxide
  - C diamond and graphite
  - **D**  $C_2H_4$  and  $C_3H_6$
- 15 Which step in the diagram shows the process of photosynthesis?



16 Dissolving ammonium nitrate in water is endothermic.


Which graph shows how the temperature alters as the ammonium nitrate is added to water and then the solution is left to stand?



17 If a strip of magnesium is dropped into excess hydrochloric acid an exothermic reaction occurs.

The rate of this reaction increases during the first few seconds because

- **A** the amount of magnesium is decreasing.
- **B** the magnesium is acting as a catalyst.
- **C** the solution is becoming hotter.
- **D** the surface area of the magnesium is increasing.
- 18 The energy profile diagrams show how adding a substance **X** to a reaction mixture changes the reaction pathway.



Which change is likely to be observed when **X** is added to the reaction mixture?

- A The reaction becomes less exothermic.
- **B** The reaction becomes more exothermic.
- **C** The speed of the reaction decreases.
- **D** The speed of the reaction increases.
- **19** Which process does **not** involve either oxidation or reduction?
  - A formation of ammonium sulphate from ammonia and sulphuric acid
  - **B** formation of nitrogen monoxide from ammonia
  - **C** formation of sulphuric acid from sulphur
  - **D** formation of zinc from zinc blende (ZnS)

						9						
20		separate exp dually added			s of aqu	eous	sodium	n hydroxid	de or	aqueous	ammoni	a was
		ooth experim gent.	nents, a pr	ecipitate	was obta	ained	d which	dissolved	d in a	an excess	s of the	added
	Wh	at could <b>X</b> co	ontain?									
	Α	copper(II) r	nitrate									
	В	iron(II) nitra	ate									
	С	iron(III) nitr	ate									
	D	zinc nitrate										
21		excess of d ium chloride.	-				-		rium	hydroxide	and aq	ueous
	Α	A gas is pro	oduced.									
	В	An insoluble	e salt is pro	duced.								
	С	The final pl	H is 7.									
	D	Water is pro	oduced.									
22	Wh	ich property	decides the	order of	the elem	ents	in the P	eriodic Ta	ble?			
	Α	the masses	of their ato	oms								
	В	the number	of electron	s in the c	outer shell	l						
	С	the number	of neutron	s in the n	ucleus							
	D	the number	of protons	in the nu	cleus							
23	The	proton num	ber of indiu	m, In, is	49.							
	Wh	at is the mos	st likely form	nula for th	ne oxide o	of indi	ium?					
	Α	In <sub>2</sub> O	В	$In_2O_3$		С	InO		D	InO <sub>2</sub>		
24	Wh	ich element i	in the table	is likely t	o be a tra	nsitic	on meta	l?				
		ment	melting po	-	colour (							
		A	high			olue	ondo					

[Turn over www.theallpapers.com

green

white

white

В

C

D

low

high

low

- 10 25 Which feature of a metal's structure is responsible for it conducting electricity? Α It contains positive ions. В It has a "sea of electrons". C Its ions are tightly packed together. D Its positive ions attract electrons. 26 Aluminium is extracted from purified bauxite by electrolysis but iron is extracted from haematite by reduction with coke. Why is iron not extracted by electrolysis? Haematite needs to be purified but bauxite does not. В Iron is less reactive than aluminium. C Reduction with coke is cheaper than electrolysis. D Reduction with coke gives a purer product than electrolysis.
  - **27** Old steel drums corrode quickly in a damp atmosphere but aluminium cans do not.


Which of the following correct statements explains this behaviour of aluminium?

- **A** Aluminium has only one valency.
- **B** Aluminium has a lower density than iron.
- **C** Aluminium is above iron in the activity series.
- **D** Aluminium is protected by its oxide layer.
- 28 Caesium is a metal that is more reactive than aluminium.


Which reaction would produce caesium?

- A electrolysing aqueous caesium chloride
- B electrolysing molten caesium chloride
- **C** heating caesium carbonate
- **D** heating caesium oxide with carbon
- **29** Which of the following gases **cannot** be removed from the exhaust gases of a petrol powered car by its catalytic converter?
  - A carbon dioxide
  - B carbon monoxide
  - C hydrocarbons
  - **D** nitrogen dioxide

**30** A 100 cm<sup>3</sup> sample of bottled gas used for diving was placed in a gas syringe in the apparatus shown.



The gas was passed backward and forward over heated copper turnings. The results obtained were used to plot the graph.



What is the percentage of oxygen in the bottled gas?

**A** 20%

**B** 30%

**C** 70%

80%

31 In the Haber process, nitrogen and hydrogen react to form ammonia.

$$N_2(g) + 3H_2(g) \rightleftharpoons 2 NH_3(g) \Delta H = -92 kJ/mol$$

Which factor increases both the speed of reaction and the amount of ammonia produced?

- A addition of a catalyst
- **B** decreasing the temperature
- **C** increasing the pressure
- **D** increasing the temperature

**32** Nitrates from fertilisers used on farmland can cause pollution.


Why do nitrates pollute rivers?

- A Nitrates are salts.
- **B** Nitrates are very soluble in water.
- C Nitrates contain oxygen.
- **D** Nitrate ions are negatively charged.
- 33 Which representation of dilute sulphuric acid is correct?
  - **A**  $H_2(aq) + SO_4^{2-}(aq)$
  - **B**  $2H^{+}(aq) + SO_4^{2-}(aq)$
  - **C**  $2H^{+}(aq) + SO_{4}^{-}(aq)$
  - $\mathbf{D}$   $H_2SO_4(I)$
- 34 Which statement describes what happens when hydrogen and oxygen are used in a fuel cell?
  - A Electricity is generated directly.
  - **B** Electricity is used to produce water.
  - **C** Hydrogen is burned to form steam.
  - **D** Hydrogen reacts to form a hydrocarbon fuel.
- 35 The structures of an acid and an alcohol are shown.

Which pairing of names correctly identify the two compounds?

	acid	alcohol
Α	ethanoic	butanol
В	ethanoic	propanol
С	propanoic	propanol
D	propanoic	butanol

- 36 Which physical property of the alkanes does not increase as relative molecular mass increases?
  - A boiling point
  - **B** flammability
  - C melting point
  - **D** viscosity
- 37 The structures of four organic compounds are shown.



Which compounds decolourise bromine water?


- **A** 1 and 2
- **B** 1, 2 and 4
- **C** 2 and 4
- **D** 3 and 4
- 38 A polymer X was hydrolysed and the two products were

and

What can be deduced about X?

- A It was a condensation polymer.
- B It was starch.
- **C** It was made by addition polymerisation.
- **D** It was *Terylene*.

**39** Which polymer has the empirical formula CH?



- 40 In the polymerisation of ethene to form poly(ethene), there is no change in
  - A boiling point.
  - B density.
  - C mass.
  - **D** molecular formula.

# **BLANK PAGE**

The Periodic Table of the Elements **DATA SHEET** 

							ב ב ב	OIC IAU	Group  Group		2							_
	=											=	≥	>	5	<b>=</b>	0	
							Hydrogen										Helium	
7 <b>Li</b> thium	Be Beryllium					-		_				11 Boron 5	12 Carbon	Nitrogen 7	Oxygen 8	19 Fluorine	20 <b>Ne</b> on 10	
Na Sodium	Mg Magnesium											27 <b>A 1</b> Aluminium 13	28 <b>Si</b> Silicon	31 Phosphorus	32 <b>Sulphur</b> 16	35.5 <b>C1</b> Chlorine	40 <b>Ar</b> Argon 18	
39 <b>K</b> Potassium	40 Calcium	Scandium	48 <b>T</b> itanium 22	51 Vanadium 23	52 <b>Cr</b> Chromium 24	Mn Manganese	56 From Iron	59 <b>Co</b> Cobalt	59 Nickel	64 Copper	65 <b>Zn</b> Zinc 30	70 <b>Gal</b> ium 31	73 Germanium	AS Arsenic	Selenium 34	80 Bromine	84 <b>K</b> rypton 36	
Rubidium	88 <b>Sr</b> Strontium	89 <b>Y</b> ttrium 39	2 Zrconium	Niobium 41	96 Mo Molybdenum 42	Tc Technetium	Ruthenium	Rh Rhodium	Pd Palladium	108 <b>Ag</b> Silver 47	Cadmium 48	115 Indium 49	119 <b>Sn</b> Tin	Sb Antimony	128 <b>Te</b> Tellurium 52	127 <b>I</b> lodine 53	131 <b>Xe</b> Xenon 54	
Caesium	137 <b>Ba</b> Barium 56	139 <b>La</b> Lanthanum 57 *	178 <b>‡</b> Hafnium	181 <b>Ta</b> Tantalum 73	184 <b>W</b> Tungsten 74	Re Rhenium	190 <b>OS</b> Osmium 76	192 <b>Ir</b> Iridium	195 <b>Pt</b> Platinum 78	Au Gold 799	Hg Mercury	204 <b>T L</b> Thallium	207 <b>Pb</b> Lead	209 <b>Bi</b> Bismuth	<b>Po</b> Polonium 84	At Astatine	Radon 86	
<b>Fr</b> Francium	226 <b>Ra</b> Radium 88	AC Actinium 89 t		_							_	-			_	_		
1 L;	*58-71 Lanthanoid series †90-103 Actinoid series	l series series	1	140 <b>Ce</b>	Praseodymium	Neodymium	<b>Pm</b> Promethium	Samarium	152 <b>Eu</b> Europium	157 <b>Gd</b> Gadolinium	159 <b>Tb</b> Terbium	162 <b>Dy</b> Dysprosium	165 <b>Ho</b> Holmium	167 <b>Er</b> Erbium	Tm Thulium	Yb Ytterbium	Lutetium	

169	Ę	Thulium	69	Md Mendelevium
167	ங்	Erbium	89	Fm Fermium
165	웃	Holmium	29	Einsteinium
162	ò	Dysprosium	99	Californium
159	Д	Terbium	92	<b>Bk</b> Berkelium
157	В	Gadolinium	64	Curium 96
152	En	Europium	63	Americium
150	Sm	Samarium	62	<b>Pu</b> Plutonium
	Pm	Promethium	61	Neptunium
144	PZ	Neodymium	09	L Uranium
141	፵	Praseodymium	59	<b>Pa</b> Protactinium
140	පී	Cerium	58	232 <b>Th</b> Thorium

۲

Nobelium

The volume of one mole of any gas is 24 dm<sup>3</sup> at room temperature and pressure (r.t.p.).

Key

b = proton (atomic) number

a = relative atomic mass X = atomic symbol