UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS Pre-U Certificate

MARK SCHEME for the May/June 2011 question paper for the guidance of teachers

9790 BIOLOGY

9790/02

Paper 2 (Structured), maximum raw mark 85

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

• Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, Pre-U, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02

Abbreviations, annotations and conventions used in the Mark Scheme

1	=	alternative and acceptable answers for the same marking point
;	=	separates marking points
R	=	reject
()	=	words which are not essential to gain credit
	=	(underlining) key words which <u>must</u> be used to gain credit
ecf	=	error carried forward
AW	=	alternative wording
Α	=	accept
ora	=	or reverse argument
1		

1 (a) (i) bacterium / prokaryote; A bacteria / bacillus

[1]

[3]

- (ii) A cell wall;
 - **B** cell membrane;
 - C (bacterial) chromosome / DNA / chromoneme ; A nucleoid
- (iii) (70S) ribosome; [1]
- (iv) award two marks for the correct answer with appropriate unit $(0.35 \ \mu m, 350 \ nm, 0.00035 \ mm)$

if incorrect answer or no answer award one mark for dividing measurement by magnification, accept \pm 1 mm

actual size = 35 mm

35 / 35 000 / AW 100 000

 $0.35 \mu m / 350 nm / 0.00035 mm$;;

A
$$0.34 - 0.36 \mu \text{m} / 340 - 360 \text{ nm} / 0.00034 - 0.00036 \text{ mm} / 3.4 - 3.6 x 10^{-4} mm [2]$$

- (b) 1 (bacteria stained) blue / purple, are Gram +ve;
 - 2 Gram +ve bacteria have a thick(er), cell wall / peptidoglycan layer (than Gram -ve);
 - 3 Gram +ve lack outer, lipid / lipopolysaccharide, membrane / layer; ora
 - 4 crystal violet / iodine, stains, cell wall / peptidoglycan, blue / purple;
 - 5 not washed away when, decolourising solution / alcohol / propanone, is added;
 - **6** Gram -ve stains, red / pink ; **A** not blue *idea that*
 - 7 membrane is removed when, alcohol / acetone, is added so, crystal violet / iodine, does not remain;
 - 8 wall is thinner so any, crystal violet / iodine, is washed out;
 - **9** AVP; [max 4]

[Total:11]

Page 3	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02
	-	•	

2 (a) (i) glycosidic bond shown as forming between OH on C1 and OH on C4, and glucose residues rotated so that -O- in the rings are opposite each other;

bond formation may be shown diagonally between C1 and C4 or by rotating one of the glucose molecules so the OH groups are adjacent

water molecule formed; [2]

(ii) $(\beta 1-4)$, glycosidic; [1]

- (b) similarities
 - 1 have –OH groups / both have more than one –OH group;
 - 2 have an aldehyde group; A both are aldoses;
 - 3 have a ring structure / have an O in the ring;
 - 4 can also be straight chains;
 - **5** CH₍₂₎O;

6 AVP; [max 2]

difference

(ribose is) pentose / 5C, (β glucose is) hexose / 6C; A C₅H₁₀O₅, C₆H₁₂O₆ [1]

- (c) (i) 1 compact;
 - 2 insoluble;
 - 3 osmotically, inactive / inert;
 - 4 ref to water potential;
 - 5 easily formed / easily recovered or mobilised;
 - 6 ref to free ends;
 - 7 ref to branching of amylopectin providing, free ends / easy mobilisation;
 - 8 amylopectin ideal storage molecule;
 - 9 ref to energy storage; A respiratory substrate [max 4]
 - (ii) more branching points allows fast, storage / mobilisation, of glucose; idea that may have arisen by mutation / evolutionary accident; [1]

[Total: 11]

Page 4			Mark Scheme: Teachers' version	Syllabus	Paper
			Pre-U – May/June 2011	9790	02
(a)			nisms share the same genetic code / AW ; ription is possible / promoters work in the same way		[1]
(b)	(i)	1 2 3 4 5	role of promoter is to switch on genes; ref to, transcription; A described only in the rice endosperm; not expressed anywhere else (would be a waste of en in edible part of seed;	ergy / resources); [max 2]
	(ii)	1 2 3 4 5	use restriction, enzyme / endonuclease; cut plasmid / open plasmid; sticky ends / complementary bases; ref to hydrogen bonding / A - T / C - G; role of ligase, in sealing sugar-phosphate backbone / /	AW;	[max 3]
	(iii)	1 2 3	can insert into plant DNA / carries other genes into plant ref to vector;	_	[max 2]
		3	ref to, tumour inducing / Ti / genes concerned with inse	eruon ;	[IIIax 2]
	(iv)	mic	roprojectile / biolistics / gene gun / micropipette;		[1]
(c)	1 2 3	slow othe	o (of several) with lowest rate; ving entire pathway / determines rate of rest of pathwer steps / AW; o that controls rate at which product is made;	ay / step that c	ontrols rate of
(d)	1 2 3 4 5	by g by, I by n	ease expression of <i>psy</i> ; genes from other sources; better / different, promoter; nore than one copy of gene; P; e.g. increase rate of production of enzyme, ref to rol	e of gene produ	[max 2] ct
(e)	(i)	1 2 3 4 5 5	gene transfer to other rice; contamination of other crop (e.g. ref to organic crop); gene transfer to wild relative; gene transfer via bacteria or viruses with unknown efferecipient plant outcompetes others / ref to 'superweed toxic to non pests; AVP; must be something specific not a vague answer.	,	[max 1]
	(ii)	1 2 3 4 5	allergy; long term toxicity; antibiotic resistance of gut bacteria if antibiotic marker gene transfer to, virus to human cells / gut bacteria, w if idea not used in (i) AVP;		ect; [max 1]

Mark Scheme: Teachers' version

Syllabus

Paper

Page 4

3

[Total: 15]

Page 5	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02

- 4 (a) (i) multipotent cells can differentiate into a limited number of cell types (but not into whole organism);
 - (ii) 1 external growth factors (first messengers) / mitogens / cytokines / AW;
 - 2 receptors on cell (surface) membrane;
 - 3 activate kinases;
 - 4 kinases phosphorylate enzymes (and alter their activity);
 - 5 activate, transcription factors / promoters;
 - 6 increase production of cyclins;
 - 7 cyclins / kinases control stages of, cell cycle / mitosis;

A ref to checkpoints during cell cycle

[max 3]

- (iii) 1 each stem cell contains a complete genome;
 - 2 genes for all features required by different cell types;
 - a example of cell required in tendons;
 - e.g. cells in wall of blood vessels, connective tissue cells, fibroblasts
 - **4** cells will be in slightly different conditions;
 - 5 different genes will be expressed / AW, in different conditions;
 - 6 ref to role of gene product;
 - 7 gene expression alters the possibility of expression of further genes;
 - 8 influence of signals from adjoining cells;
 - 9 influence of signals from distant cells; [max 3]
- (b) 1 diagram shows (TS) of, sarcomere / myofibril / thick and thin filaments;
 - 2 in relaxed muscle;

credit these points if they are on a diagram of LS sarcomere / thick and thin filaments

- 3 Q shows, actin / thin, filaments, alone:
- 4 R shows, myosin / thick, filaments alone;
- **5** P is overlap between thick and thin filaments;
- 6 Q is I band;
- 7 P is (overlap region of) A band;
- 8 R is H zone;
- **9** if contracted muscle, then, no / little, I band and H zone;

[max 4]

- (c) at end of motor neurone to max 3
 - 1 neuromuscular junction / motor end plate;
 - 2 calcium ions enter / AW, when action potential arrives;
 - 3 stimulates vesicles to, move towards / fuse with, pre-synaptic membrane;
 - 4 further detail; e.g. voltage gated channel proteins for calcium ions

muscle to max 3

- 5 impulse / action potential, in, sarcolemma / T-tubules, stimulates release of calcium ions from sarcoplasmic reticulum;
- 6 calcium ions combine with troponin;
- 7 movement of tropomyosin;
- 8 exposing myosin binding sites on thin filaments;
- 9 actin and myosin interact;
- 10 further detail of role of calcium ions in muscle;
- 11 when no action potential calcium ions pumped back into sarcoplasmic reticulum; [max 5]

[Total: 16]

Page 6	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02

- 5 (a) flightless birds in New Zealand
 - 1 evolved from birds that could fly / flighted birds migrated to New Zealand;
 - 2 ref to founder effect / rapid evolution on islands;
 - 3 absence of mammals;
 - 4 available niches;

credit ref to takahē as a ground-dwelling, herbivorous animal / grazer, occupying a niche that is associated with mammals elsewhere

- 5 abundant, food / resources, on ground;
- 6 nesting sites available on the ground;
- 7 few / no, predators;
- 8 advantage in being flightless as flight requires considerable energy;
- 9 ref to (natural) selection;
- **10** AVP; e.g. ref to non-flying, trait / allele(s)

[max 4]

- (b) 1 humans introduced, predators / invasive species / domesticated species;
 - 2 named example(s), e.g. stoats, rats;
 - 3 habitat change as a result of environmental changes;
 - 4 isolated / trapped, populations;

A geographical isolation or ref from passage about population on mountains

- 5 cannot fly to other areas;
- 6 ref to effects of inbreeding in small populations;
- 7 ref to, pathogens / disease;
- 8 ref to, habitat destruction by humans / other appropriate human influence(s);
- **9** AVP; [max 3]
- (c) 1 fundamental niche represents the possible range of a species;
 - 2 realised niche may be less than ideal;
 - 3 range restricted by limiting factors;
 - 4 including interactions with other organisms;
 - 5 e.g. interspecific competition / named example;
 - 6 e.g. food availability;
 - 7 e.g. presence of predators / disease;
 - 8 realised niche has features of fundamental niche;
 - **9** features must be identified before endangered species is transferred;
 - 10 use of information about the fundamental niche of the takahē;
 - 11 use of information about the realised niche of the takahē;
 - 12 realised niche *now* may not be optimal as a result of changes in the past;
 - **13** AVP:

[Total: 11]

Page 7	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02

6 (a) award one mark for each feature plus its adaptation. Some features have the same adaptation.

	features	adaptations	
1	wide / large cross sectional area / large diameter	transport large volume of water / low resistance to flow;	
2	lack of cross walls / continuous columns	low resistance to flow / allows continuous flow of water;	
3	empty / no cytoplasm / AW	low resistance to flow;	
4	pits R 'gaps'	permit lateral movement / AW; allows movement around cavitation;	
5	thickening of (secondary) wall A lignification	provides mechanical support / strengthens wall to prevent collapse under tension; R strength unqualified / rigid R resisting positive pressure	
6	annular / spiral, thickening	allows xylem vessels to stretch during growth without collapsing;	
7	lignin in walls	waterproofing wall (so water remains inside lumen);	
8	AVP ; e.g. hydrophilic nature of cellulose	e.g. ref to adhesion of water to walls of xylem vessels	

R adhesion of water to lignin as it is hydrophobic

R refs to capillarity [max 4]

- (b) (i) loblolly pine
 - 1 comparative descriptive comment;
 - e.g. time delay between high rates of transpiration and water absorption
 - 2 comparative data quote (transpiration and water absorption);
 - 3 environmental conditions promote stomatal opening and loss of water vapour;
 - 4 idea that transpiration drives water absorption;
 - 5 transpiration sets up a water potential gradient;
 - 6 ref to cohesion-tension from leaf to root;
 - 7 water enters roots;

rate of transpiration decreases after 1400 hours

- 8 reaches water stress at 1400 hours;
- 9 stomata close;
- 10 stated environmental conditions change after 1400 hours;
- **11** AVP; [max 4]

Page 8	Mark Scheme: Teachers' version	Syllabus	Paper
	Pre-U – May/June 2011	9790	02

- (ii) 1 prickly pear / Opuntia ficus indica, is a xerophyte;
 - 2 example of xeromorphic feature;
 - 3 very low rate of transpiration / data quote;
 - 4 rate of transpiration peaks at 1800 hrs, after hottest part of the day;
 - 5 (suggests that) prickly pear / Opuntia ficus indica, is a CAM plant;
 - 6 stomata closed during the day / only open at night;

[max 3]

[Total:11]

7 (a) (i) (theca of) follicle (in ovary);

A adrenal cortex

[1]

(ii) 1 increase in concentration of 17β -oestradiol (during first half of cycle);

oestrogen / 17β-oestradiol

- 2 acts on endometrium;
- 3 to stimulate, repair / cell division / growth of blood vessels / growth of glands;
- 4 thickening;
- 5 AVP; [max 3]
- **(b) 1** hormones act by <u>negative feedback</u> on, anterior pituitary / hypothalamus;
 - **2** decrease release of gonadotrophin-releasing hormone / GnRH;
 - 3 decrease release of FSH, so development of follicles inhibited;
 - 4 no oestrogen for positive feedback which would release (surge) of LH (mid cycle);
 - 5 no, ovulation / release of oocyte;
 - 6 stimulates thickening of cervical mucus;
 - 7 prevents thickening of endometrium / AW;

[max 3]

- (c) woman may be infertile because of high concentration of oestrogen / failure of negative feedback control of oestrogen secretion
 - 1 clomiphene, interacts with / blocks, oestrogen receptors;
 - 2 acts as a competitive oestrogen receptor antagonist on, gonadotrophin-secreting cells in anterior pituitary / GnRH secreting cells in hypothalamus;
 - 3 reduces negative feedback effect of oestrogen;
 - 4 gonadotrophin-releasing hormone / GnRH, secreted from hypothalamus;
 - 5 increases secretion of, FSH / LH, from anterior pituitary;

[max 3]

[Total: 10]