



# Victorian Certificate of Education 2009

SUPERVISOR TO ATTACH PROCESSING LABEL HERE

## STUDENT NUMBER

Letter

Figures

Words

|  |  |  |  |  |  |  |  |  |  |
|--|--|--|--|--|--|--|--|--|--|
|  |  |  |  |  |  |  |  |  |  |
|  |  |  |  |  |  |  |  |  |  |

# CHEMISTRY

## Written examination 2

Thursday 12 November 2009

Reading time: 9.00 am to 9.15 am (15 minutes)

Writing time: 9.15 am to 10.45 am (1 hour 30 minutes)

## QUESTION AND ANSWER BOOK

### Structure of book

| Section | Number of questions | Number of questions to be answered | Number of marks |
|---------|---------------------|------------------------------------|-----------------|
| A       | 20                  | 20                                 | 20              |
| B       | 7                   | 7                                  | 56              |
|         |                     |                                    | Total 76        |

- Students are permitted to bring into the examination room: pens, pencils, highlighters, erasers, sharpeners, rulers and one scientific calculator.
- Students are NOT permitted to bring into the examination room: blank sheets of paper and/or white out liquid/tape.

### Materials supplied

- Question and answer book of 20 pages.
- A data book.
- Answer sheet for multiple-choice questions.

### Instructions

- Write your **student number** in the space provided above on this page.
- Check that your **name** and **student number** as printed on your answer sheet for multiple-choice questions are correct, **and** sign your name in the space provided to verify this.
- All written responses must be in English.

### At the end of the examination

- Place the answer sheet for multiple-choice questions inside the front cover of this book.
- You may keep the data book.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

**SECTION A – Multiple-choice questions****Instructions for Section A**

Answer **all** questions in pencil on the answer sheet provided for multiple-choice questions.

Choose the response that is **correct** or that **best answers** the question.

A correct answer scores 1, an incorrect answer scores 0.

Marks will **not** be deducted for incorrect answers.

No marks will be given if more than one answer is completed for any question.

**Question 1**

The addition of a catalyst to a chemical reaction

- A. lowers the activation energy required for the reaction to occur.
- B. lowers the chemical energy of the products.
- C. lowers the chemical energy of the reactants.
- D. lowers the value of the enthalpy change for the reaction.

**Question 2**

The two statements below give possible explanations for changes that occur when the temperature of a reaction mixture is increased.

I At a higher temperature, particles move faster and the reactant particles collide more frequently.


II At a higher temperature, more particles have energy greater than the activation energy.

Which alternative below best explains why the observed reaction rate is greater at higher temperatures?

- A. I only
- B. II only
- C. I and II to an equal extent
- D. I and II, but II to a greater extent than I

**Question 3**

The change in energy during a reaction is represented in the following energy profile diagram.




The change in energy labelled  $\Delta E$  above is


- A. the energy absorbed when bonds in the reactants break.
- B. the activation energy of the forward reaction.
- C. the activation energy for the reverse reaction.
- D. the heat of reaction.

**Question 4**

If, for the reaction



then the  $\Delta H$  value for



would be

- A. +2728 kJ mol<sup>-1</sup>
- B. +1364 kJ mol<sup>-1</sup>
- C. +682 kJ mol<sup>-1</sup>
- D. -1364 kJ mol<sup>-1</sup>

**Question 5**

The concentrations of reactants and products were studied for the following reaction.



In an experiment, the initial concentrations of the gases were


$$[\text{H}_2] = 0.0200 \text{ M}, [\text{F}_2] = 0.0100 \text{ M} \text{ and } [\text{HF}] = 0.400 \text{ M}$$

When the reaction reaches equilibrium at 25°C, the concentration of HF will be

- A. 0.400 M
- B. 0.420 M
- C. between 0.400 M and 0.420 M
- D. less than 0.400 M

**Question 6**

The anaesthetic, nitrous oxide, N<sub>2</sub>O, decomposes to form an equilibrium mixture of N<sub>2</sub>O, N<sub>2</sub> and O<sub>2</sub> according to the following equation.



At 25°C,  $K = 7.3 \times 10^{37}$  M and at 40°C,  $K = 2.7 \times 10^{36}$  M

What valid conclusion can be made from this?

- A. The equilibrium concentrations of N<sub>2</sub> and O<sub>2</sub> are equal at 25°C.
- B. The equilibrium concentration of N<sub>2</sub>O is higher at 25°C than at 40°C.
- C. N<sub>2</sub>O is less stable at the higher temperature.
- D. The forward reaction is exothermic.

**Question 7**


In a flask, 10.0 mL of a 0.100 M HCl solution is diluted to 1.00 L. In a second flask, 10.0 mL of a 0.100 M KOH solution is also diluted to 1.00 L.

Which statement best describes the changes in pH in these flasks?

- |    | pH change of the HCl solution | pH change of the KOH solution |
|----|-------------------------------|-------------------------------|
| A. | increases by 2                | decreases by 2                |
| B. | increases by 2                | increases by 2                |
| C. | decreases by 2                | increases by 2                |
| D. | decreases by 2                | decreases by 2                |

**Question 8**

The value of the ionisation constant,  $K_w$ , of a sample of pure water at different temperatures is shown in the graph below.



Which one of the following statements about the effect of increasing temperature on the pH and acidity of water is correct?

- A. The pH is always 7 and the water remains neutral.
- B. The pH decreases and the water remains neutral.
- C. The pH decreases and the water becomes acidic.
- D. The pH increases and the water remains neutral.

**Question 9**

The following table contains information about three experiments. In each experiment 0.10 mol of an alkane is burned completely and all the energy released is used to heat 1.00 L of water which was initially at 20°C.


| experiment | alkane  | molecular formula              |
|------------|---------|--------------------------------|
| I          | butane  | C <sub>4</sub> H <sub>10</sub> |
| II         | pentane | C <sub>5</sub> H <sub>12</sub> |
| III        | hexane  | C <sub>6</sub> H <sub>14</sub> |

In which experiment(s) will the water be heated to its boiling temperature?

- A. III only
- B. II and III only
- C. I and II only
- D. I, II and III

**Question 10**

Potassium hydroxide and hydrochloric acid react in aqueous solution according to the following equation.




A 50 mL solution containing 0.025 mol of KOH was mixed rapidly in an insulated vessel with a 50 mL solution containing 0.025 mol of HCl. The temperature increased by 3.5°C.

Assuming that the specific heat capacity of the solution is the same as that of water, the enthalpy change,  $\Delta H$ , of this reaction, in  $\text{kJ mol}^{-1}$ , is closest to

- A. -29
- B. -59
- C.  $-2.9 \times 10^4$
- D.  $-5.9 \times 10^4$

**Question 11**

Two standard galvanic cells are shown below.




On the basis of the polarity of the electrodes shown above, which one of the following reactions would **not** be expected to occur spontaneously?


- A.  $\text{Co}^{2+}(\text{aq}) + \text{Cd}(\text{s}) \rightarrow \text{Co}(\text{s}) + \text{Cd}^{2+}(\text{aq})$
- B.  $2\text{Mn}^{3+}(\text{aq}) + \text{Co}(\text{s}) \rightarrow 2\text{Mn}^{2+}(\text{aq}) + \text{Co}^{2+}(\text{aq})$
- C.  $2\text{Mn}^{3+}(\text{aq}) + \text{Cd}(\text{s}) \rightarrow 2\text{Mn}^{2+}(\text{aq}) + \text{Cd}^{2+}(\text{aq})$
- D.  $2\text{Mn}^{2+}(\text{aq}) + \text{Co}^{2+}(\text{aq}) \rightarrow 2\text{Mn}^{3+}(\text{aq}) + \text{Co}(\text{s})$

**Question 12**

The sodium-sulfur cell shown below is a secondary galvanic cell with the overall cell reaction



The cell produces 2.1 volts.



The cell is to be recharged by connecting it to the power supply.


Which one of the following best describes the arrangement for recharging the cell?

- |    | <b>Power supply voltage</b> | <b>Connect terminals</b> |
|----|-----------------------------|--------------------------|
| A. | 2.1 volts                   | A to C and B to D        |
| B. | 2.1 volts                   | A to D and B to C        |
| C. | more than 2.1 volts         | A to C and B to D        |
| D. | more than 2.1 volts         | A to D and B to C        |

*Questions 13 and 14 refer to the following information.*

Iron pipes are used to transport natural gas to cities. Corrosion occurs when water droplets sit on the outer surface of the iron pipe.

Miniature galvanic cells are created, with regions such as those shown below, that act as anodes and cathodes.

**Question 13**

The type of region and reaction occurring at X in the cell is

- |    | <b>Region</b> | <b>Reaction</b>   |
|----|---------------|---|
| A. | anode         | $\text{Fe(s)} \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^{-}$                                  |
| B. | cathode       | $\text{Fe(s)} \rightarrow \text{Fe}^{2+}(\text{aq}) + 2\text{e}^{-}$                                  |
| C. | anode         | $\text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^{-} \rightarrow 4\text{OH}^{-}(\text{aq})$ |
| D. | cathode       | $\text{O}_2(\text{g}) + 2\text{H}_2\text{O(l)} + 4\text{e}^{-} \rightarrow 4\text{OH}^{-}(\text{aq})$ |

**Question 14**


Corrosion of an iron pipe can be prevented by connecting it to a magnesium bar buried in the ground. The magnesium corrodes in preference to the iron.

If the average current flowing between the two metals is  $2.0 \times 10^{-6}$  A, the amount of magnesium metal, in mol, reacting each second, would be

- A.  $1.0 \times 10^{-11}$
- B.  $2.1 \times 10^{-11}$
- C.  $4.1 \times 10^{-11}$
- D. 0.19

*Questions 15 and 16 refer to the following information.*

A fuel cell can be constructed that uses the following two half-reactions.

**Question 15**

Which one of the following would occur at the negative electrode of the cell as it generates electricity?

- A. production of  $\text{H}^+$
- B. formation of  $\text{H}_2\text{O}$
- C. consumption of  $\text{CO}_2$
- D. reduction of  $\text{CH}_3\text{OH}$


**Question 16**

Which one of the following statements about this fuel cell is most likely to be correct?

- A. An external power supply is used to recharge the cell.
- B. Gaseous products are recycled into the cell to improve efficiency.
- C. Chemical energy is not completely converted into electrical energy.
- D. More  $\text{H}^+$  ions are produced at the anode than are consumed at the cathode.

**Question 17**

Many reactions occurring in plant and animal cells involve a chemical called nicotinamide adenine dinucleotide,  $\text{NAD}^+$ . One such reaction is



It has been suggested that this reaction could be used in biochemical fuel cells to power pacemakers used to control irregular heartbeats.

If this reaction were performed in a fuel cell, NADH would

- A. undergo oxidation at the anode.
- B. undergo reduction at the cathode.
- C. undergo reduction at the anode.
- D. undergo oxidation at the cathode.

**Question 18**

Which one of the following describes the polarity of the anodes in electrolytic and galvanic cells?

- |    | <b>electrolytic cells</b> | <b>galvanic cells</b> |
|----|---------------------------|-----------------------|
| A. | positive                  | positive              |
| B. | positive                  | negative              |
| C. | negative                  | negative              |
| D. | negative                  | positive              |

**Question 19**

An aqueous solution containing a mixture of 1.0 M KI and 1.0 M CaBr<sub>2</sub> was electrolysed using unreactive electrodes.

Which one of the following reactions is most likely to occur at the anode?

- A.  $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightarrow \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$
- B.  $2\text{Br}^-(\text{aq}) \rightarrow \text{Br}_2(\text{aq}) + 2\text{e}^-$
- C.  $\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightarrow \text{Ca}(\text{s})$
- D.  $2\text{I}^-(\text{aq}) \rightarrow \text{I}_2(\text{aq}) + 2\text{e}^-$

**Question 20**

Lithium metal is manufactured by electrolysis of lithium salts.

Which of the following would be the best choice for the electrolyte and the anode in a commercial cell?

- |    | <b>electrolyte</b> | <b>anode</b> |
|----|--------------------|--------------|
| A. | LiCl solution      | iron rod     |
| B. | molten LiCl        | iron rod     |
| C. | LiCl solution      | carbon rod   |
| D. | molten LiCl        | carbon rod   |



**SECTION B – Short answer questions****Instructions for Section B**

Answer **all** questions in the spaces provided.

To obtain full marks for your responses you should

- give simplified answers with an appropriate number of significant figures to all numerical questions; unsimplified answers will not be given full marks.
- show all working in your answers to numerical questions. No credit will be given for an incorrect answer unless it is accompanied by details of the working.
- make sure chemical equations are balanced and that the formulas for individual substances include an indication of state; for example,  $\text{H}_2(\text{g})$ ;  $\text{NaCl}(\text{s})$

**Question 1**

- a. Use information from the electrochemical series in the Data Book to write a balanced overall equation that shows hydrogen peroxide,  $\text{H}_2\text{O}_2$ , reacting as a reductant.

---

---

---

2 marks

- b. Using data from the electrochemical series, a student suggests that a reaction will occur between  $\text{Cu}^{2+}$  ions and  $\text{H}_2$  gas. To test this prediction, hydrogen gas was bubbled into an aqueous solution of copper(II) sulfate,  $\text{CuSO}_4$ . No reaction was observed after 5 minutes. Provide one possible chemical reason that explains why the predicted reaction was not observed.

---


---

1 mark

Total 3 marks

**Question 2**

A 'QwikCure' pack, used to treat sporting injuries, contains a bag of water inside a larger bag of finely powdered ammonium nitrate,  $\text{NH}_4\text{NO}_3$ . Squeezing the pack causes the bag of water to break and the  $\text{NH}_4\text{NO}_3$  to dissolve. The change of energy that occurs can be used to treat an injury.



a. Suppose the activation energy of the **reverse reaction** is  $35 \text{ kJ mol}^{-1}$ .


i. Explain the meaning of the term 'activation energy'.

---

---

---


ii. On the graph below, sketch an energy profile diagram showing the changes that occur in chemical energy as the  $\text{NH}_4\text{NO}_3$  powder dissolves.



1 + 2 = 3 marks

- b. A chemist investigates the equilibrium reaction of ammonium ions with water. In this reaction the ammonium ion acts as a weak acid.
- i. Write an equation for the equilibrium reaction of ammonium ions with water.

While keeping the **temperature constant**, the chemist makes a change to a solution of ammonium ions in water that is initially at equilibrium. The following graph shows the effect of this change, which was made at time  $t_2$ , on the **rates** of the forward and back reactions.



- ii. What could have caused the change that occurred at time  $t_2$ ? Explain why the rate of the back reaction is affected by this change.

---



---



---



---



---

- iii. Would the value of the equilibrium constant at time  $t_3$  be **less than**, **equal to** or **greater than** the value of the equilibrium constant at time  $t_1$ ? Circle the correct response.

less than      equal to      greater than

1 + 2 + 1 = 4 marks

- c. The  $\text{NH}_4\text{NO}_3$  powder in a QwikCure pack dissolves completely to form 300 mL of solution, with a pH of 5.04.
- i. Write an expression for the acidity constant,  $K_a$ , for the reaction between ammonium ions and water.

- ii. Calculate the concentration, in  $\text{mol L}^{-1}$ , of  $\text{H}_3\text{O}^+$  ions in the 300 mL of solution.

---

---

---

- iii. Calculate the mass, in grams, of  $\text{NH}_4\text{NO}_3$  in the pack.

---

---

---

---

---

---

---

1 + 1 + 3 = 5 marks

Total 12 marks

**Question 3**

Dimethyl ether,  $\text{CH}_3\text{OCH}_3$ , is used as an environmentally friendly propellant in spray cans. It can be synthesised from methanol according to the following equation.



The equilibrium constant,  $K$ , for this reaction at  $350^\circ\text{C}$  is 5.74.

- a. Write an expression for  $K$  for this reaction.

1 mark

- b. Calculate the value of  $K$  at  $350^\circ\text{C}$  for the following reaction.




---



---

1 mark

- c. Methanol is pumped into an empty 20.0 L reactor vessel. At equilibrium the vessel contains 0.340 mol of methanol at  $350^\circ\text{C}$ .

- i. Calculate the concentration, in  $\text{mol L}^{-1}$ , of methanol at equilibrium.

---



---

- ii. Calculate the amount, in mol, of dimethyl ether present at equilibrium.

---



---



---



---



---



---

- iii. Calculate the amount, in mol, of methanol initially pumped into the reaction vessel.

---



---



---



---

1 + 2 + 2 = 5 marks

Total 7 marks

**Question 4**

Methyl palmitate,  $C_{17}H_{34}O_2$ , is a component of one type of biochemical fuel. It is a liquid at room temperature.

The molar enthalpy of combustion of methyl palmitate was determined using a bomb calorimeter.

The calorimeter was calibrated by passing a current of 4.40 amperes at a potential difference of 5.61 volts through an electric heater for 240 seconds. The temperature of the calorimeter rose by  $1.75^\circ\text{C}$ .

- a. Calculate the calibration factor of the calorimeter. Include the units of the calibration factor with your answer.

---

---

---

---

---

3 marks

A 0.529 g sample of methyl palmitate was then burned in excess oxygen in the calorimeter and the temperature rose by a further  $6.19^\circ\text{C}$ . The molar mass of methyl palmitate is  $270\text{ g mol}^{-1}$ .

- b. Calculate the amount of energy, in kJ, absorbed by the calorimeter when the sample of methyl palmitate was burned.

---

---

---

1 mark

- c. Calculate the amount of energy released, in kJ, by the combustion of 1.00 mol of methyl palmitate.

---

---


---

---

---

2 marks

- d. The balanced equation for the combustion of liquid methyl palmitate in excess oxygen is



Write the value of  $\Delta H$  for this reaction, in  $\text{kJ mol}^{-1}$ .

---

2 marks

Most of Victoria's electricity is generated by burning fossil fuels such as coal and natural gas. Alternative methods of generating electricity are currently being developed.

- e. Biochemical fuels are an alternative fuel for generating electricity.
- Name one biochemical fuel, other than methyl palmitate, and the raw material used in its production.

Biochemical fuel \_\_\_\_\_

Raw material used in its production \_\_\_\_\_

- Identify **one** disadvantage or limitation of the use of this biochemical fuel for the large-scale generation of electricity.

\_\_\_\_\_

\_\_\_\_\_

2 + 1 = 3 marks

- f. Some countries rely on nuclear fission for the large-scale production of electricity.

- State one advantage of using nuclear fission.

\_\_\_\_\_

- State one disadvantage of using nuclear fission.

\_\_\_\_\_


1 + 1 = 2 marks

Total 13 marks


**Question 5**

A vanadium redox battery is used to store electrical energy generated at a wind farm in Tasmania. The battery supplies electricity to the power grid as required through a control switch.

The diagram below shows the structure of a cell in a vanadium redox battery. The reactants are dissolved in an acidic solution, stored in large tanks and pumped through the cell. The cell is recharged using electricity generated by the wind turbines. A polymer membrane allows the movement of particular ions.



The two relevant half-equations for the vanadium redox battery are



- a. State the polarity of each electrode as the battery is discharged.

Electrode A \_\_\_\_\_

Electrode B \_\_\_\_\_

1 mark

- b. Circle the vanadium-containing ion that would have the highest concentration at the anode when the cell is **fully charged**.



1 mark



- c. Write a balanced overall equation for the reaction that occurs when the cell is being **recharged**.

---

---

1 mark

- d. Compare the vanadium redox cell to a fuel cell by describing **one** major way in which they differ.

---

---

---

---

1 mark

- e. Write a balanced overall equation to show why iron would be an unsuitable material to use as electrode B in the vanadium redox cell.

---

---


---

1 mark

Total 5 marks

**Question 6**

A particular industrial process involves the following steps.



- a. It is possible to alter the temperature and pressure at which reaction 2 occurs. In the table below, indicate what effect the following changes to temperature and pressure would have on the **rate**, **equilibrium yield** and value of the **equilibrium constant,  $K$** , for reaction 2.

|   | Would the <b>rate</b> of reaction 2 become <b>higher, lower</b> or <b>remain unchanged</b> ? | Would the <b>equilibrium yield</b> of reaction 2 become <b>higher, lower</b> or <b>remain unchanged</b> ? | Would the <b>value of the equilibrium constant, <math>K</math></b> , of reaction 2 become <b>higher, lower</b> or <b>remain unchanged</b> ? |
|---|--|---|---|
| The temperature of reaction 2 is lowered to 150°C.  |  |   |   |
| The pressure of reaction 2 is increased to 5 atm by pumping more B(g) and C(g) into the reaction vessel, at constant temperature. |  |   |   |

6 marks

- b. Heat energy is released by reaction 2. Describe how the heat energy could be used within this industrial process.

---



---



---

1 mark

- c. During this semester you have studied the production of one of the following chemicals.  
Circle the chemical you have studied in detail this semester.

ammonia      ethene      sulfuric acid      nitric acid

- i. Describe one waste management strategy, other than recycling heat, employed in the industrial production of your selected chemical.

---






---



---

- ii. The following table includes a selection of HAZCHEM labels used to identify dangerous goods.

|   |   |  |   |
|---|---|--|---|
|  |  |  |  |
|  |  |  |  |

Circle one label that could be used to identify the hazardous nature of your selected chemical.

- iii. State two uses of your selected chemical.

Use 1 \_\_\_\_\_

Use 2 \_\_\_\_\_

1 + 1 + 2 = 4 marks

Total 11 marks

**Question 7**

A classroom experiment was set up to simulate the industrial extraction of zinc metal from an aqueous solution of zinc ions by electrolysis. In this experiment 150 mL of 1.00 M  $\text{ZnSO}_4$  solution was electrolysed at  $25^\circ\text{C}$  using inert carbon electrodes.

- a. Write a half-equation for the oxidation reaction.

---

1 mark

- b. A mass of 0.900 g of zinc is produced in 30.0 minutes.

Calculate the electric current, in A, supplied to the cell during the electrolysis. Express your answer to an appropriate number of significant figures.

---

---

---

---

---

---

---

---

---

4 marks

Total 5 marks



[www.theallpapers.com](http://www.theallpapers.com)



**Victorian Certificate of Education  
2009**

**CHEMISTRY**  
**Written examination**

**Thursday 12 November 2009**

**Reading time: 9.00 am to 9.15 am (15 minutes)**

**Writing time: 9.15 am to 10.45 am (1 hour 30 minutes)**

**DATA BOOK**

**Directions to students**

- A question and answer book is provided with this data book.

**Students are NOT permitted to bring mobile phones and/or any other unauthorised electronic devices into the examination room.**

**Table of contents**

|  | page |
|--|------|
| 1. Periodic table of the elements  | 3    |
| 2. The electrochemical series  | 4    |
| 3. Physical constants  | 5    |
| 4. SI prefixes, their symbols and values   | 5    |
| 5. $^1\text{H}$ NMR data   | 5–6  |
| 6. $^{13}\text{C}$ NMR data  | 7    |
| 7. Infrared absorption data  | 7    |
| 8. 2-amino acids ( $\alpha$ -amino acids)  | 8–9  |
| 9. Formulas of some fatty acids  | 10   |
| 10. Structural formulas of some important biomolecules                                 | 10   |
| 11. Acid-base indicators   | 11   |
| 12. Acidity constants, $K_a$ , of some weak acids                                      | 11   |
| 13. Values of molar enthalpy of combustion of some common fuels at 298 K and 101.3 kPa | 11   |

## 1. Periodic table of the elements

| 1<br><b>H</b><br>Hydrogen<br>1.0          |  | 2<br><b>He</b><br>Helium<br>4.0           |  |
|---|--|---|--|
| 3<br><b>Li</b><br>Lithium<br>6.9          |  | 4<br><b>Be</b><br>Beryllium<br>9.0        |  |
| 11<br><b>Na</b><br>Sodium<br>23.0         |  | 12<br><b>Mg</b><br>Magnesium<br>24.3      |  |
| 19<br><b>K</b><br>Potassium<br>39.1       |  | 20<br><b>Ca</b><br>Calcium<br>40.1        |  |
| 37<br><b>Rb</b><br>Rubidium<br>85.5       |  | 38<br><b>Sr</b><br>Strontium<br>87.6      |  |
| 55<br><b>Cs</b><br>Caesium<br>132.9       |  | 56<br><b>Ba</b><br>Barium<br>137.3        |  |
| 87<br><b>Fr</b><br>Francium<br>(223)      |  | 88<br><b>Ra</b><br>Radium<br>(226)        |  |
| 21<br><b>Sc</b><br>Scandium<br>44.9       |  | 22<br><b>Ti</b><br>Titanium<br>47.9       |  |
| 39<br><b>Y</b><br>Yttrium<br>88.9         |  | 40<br><b>Zr</b><br>Zirconium<br>91.2      |  |
| 71<br><b>La</b><br>Lanthanum<br>138.9     |  | 72<br><b>Hf</b><br>Hafnium<br>178.5       |  |
| 89<br><b>Ac</b><br>Actinium<br>(227)      |  | 90<br><b>Rf</b><br>Rutherfordium<br>(261) |  |
| 23<br><b>V</b><br>Vanadium<br>50.9        |  | 24<br><b>Cr</b><br>Chromium<br>52.0       |  |
| 41<br><b>Nb</b><br>Niobium<br>92.9        |  | 42<br><b>Mo</b><br>Molybdenum<br>95.9     |  |
| 73<br><b>Ta</b><br>Tantalum<br>180.9      |  | 74<br><b>W</b><br>Tungsten<br>183.8       |  |
| 105<br><b>Db</b><br>Dubnium<br>(262)      |  | 106<br><b>Sg</b><br>Seaborgium<br>(266)   |  |
| 25<br><b>Mn</b><br>Manganese<br>54.9      |  | 26<br><b>Fe</b><br>Iron<br>55.9           |  |
| 43<br><b>Tc</b><br>Technetium<br>98.1     |  | 44<br><b>Ru</b><br>Ruthenium<br>101.1     |  |
| 75<br><b>Re</b><br>Rhenium<br>186.2       |  | 76<br><b>Os</b><br>Osmium<br>190.2        |  |
| 107<br><b>Bh</b><br>Bohrium<br>(264)      |  | 108<br><b>Hs</b><br>Hassium<br>(277)      |  |
| 27<br><b>Co</b><br>Cobalt<br>58.9         |  | 28<br><b>Ni</b><br>Nickel<br>58.7         |  |
| 45<br><b>Rh</b><br>Rhodium<br>102.9       |  | 46<br><b>Pd</b><br>Palladium<br>106.4     |  |
| 77<br><b>Ir</b><br>Iridium<br>192.2       |  | 78<br><b>Pt</b><br>Platinum<br>195.1      |  |
| 109<br><b>Mt</b><br>Meitnerium<br>(268)   |  | 110<br><b>Ds</b><br>Darmstadtium<br>(271) |  |
| 29<br><b>Cu</b><br>Copper<br>63.6         |  | 30<br><b>Zn</b><br>Zinc<br>65.4           |  |
| 47<br><b>Ag</b><br>Silver<br>107.9        |  | 48<br><b>Cd</b><br>Cadmium<br>112.4       |  |
| 79<br><b>Au</b><br>Gold<br>197.0          |  | 80<br><b>Hg</b><br>Mercury<br>200.6       |  |
| 111<br><b>Rg</b><br>Roentgenium<br>(272)  |  | 112<br><b>Uub</b><br>Ununbium<br>(271)    |  |
| 13<br><b>Al</b><br>Aluminium<br>27.0      |  | 14<br><b>Si</b><br>Silicon<br>28.1        |  |
| 31<br><b>Ga</b><br>Gallium<br>69.7        |  | 32<br><b>Ge</b><br>Germanium<br>72.6      |  |
| 49<br><b>In</b><br>Indium<br>114.8        |  | 50<br><b>Sn</b><br>Tin<br>118.7           |  |
| 81<br><b>Tl</b><br>Thallium<br>204.4      |  | 82<br><b>Pb</b><br>Lead<br>207.2          |  |
| 113<br><b>Uut</b><br>Ununtrium<br>(273)   |  | 114<br><b>Uuq</b><br>Ununquadium<br>(274) |  |
| 15<br><b>P</b><br>Phosphorus<br>31.0      |  | 16<br><b>S</b><br>Sulfur<br>32.1          |  |
| 33<br><b>As</b><br>Arsenic<br>74.9        |  | 34<br><b>Se</b><br>Selenium<br>79.0       |  |
| 51<br><b>Sb</b><br>Antimony<br>121.8      |  | 52<br><b>Te</b><br>Tellurium<br>127.6     |  |
| 83<br><b>Bi</b><br>Bismuth<br>209.0       |  | 84<br><b>Po</b><br>Polonium<br>(209)      |  |
| 115<br><b>Uup</b><br>Ununpentium<br>(275) |  | 116<br><b>Uuh</b><br>Ununhexium<br>(276)  |  |
| 7<br><b>N</b><br>Nitrogen<br>14.0         |  | 8<br><b>O</b><br>Oxygen<br>16.0           |  |
| 15<br><b>P</b><br>Phosphorus<br>31.0      |  | 16<br><b>S</b><br>Sulfur<br>32.1          |  |
| 33<br><b>As</b><br>Arsenic<br>74.9        |  | 34<br><b>Se</b><br>Selenium<br>79.0       |  |
| 51<br><b>Sb</b><br>Antimony<br>121.8      |  | 52<br><b>Te</b><br>Tellurium<br>127.6     |  |
| 83<br><b>Bi</b><br>Bismuth<br>209.0       |  | 84<br><b>Po</b><br>Polonium<br>(209)      |  |
| 115<br><b>Uup</b><br>Ununpentium<br>(275) |  | 116<br><b>Uuh</b><br>Ununhexium<br>(276)  |  |
| 6<br><b>C</b><br>Carbon<br>12.0           |  | 7<br><b>N</b><br>Nitrogen<br>14.0         |  |
| 14<br><b>Si</b><br>Silicon<br>28.1        |  | 15<br><b>P</b><br>Phosphorus<br>31.0      |  |
| 32<br><b>Ge</b><br>Germanium<br>72.6      |  | 33<br><b>As</b><br>Arsenic<br>74.9        |  |
| 50<br><b>Sn</b><br>Tin<br>118.7           |  | 51<br><b>Sb</b><br>Antimony<br>121.8      |  |
| 82<br><b>Pb</b><br>Lead<br>207.2          |  | 83<br><b>Bi</b><br>Bismuth<br>209.0       |  |
| 114<br><b>Uuq</b><br>Ununquadium<br>(274) |  | 115<br><b>Uup</b><br>Ununpentium<br>(275) |  |
| 9<br><b>F</b><br>Fluorine<br>19.0         |  | 10<br><b>Ne</b><br>Neon<br>20.1           |  |
| 17<br><b>Cl</b><br>Chlorine<br>35.5       |  | 18<br><b>Ar</b><br>Argon<br>39.9          |  |
| 35<br><b>Br</b><br>Bromine<br>79.9        |  | 36<br><b>Kr</b><br>Krypton<br>83.8        |  |
| 53<br><b>I</b><br>Iodine<br>126.9         |  | 54<br><b>Xe</b><br>Xenon<br>131.3         |  |
| 85<br><b>At</b><br>Astatine<br>(210)      |  | 86<br><b>Rn</b><br>Radon<br>(222)         |  |
| 117<br><b>Uue</b><br>Ununseptium<br>(277) |  | 118<br><b>Uuo</b><br>Ununoctium<br>(278)  |  |

|   |   |   |   |
|---|---|---|---|
| 68<br><b>Er</b><br>Erbium<br>167.3        | 69<br><b>Tm</b><br>Thulium<br>168.9       | 70<br><b>Yb</b><br>Ytterbium<br>173.0     | 71<br><b>Lu</b><br>Lutetium<br>175.0      |
| 99<br><b>Es</b><br>Einsteinium<br>(252)   | 100<br><b>Fm</b><br>Fermium<br>(257)      | 101<br><b>Md</b><br>Mendelevium<br>(258)  | 102<br><b>No</b><br>Nobelium<br>(259)     |
| 112<br><b>Ubn</b><br>Unbinilium<br>(272)  | 113<br><b>Ubt</b><br>Unbinilium<br>(273)  | 114<br><b>Ubu</b><br>Unbinilium<br>(274)  | 115<br><b>Ubu</b><br>Unbinilium<br>(275)  |
| 67<br><b>Ho</b><br>Holmium<br>164.9       | 68<br><b>Er</b><br>Erbium<br>167.3        | 69<br><b>Tm</b><br>Thulium<br>168.9       | 70<br><b>Yb</b><br>Ytterbium<br>173.0     |
| 98<br><b>Cf</b><br>Californium<br>(251)   | 99<br><b>Es</b><br>Einsteinium<br>(252)   | 100<br><b>Fm</b><br>Fermium<br>(257)      | 101<br><b>Md</b><br>Mendelevium<br>(258)  |
| 110<br><b>Ds</b><br>Darmstadtium<br>(271) | 111<br><b>Rg</b><br>Roentgenium<br>(272)  | 112<br><b>Uub</b><br>Ununbium<br>(271)    | 113<br><b>Uut</b><br>Ununtrium<br>(273)   |
| 65<br><b>Tb</b><br>Terbium<br>158.9       | 66<br><b>Dy</b><br>Dysprosium<br>162.5    | 67<br><b>Ho</b><br>Holmium<br>164.9       | 68<br><b>Er</b><br>Erbium<br>167.3        |
| 97<br><b>Bk</b><br>Berkelium<br>(247)     | 98<br><b>Cf</b><br>Californium<br>(251)   | 99<br><b>Es</b><br>Einsteinium<br>(252)   | 100<br><b>Fm</b><br>Fermium<br>(257)      |
| 109<br><b>Mt</b><br>Meitnerium<br>(268)   | 110<br><b>Ds</b><br>Darmstadtium<br>(271) | 111<br><b>Rg</b><br>Roentgenium<br>(272)  | 112<br><b>Uub</b><br>Ununbium<br>(271)    |
| 64<br><b>Gd</b><br>Gadolinium<br>157.2    | 65<br><b>Tb</b><br>Terbium<br>158.9       | 66<br><b>Dy</b><br>Dysprosium<br>162.5    | 67<br><b>Ho</b><br>Holmium<br>164.9       |
| 96<br><b>Cm</b><br>Curium<br>(247)        | 97<br><b>Bk</b><br>Berkelium<br>(247)     | 98<br><b>Cf</b><br>Californium<br>(251)   | 99<br><b>Es</b><br>Einsteinium<br>(252)   |
| 108<br><b>Hs</b><br>Hassium<br>(277)      | 109<br><b>Mt</b><br>Meitnerium<br>(268)   | 110<br><b>Ds</b><br>Darmstadtium<br>(271) | 111<br><b>Rg</b><br>Roentgenium<br>(272)  |
| 63<br><b>Eu</b><br>Europium<br>152.0      | 64<br><b>Gd</b><br>Gadolinium<br>157.2    | 65<br><b>Tb</b><br>Terbium<br>158.9       | 66<br><b>Dy</b><br>Dysprosium<br>162.5    |
| 95<br><b>Am</b><br>Americium<br>(243)     | 96<br><b>Cm</b><br>Curium<br>(247)        | 97<br><b>Bk</b><br>Berkelium<br>(247)     | 98<br><b>Cf</b><br>Californium<br>(251)   |
| 107<br><b>Bh</b><br>Bohrium<br>(264)      | 108<br><b>Hs</b><br>Hassium<br>(277)      | 109<br><b>Mt</b><br>Meitnerium<br>(268)   | 110<br><b>Ds</b><br>Darmstadtium<br>(271) |
| 62<br><b>Sm</b><br>Samarium<br>150.3      | 63<br><b>Eu</b><br>Europium<br>152.0      | 64<br><b>Gd</b><br>Gadolinium<br>157.2    | 65<br><b>Tb</b><br>Terbium<br>158.9       |
| 94<br><b>Pu</b><br>Plutonium<br>(244)     | 95<br><b>Am</b><br>Americium<br>(243)     | 96<br><b>Cm</b><br>Curium<br>(247)        | 97<br><b>Bk</b><br>Berkelium<br>(247)     |
| 106<br><b>Sg</b><br>Seaborgium<br>(266)   | 107<br><b>Bh</b><br>Bohrium<br>(264)      | 108<br><b>Hs</b><br>Hassium<br>(277)      | 109<br><b>Mt</b><br>Meitnerium<br>(268)   |
| 61<br><b>Pm</b><br>Promethium<br>(145)    | 62<br><b>Sm</b><br>Samarium<br>150.3      | 63<br><b>Eu</b><br>Europium<br>152.0      | 64<br><b>Gd</b><br>Gadolinium<br>157.2    |
| 93<br><b>Np</b><br>Neptunium<br>(237.1)   | 94<br><b>Pu</b><br>Plutonium<br>(244)     | 95<br><b>Am</b><br>Americium<br>(243)     | 96<br><b>Cm</b><br>Curium<br>(247)        |
| 85<br><b>At</b><br>Astatine<br>(210)      | 86<br><b>Rn</b><br>Radon<br>(222)         | 87<br><b>Fr</b><br>Francium<br>(223)      | 88<br><b>Ra</b><br>Radium<br>(226)        |
| 116<br><b>Uuh</b><br>Ununhexium<br>(276)  | 117<br><b>Uue</b><br>Ununseptium<br>(277) | 118<br><b>Uuo</b><br>Ununoctium<br>(278)  | 119<br><b>Uuq</b><br>Ununquadium<br>(279) |

|                                     |  |                                       |   |                                       |                                       |  |                                       |   |   |                                      |  |                                       |   |
|-------------------------------------|--|---------------------------------------|---|---------------------------------------|---------------------------------------|--|---------------------------------------|---|---|--------------------------------------|--|---------------------------------------|---|
| 58<br><b>Ce</b><br>Cerium<br>140.1  | 59<br><b>Pr</b><br>Praseodymium<br>140.9 | 60<br><b>Nd</b><br>Neodymium<br>144.2 | 61<br><b>Pm</b><br>Promethium<br>(145)  | 62<br><b>Sm</b><br>Samarium<br>150.3  | 63<br><b>Eu</b><br>Europium<br>152.0  | 64<br><b>Gd</b><br>Gadolinium<br>157.2 | 65<br><b>Tb</b><br>Terbium<br>158.9   | 66<br><b>Dy</b><br>Dysprosium<br>162.5  | 67<br><b>Ho</b><br>Holmium<br>164.9     | 68<br><b>Er</b><br>Erbium<br>167.3   | 69<br><b>Tm</b><br>Thulium<br>168.9      | 70<br><b>Yb</b><br>Ytterbium<br>173.0 | 71<br><b>Lu</b><br>Lutetium<br>175.0    |
| 90<br><b>Th</b><br>Thorium<br>232.0 | 91<br><b>Pa</b><br>Protactinium<br>231.0 | 92<br><b>U</b><br>Uranium<br>238.0    | 93<br><b>Np</b><br>Neptunium<br>(237.1) | 94<br><b>Pu</b><br>Plutonium<br>(244) | 95<br><b>Am</b><br>Americium<br>(243) | 96<br><b>Cm</b><br>Curium<br>(247)     | 97<br><b>Bk</b><br>Berkelium<br>(247) | 98<br><b>Cf</b><br>Californium<br>(251) | 99<br><b>Es</b><br>Einsteinium<br>(252) | 100<br><b>Fm</b><br>Fermium<br>(257) | 101<br><b>Md</b><br>Mendelevium<br>(258) | 102<br><b>No</b><br>Nobelium<br>(259) | 103<br><b>Lr</b><br>Lawrencium<br>(262) |

TURN OVER



**2. The electrochemical series**

|   | $E^\circ$ in volt |
|---|-------------------|
| $\text{F}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{F}^-(\text{aq})$  | +2.87             |
| $\text{H}_2\text{O}_2(\text{aq}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$ | +1.77             |
| $\text{Au}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Au}(\text{s})$  | +1.68             |
| $\text{Cl}_2(\text{g}) + 2\text{e}^- \rightleftharpoons 2\text{Cl}^-(\text{aq})$  | +1.36             |
| $\text{O}_2(\text{g}) + 4\text{H}^+(\text{aq}) + 4\text{e}^- \rightleftharpoons 2\text{H}_2\text{O}(\text{l})$            | +1.23             |
| $\text{Br}_2(\text{l}) + 2\text{e}^- \rightleftharpoons 2\text{Br}^-(\text{aq})$  | +1.09             |
| $\text{Ag}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Ag}(\text{s})$  | +0.80             |
| $\text{Fe}^{3+}(\text{aq}) + \text{e}^- \rightleftharpoons \text{Fe}^{2+}(\text{aq})$                                     | +0.77             |
| $\text{O}_2(\text{g}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{O}_2(\text{aq})$          | +0.68             |
| $\text{I}_2(\text{s}) + 2\text{e}^- \rightleftharpoons 2\text{I}^-(\text{aq})$  | +0.54             |
| $\text{O}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l}) + 4\text{e}^- \rightleftharpoons 4\text{OH}^-(\text{aq})$           | +0.40             |
| $\text{Cu}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Cu}(\text{s})$  | +0.34             |
| $\text{Sn}^{4+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}^{2+}(\text{aq})$                                    | +0.15             |
| $\text{S}(\text{s}) + 2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2\text{S}(\text{g})$               | +0.14             |
| $2\text{H}^+(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g})$  | 0.00              |
| $\text{Pb}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Pb}(\text{s})$  | -0.13             |
| $\text{Sn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Sn}(\text{s})$  | -0.14             |
| $\text{Ni}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ni}(\text{s})$  | -0.23             |
| $\text{Co}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Co}(\text{s})$  | -0.28             |
| $\text{Fe}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Fe}(\text{s})$  | -0.44             |
| $\text{Zn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Zn}(\text{s})$  | -0.76             |
| $2\text{H}_2\text{O}(\text{l}) + 2\text{e}^- \rightleftharpoons \text{H}_2(\text{g}) + 2\text{OH}^-(\text{aq})$           | -0.83             |
| $\text{Mn}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mn}(\text{s})$  | -1.03             |
| $\text{Al}^{3+}(\text{aq}) + 3\text{e}^- \rightleftharpoons \text{Al}(\text{s})$  | -1.67             |
| $\text{Mg}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Mg}(\text{s})$  | -2.34             |
| $\text{Na}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Na}(\text{s})$  | -2.71             |
| $\text{Ca}^{2+}(\text{aq}) + 2\text{e}^- \rightleftharpoons \text{Ca}(\text{s})$  | -2.87             |
| $\text{K}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{K}(\text{s})$  | -2.93             |
| $\text{Li}^+(\text{aq}) + \text{e}^- \rightleftharpoons \text{Li}(\text{s})$  | -3.02             |

### 3. Physical constants

Avogadro's constant ( $N_A$ ) =  $6.02 \times 10^{23} \text{ mol}^{-1}$

Charge on one electron =  $-1.60 \times 10^{-19} \text{ C}$

Faraday constant ( $F$ ) =  $96\,500 \text{ C mol}^{-1}$

Gas constant ( $R$ ) =  $8.31 \text{ J K}^{-1} \text{ mol}^{-1}$

Ionic product for water ( $K_w$ ) =  $1.00 \times 10^{-14} \text{ mol}^2 \text{ L}^{-2}$  at 298 K  
(Self ionisation constant)

Molar volume ( $V_m$ ) of an ideal gas at 273 K, 101.3 kPa (STP) =  $22.4 \text{ L mol}^{-1}$

Molar volume ( $V_m$ ) of an ideal gas at 298 K, 101.3 kPa (SLC) =  $24.5 \text{ L mol}^{-1}$

Specific heat capacity ( $c$ ) of water =  $4.18 \text{ J g}^{-1} \text{ K}^{-1}$

Density ( $d$ ) of water at 25°C =  $1.00 \text{ g mL}^{-1}$

1 atm = 101.3 kPa = 760 mm Hg

0°C = 273 K

### 4. SI prefixes, their symbols and values

| SI prefix | Symbol | Value      |
|-----------|--------|------------|
| giga      | G      | $10^9$     |
| mega      | M      | $10^6$     |
| kilo      | k      | $10^3$     |
| deci      | d      | $10^{-1}$  |
| centi     | c      | $10^{-2}$  |
| milli     | m      | $10^{-3}$  |
| micro     | $\mu$  | $10^{-6}$  |
| nano      | n      | $10^{-9}$  |
| pico      | p      | $10^{-12}$ |




### 5. $^1\text{H}$ NMR data

Typical proton shift values relative to TMS = 0

These can differ slightly in different solvents. Where more than one proton environment is shown in the formula, the shift refers to the ones in bold letters.

| Type of proton  | Chemical shift (ppm) |
|---|----------------------|
| R-CH <sub>3</sub>   | 0.9                  |
| R-CH <sub>2</sub> -R  | 1.3                  |
| RCH = CH- <b>CH<sub>3</sub></b>   | 1.7                  |
| R <sub>3</sub> -CH  | 2.0                  |
| $\text{CH}_3-\text{C} \begin{array}{l} \text{O} \\ \parallel \\ \text{OR} \end{array}$ or $\text{CH}_3-\text{C} \begin{array}{l} \text{O} \\ \parallel \\ \text{NHR} \end{array}$ | 2.0                  |

TURN OVER

| Type of proton   | Chemical shift (ppm)                                 |
|--|--|
| $\begin{array}{c} \text{R} \quad \text{CH}_3 \\ \quad \diagdown \quad / \\ \quad \text{C} \\ \quad    \\ \quad \text{O} \end{array}$ | 2.1  |
| R-CH <sub>2</sub> -X (X = F, Cl, Br or I)  | 3-4  |
| R-CH <sub>2</sub> -OH  | 3.6  |
| $\begin{array}{c} \quad \text{O} \\ \quad // \\ \text{R}-\text{C} \\ \quad \backslash \\ \quad \text{NHCH}_2\text{R} \end{array}$    | 3.2  |
| R-O-CH <sub>3</sub> or R-O-CH <sub>2</sub> R   | 3.3  |
|   | 2.3  |
| $\begin{array}{c} \quad \text{O} \\ \quad // \\ \text{R}-\text{C} \\ \quad \backslash \\ \quad \text{OCH}_2\text{R} \end{array}$     | 4.1  |
| R-O-H  | 1-6 (varies considerably under different conditions) |
| R-NH <sub>2</sub>  | 1-5  |
| RHC = CH <sub>2</sub>  | 4.6-6.0  |
|   | 7.0  |
|   | 7.3  |
| $\begin{array}{c} \quad \text{O} \\ \quad // \\ \text{R}-\text{C} \\ \quad \backslash \\ \quad \text{NHCH}_2\text{R} \end{array}$    | 8.1  |
| $\begin{array}{c} \quad \text{O} \\ \quad // \\ \text{R}-\text{C} \\ \quad \backslash \\ \quad \text{H} \end{array}$                 | 9-10   |
| $\begin{array}{c} \quad \text{O} \\ \quad // \\ \text{R}-\text{C} \\ \quad \backslash \\ \quad \text{O}-\text{H} \end{array}$        | 11.5   |

**6.  $^{13}\text{C}$  NMR data**

| Type of carbon                   | Chemical shift (ppm) |
|----------------------------------|----------------------|
| R-CH <sub>3</sub>                | 8–25                 |
| R-CH <sub>2</sub> -R             | 20–45                |
| R <sub>3</sub> -CH               | 40–60                |
| R <sub>4</sub> -C                | 36–45                |
| R-CH <sub>2</sub> -X             | 15–80                |
| R <sub>3</sub> C-NH <sub>2</sub> | 35–70                |
| R-CH <sub>2</sub> -OH            | 50–90                |
| RC≡CR                            | 75–95                |
| R <sub>2</sub> C=CR <sub>2</sub> | 110–150              |
| RCOOH                            | 160–185              |

**7. Infrared absorption data**

Characteristic range for infrared absorption


| Bond                 | Wave number (cm <sup>-1</sup> ) |
|----------------------|---------------------------------|
| C-Cl                 | 700–800                         |
| C-C                  | 750–1100                        |
| C-O                  | 1000–1300                       |
| C=C                  | 1610–1680                       |
| C=O                  | 1670–1750                       |
| O-H (acids)          | 2500–3300                       |
| C-H                  | 2850–3300                       |
| O-H (alcohols)       | 3200–3550                       |
| N-H (primary amines) | 3350–3500                       |

TURN OVER

[www.theallpapers.com](http://www.theallpapers.com)

**8. 2-amino acids ( $\alpha$ -amino acids)**

| Name          | Symbol | Structure   |
|---------------|--------|---|
| alanine       | Ala    | $\begin{array}{c} \text{CH}_3 \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$   |
| arginine      | Arg    | $\begin{array}{c} \text{NH} \\    \\ \text{CH}_2-\text{CH}_2-\text{CH}_2-\text{NH}-\text{C}-\text{NH}_2 \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$ |
| asparagine    | Asn    | $\begin{array}{c} \text{O} \\    \\ \text{CH}_2-\text{C}-\text{NH}_2 \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$                                    |
| aspartic acid | Asp    | $\begin{array}{c} \text{CH}_2-\text{COOH} \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$   |
| cysteine      | Cys    | $\begin{array}{c} \text{CH}_2-\text{SH} \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$   |
| glutamine     | Gln    | $\begin{array}{c} \text{O} \\    \\ \text{CH}_2-\text{CH}_2-\text{C}-\text{NH}_2 \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$                        |
| glutamic acid | Glu    | $\begin{array}{c} \text{CH}_2-\text{CH}_2-\text{COOH} \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$   |
| glycine       | Gly    | $\text{H}_2\text{N}-\text{CH}_2-\text{COOH}$  |
| histidine     | His    | $\begin{array}{c} \text{N} \\ // \quad \backslash \\ \text{CH}_2-\text{C} \quad \text{N}-\text{H} \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$       |
| isoleucine    | Ile    | $\begin{array}{c} \text{CH}_3-\text{CH}-\text{CH}_2-\text{CH}_3 \\   \\ \text{H}_2\text{N}-\text{CH}-\text{COOH} \end{array}$   |


| Name          | Symbol | Structure   |
|---------------|--------|---|
| leucine       | Leu    | $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\   \\ \text{CH}_2 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$           |
| lysine        | Lys    | $\begin{array}{c} \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{CH}_2 - \text{NH}_2 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$ |
| methionine    | Met    | $\begin{array}{c} \text{CH}_2 - \text{CH}_2 - \text{S} - \text{CH}_3 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                  |
| phenylalanine | Phe    | $\begin{array}{c} \text{CH}_2 - \text{C}_6\text{H}_5 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                                  |
| proline       | Pro    |   |
| serine        | Ser    | $\begin{array}{c} \text{CH}_2 - \text{OH} \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$   |
| threonine     | Thr    | $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{OH} \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                                 |
| tryptophan    | Trp    | $\begin{array}{c} \text{CH}_2 - \text{C}_8\text{H}_6\text{N}_2 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                        |
| tyrosine      | Tyr    | $\begin{array}{c} \text{CH}_2 - \text{C}_6\text{H}_4 - \text{OH} \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                      |
| valine        | Val    | $\begin{array}{c} \text{CH}_3 - \text{CH} - \text{CH}_3 \\   \\ \text{H}_2\text{N} - \text{CH} - \text{COOH} \end{array}$                               |

TURN OVER


### 9. Formulas of some fatty acids

| Name        | Formula            |
|-------------|--------------------|
| Lauric      | $C_{11}H_{23}COOH$ |
| Myristic    | $C_{13}H_{27}COOH$ |
| Palmitic    | $C_{15}H_{31}COOH$ |
| Palmitoleic | $C_{15}H_{29}COOH$ |
| Stearic     | $C_{17}H_{35}COOH$ |
| Oleic       | $C_{17}H_{33}COOH$ |
| Linoleic    | $C_{17}H_{31}COOH$ |
| Linolenic   | $C_{17}H_{29}COOH$ |
| Arachidic   | $C_{19}H_{39}COOH$ |
| Arachidonic | $C_{19}H_{31}COOH$ |


### 10. Structural formulas of some important biomolecules




sucrose




glycerol




deoxyribose




adenine




guanine



cytosine



thymine



phosphate

**11. Acid-base indicators**

| Name             | pH range | Colour change |        | $K_a$               |
|------------------|----------|---------------|--------|---------------------|
|                  |          | Acid          | Base   |                     |
| Thymol blue      | 1.2–2.8  | red           | yellow | $2 \times 10^{-2}$  |
| Methyl orange    | 3.1–4.4  | red           | yellow | $2 \times 10^{-4}$  |
| Bromophenol blue | 3.0–4.6  | yellow        | blue   | $6 \times 10^{-5}$  |
| Methyl red       | 4.2–6.3  | red           | yellow | $8 \times 10^{-6}$  |
| Bromothymol blue | 6.0–7.6  | yellow        | blue   | $1 \times 10^{-7}$  |
| Phenol red       | 6.8–8.4  | yellow        | red    | $1 \times 10^{-8}$  |
| Phenolphthalein  | 8.3–10.0 | colourless    | red    | $5 \times 10^{-10}$ |

**12. Acidity constants,  $K_a$ , of some weak acids**

| Name         | Formula                           | $K_a$                 |
|--------------|-----------------------------------|-----------------------|
| Ammonium ion | $\text{NH}_4^+$                   | $5.6 \times 10^{-10}$ |
| Benzoic      | $\text{C}_6\text{H}_5\text{COOH}$ | $6.4 \times 10^{-5}$  |
| Boric        | $\text{H}_3\text{BO}_3$           | $5.8 \times 10^{-10}$ |
| Ethanoic     | $\text{CH}_3\text{COOH}$          | $1.7 \times 10^{-5}$  |
| Hydrocyanic  | $\text{HCN}$                      | $6.3 \times 10^{-10}$ |
| Hydrofluoric | $\text{HF}$                       | $7.6 \times 10^{-4}$  |
| Hypobromous  | $\text{HOBr}$                     | $2.4 \times 10^{-9}$  |
| Hypochlorous | $\text{HOCl}$                     | $2.9 \times 10^{-8}$  |
| Lactic       | $\text{HC}_3\text{H}_5\text{O}_3$ | $1.4 \times 10^{-4}$  |
| Methanoic    | $\text{HCOOH}$                    | $1.8 \times 10^{-4}$  |
| Nitrous      | $\text{HNO}_2$                    | $7.2 \times 10^{-4}$  |
| Propanoic    | $\text{C}_2\text{H}_5\text{COOH}$ | $1.3 \times 10^{-5}$  |

**13. Values of molar enthalpy of combustion of some common fuels at 298 K and 101.3 kPa**

| Substance         | Formula                                      | State | $\Delta H_c$ (kJ mol <sup>-1</sup> ) |
|-------------------|--|-------|--------------------------------------|
| hydrogen          | $\text{H}_2$                                 | g     | -286                                 |
| carbon (graphite) | C  | s     | -394                                 |
| methane           | $\text{CH}_4$                                | g     | -889                                 |
| ethane            | $\text{C}_2\text{H}_6$                       | g     | -1557                                |
| propane           | $\text{C}_3\text{H}_8$                       | g     | -2217                                |
| butane            | $\text{C}_4\text{H}_{10}$                    | g     | -2874                                |
| pentane           | $\text{C}_5\text{H}_{12}$                    | l     | -3509                                |
| hexane            | $\text{C}_6\text{H}_{14}$                    | l     | -4158                                |
| octane            | $\text{C}_8\text{H}_{18}$                    | l     | -5464                                |
| ethene            | $\text{C}_2\text{H}_4$                       | g     | -1409                                |
| methanol          | $\text{CH}_3\text{OH}$                       | l     | -725                                 |
| ethanol           | $\text{C}_2\text{H}_5\text{OH}$              | l     | -1364                                |
| 1-propanol        | $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$ | l     | -2016                                |
| 2-propanol        | $\text{CH}_3\text{CHOHCH}_3$                 | l     | -2003                                |
| glucose           | $\text{C}_6\text{H}_{12}\text{O}_6$          | s     | -2816                                |



